

Title Of The Invention

**NUCLEIC ACID AND AMINO ACID SEQUENCES RELATING TO  
*ENTEROBACTER CLOACAE* FOR DIAGNOSTICS AND THERAPEUTICS**

5

Cross-Reference to Related Applications

This application is converted from U.S. provisional application Serial Number  
10 60/074,787, filed February 18, 1998 and U.S. provisional application Serial Number  
60/094,145 filed July 24, 1998.

Field Of The Invention

The invention relates to isolated nucleic acids and polypeptides derived from  
15 *Enterobacter cloacae* that are useful as molecular targets for diagnostics, prophylaxis  
and treatment of pathological conditions, as well as materials and methods for the  
diagnosis, prevention, and amelioration of pathological conditions resulting from  
bacterial infection.

20 Background Of The Invention

*Enterobacter cloacae* (*E. cloacae*) belongs to the bacterial family  
Enterobacteriaceae, whose diverse members are Gram-negative rods that are glucose  
fermenters and nitrate reducers. These organisms are found free-living in nature and  
as part of the indigenous flora of human and animals. They grow rapidly under  
25 aerobic and anaerobic conditions and are metabolically active, utilizing a variety of  
substrates. Most species are opportunistic pathogens (Kenneth Ryan,  
Enterobacteriaceae, Chap. 20, Medical Microbiology, An Introduction to Infectious  
Diseases, Second Edition, Editor, John C. Sherris, Elsevier, New York, 1990).

*E. cloacae* is an ornithine-positive, lysine-negative pathogen that can be  
30 associated with urinary tract and respiratory tract infections. The bacteria produces

endotoxins which as aerosols can penetrate into the lungs causing fever, coughing, difficulty in breathing and wheezing (Fairley, T. and Gislason, S., 1986-1997, Environmed Research Inc). *E. cloacae* is becoming progressively common in newborns in Neonatal Intensive Care Units (NICU) (Shi, Z.Y., et al, 1996, J. Clin. Microbiol. 34:2784-2790; Cordero, L., et al, 1997, Pediatr. Infect. Dis. J. 16:18-23; Acolet, D., et al, 1994, J. Hosp. Infect. 28:273-286). A study at Children's Hospital in Michigan showed a four-fold increase in *Enterobacter* in patients with bacteremia between 1989 and 1992. *E. cloacae* accounted for 74% of the isolates. Twenty eight percent of the infected children went into shock and six percent died (Andresen, J., et al, 1994, Pediatr. Infect. Dis. J. 13: 787-792). An outbreak of multidrug-resistant *E. cloacae* lasted for 4 months in the NICU in China (Shi, Z.Y., et al, 1996, J. Clin. Microbiol. 34:2784-2790). Outbreaks have also occurred in surgical wards (Burchard, K.W., et al, 1986, Surgery 100:857-862 ) and burn units (Markowitz, S. M., et al, 1983, J. Infect. Dis. 148:18-23). *E. cloacae* has also been shown to be the causative agent in a case of gas gangrene (Fata, F., et al, 1996, South Med. J. 89:1095-1096).

Epidemiology of *E. cloacae* is not completely understood, although studies of infection and colonization point to the endogenous flora of the patients. Molecular typing results of 141 strains of *E. cloacae* from broad geographic areas in the United States (from the National Surveillance Program: SCOPE) indicated that although clonal spread of a single strain was observed within a given institution most of the episodes of bacteremia were caused by strains unique to the individual patients. Therefore, selection of mutant subpopulations within each endogenous infection can be caused by drug exposure (Pfaller, M.A., 1997, Diagn. Microbiol. Infect. Dis. 28:211-219).

Antibiotic resistance is a major problem in the control of infectious diseases. Strains of *E. cloacae* resistant to broad-spectrum penicillins and beta-lactamase-stable

cephalosporins occurs at a frequency of  $10^7$  to  $10^6$  (Kadima, T.A. and Weiner, J.H., 1997, Antimicrobiol. Agents Chemother. 41:2177-2183; Lampe, M.F., et al, Antimicrob. Agents Chemother. 21:655-660; Lindberg, F., et al, Rev. Infect. Dis. 8 [Suppl 3]:S292-S304). Selected fluoroquinolones have often been successfully  
5 administered to patients with urinary tract infections; however, *E. cloacae* has become resistant to many of them (Deguchi, T., et al, 1997, Antimicrobiol. Agents Chemother. 41: 2544-2546). Some resistance has been attributed to plasmid-containing *E. cloacae* and some to the *E. cloacae* chromosome. In Holland, two different resistant strains of *E. cloacae* have been identified. The Amsterdam strain (resistant to ceftotaxim and  
10 piperacillin) exhibits depressed chromosomal Class 1 beta-lactamase, whereas the Rotterdam strain (resistant to cefuroxime) favors the spread of a plasmid encoding TEM-2 beta-lactamase (Namavar, F., 1997, BIO 99-53 99-606615 ). Resistant strains of *E. cloacae* developed within 6 days in nearly 50% of the *E. cloacae*-infected intensive care patients with pulmonary complications treated with cefotaxime (Fussle,  
15 et al., 1994, Clin. Investig. 72:1015-1019). While several antimicrobial agents retain potent activity against the highly resistant organisms (Pfaller, M.A., 1997, Diagn. Microbiol. Infect. Dis. 28:211-219), constant exposure to these agents may eventually result in resistance.

*E. cloacae* has been shown to be beneficial to plants in the control of diseases  
20 caused by bacteria (Bacon, C.W., et al., PCT publication WO 97/24433). As a biocontrol agent, *E. cloacae* coated onto cucumber seed has protected the seed from a lethal infection of the fungus *Pythium ultimum* (Nelson, E.B., et al, 1992, Can. J. Plant Pathol. 14:106-114). Nutritional mutants of *E. cloacae* were also protective and it has been suggested that mutant strains would be beneficial for an environmental  
25 containment strategy (Roberts, D.P., et al, 1994, Plant Science [Limerick], 10183-89).

## Summary Of The Invention

The present invention fulfills the need for diagnostic tools and therapeutics by providing bacterial-specific compositions and methods for detecting *Enterobacter* species including *E. cloacae*, as well as compositions and methods useful for treating and preventing *Enterobacter* infection, in particular, *E. cloacae* infection, in vertebrates including mammals.

The present invention encompasses isolated nucleic acids and polypeptides derived from *E. cloacae* that are useful as reagents for diagnosis of bacterial disease, components of effective antibacterial vaccines, and/or as targets for antibacterial drugs including anti-*E. cloacae* drugs. They can also be used to detect the presence of *E. cloacae* and other *Enterobacter* species in a sample; and in screening compounds for the ability to interfere with the *E. cloacae* life cycle or to inhibit *E. cloacae* infection. They also have use as biocontrol agents for plants.

In one aspect, the invention features compositions of nucleic acids corresponding to entire coding sequences of *E. cloacae* proteins, including surface or secreted proteins or parts thereof, nucleic acids capable of binding mRNA from *E. cloacae* proteins to block protein translation, and methods for producing *E. cloacae* proteins or parts thereof using peptide synthesis and recombinant DNA techniques. This invention also features antibodies and nucleic acids useful as probes to detect *E. cloacae* infection. In addition, vaccine compositions and methods for the protection or treatment of infection by *E. cloacae* are within the scope of this invention.

The nucleotide sequences provided in SEQ ID NO: 1 - SEQ ID NO: 5662, a fragment thereof, or a nucleotide sequence at least about 99.5% identical to a sequence contained within SEQ ID NO: 1 - SEQ ID NO: 5662 may be "provided" in a variety of medias to facilitate use thereof. As used herein, "provided" refers to a manufacture, other than an isolated nucleic acid molecule, which contains a nucleotide



sequence of the present invention, i.e., the nucleotide sequence provided in SEQ ID NO: 1 - SEQ ID NO: 5662, a fragment thereof, or a nucleotide sequence at least about 99.5% identical to a sequence contained within SEQ ID NO: 1 - SEQ ID NO: 5662. Uses for and methods for providing nucleotide sequences in a variety of media is well known in the art (see e.g., EPO Publication No. EP 0 756 006).

In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any media which can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage media, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A person skilled in the art can readily appreciate how any of the presently known computer readable media can be used to create a manufacture comprising computer readable media having recorded thereon a nucleotide sequence of the present invention.

As used herein, "recorded" refers to a process for storing information on computer readable media. A person skilled in the art can readily adopt any of the presently known methods for recording information on computer readable media to generate manufactures comprising the nucleotide sequence information of the present invention.

A variety of data storage structures are available to a person skilled in the art for creating a computer readable media having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable media. The sequence

information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. A person skilled in the art can readily adapt any  
5 number of data processor structuring formats (e.g. text file or database) in order to obtain computer readable media having recorded thereon the nucleotide sequence information of the present invention.

By providing the nucleotide sequence of SEQ ID NO: 1 - SEQ ID NO: 5662, a fragment thereof, or a nucleotide sequence at least about 99.5% identical to SEQ ID  
10 NO: 1 - SEQ ID NO: 5662 in computer readable form, a person skilled in the art can routinely access the coding sequence information for a variety of purposes. Computer software is publicly available which allows a person skilled in the art to access sequence information provided in a computer readable media. Examples of such computer software include programs of the "Staden Package", "DNA Star",  
15 "MacVector", GCG "Wisconsin Package" (Genetics Computer Group, Madison, WI) and "NCBI Toolbox" (National Center For Biotechnology Information). Suitable programs are described, for example, in Martin J. Bishop, ed., *Guide to Human Genome Computing*, 2d Edition, Academic Press, San Diego, CA. (1998); and Leonard F. Peruski, Jr., and Anne Harwood Peruski, *The Internet and the New*  
20 *Biology: Tools for Genomic and Molecular Research*, American Society for Microbiology, Washington, D.C. (1997).

Computer algorithms enable the identification of *E. cloacae* open reading frames (ORFs) within SEQ ID NO: 1 - SEQ ID NO: 5662 which contain homology to ORFs or proteins from other organisms. Examples of such similarity-search  
25 algorithms include the BLAST [Altschul et al., J. Mol. Biol. 215:403-410 (1990)] and Smith-Waterman [Smith and Waterman (1981) *Advances in Applied Mathematics*,

2:482-489] search algorithms. Suitable search algorithms are described, for example, in Martin J. Bishop, ed., *Guide to Human Genome Computing*, 2d Edition, Academic Press, San Diego, CA. (1998); and Leonard F. Peruski, Jr., and Anne Harwood Peruski, *The Internet and the New Biology: Tools for Genomic and Molecular Research*, American Society for Microbiology, Washington, D.C. (1997). Such algorithms are utilized on computer systems as exemplified below. The ORFs so identified represent protein encoding fragments within the *E. cloacae* genome and are useful in producing commercially important proteins such as enzymes used in fermentation reactions and in the production of commercially useful metabolites.

10       The present invention further provides systems, particularly computer-based systems, which contain the sequence information described herein. Such systems are designed to identify commercially important fragments of the *E. cloacae* genome. As used herein, "a computer-based system" refers to the hardware means, software means, and data storage means used to analyze the nucleotide sequence information of the  
15       present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means, and data storage means. A person skilled in the art can readily appreciate that any one of the currently available computer-based systems is suitable for use in the present invention. The computer-based systems of the present invention comprise a  
20       data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory which can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the  
25       nucleotide sequence information of the present invention.

As used herein, "search means" refers to one or more programs which are implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of the *E. cloacae* genome which are similar to, or "match", a particular target sequence or target motif. A variety of known algorithms are known in the art and have been disclosed publicly, and a variety of commercially available software for conducting homology-based similarity searches are available and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to, FASTA (GCG Wisconsin Package), Bic\_SW (Compugen Bioccelerator), BLASTN2, BLASTP2, BLASTX2 (NCBI) and Motifs (GCG). Suitable software programs are described, for example, in Martin J. Bishop, ed., *Guide to Human Genome Computing*, 2d Edition, Academic Press, San Diego, CA. (1998); and Leonard F. Peruski, Jr., and Anne Harwood Peruski, *The Internet and the New Biology: Tools for Genomic and Molecular Research*, American Society for Microbiology, Washington, D.C. (1997). A person skilled in the art can readily recognize that any one of the available algorithms or implementing software packages for conducting homology searches can be adapted for use in the present computer-based systems.

As used herein, a "target sequence" can be any DNA or amino acid sequence of six or more nucleotides or two or more amino acids. A person skilled in the art can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. The most preferred sequence length of a target sequence is from about 10 to 100 amino acids or from about 30 to 300 nucleotide residues. However, it is well recognized that many genes are longer than 500 amino acids, or 1.5 kb in length, and that commercially important fragments

of the *E. cloacae* genome, such as sequence fragments involved in gene expression and protein processing, will often be shorter than 30 nucleotides.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequence(s) are  
5 chosen based on a specific functional domain or three-dimensional configuration which is formed upon the folding of the target polypeptide. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzymatic active sites, membrane-spanning regions, and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin  
10 structures and inducible expression elements (protein binding sequences).

A variety of structural formats for the input and output means can be used to input and output the information in the computer-based systems of the present invention. A preferred format for an output means ranks fragments of the *E. cloacae* genome possessing varying degrees of homology to the target sequence or target  
15 motif. Such presentation provides a person skilled in the art with a ranking of sequences which contain various amounts of the target sequence or target motif and identifies the degree of homology contained in the identified fragment.

A variety of comparing means can be used to compare a target sequence or target motif with the data storage means to identify sequence fragments of the *E.*  
20 *cloacae* genome. In the present examples, implementing software which implement the BLASTP2 and bic\_SW algorithms (Altschul et al., J Mol. Biol. 215:403-410 (1990); Compugen Biocellator) was used to identify open reading frames within the *E. cloacae* genome. A person skilled in the art can readily recognize that any one of the publicly available homology search programs can be used as the search means for the  
25 computer-based systems of the present invention. Suitable programs are described, for example, in Martin J. Bishop, ed., *Guide to Human Genome Computing*, 2d Edition,

Academic Press, San Diego, CA. (1998); and Leonard F. Peruski, Jr., and Anne Harwood Peruski, *The Internet and the New Biology: Tools for Genomic and Molecular Research*, American Society for Microbiology, Washington, D.C. (1997).

The invention features *E. cloacae* polypeptides, preferably a substantially pure  
5 preparation of an *E. cloacae* polypeptide, or a recombinant *E. cloacae* polypeptide. In  
preferred embodiments: the polypeptide has biological activity; the polypeptide has an  
amino acid sequence at least about 60%, 70%, 80%, 90%, 95%, 98%, or 99%  
identical to an amino acid sequence of the invention contained in the Sequence  
Listing, preferably it has about 65% sequence identity with an amino acid sequence of  
10 the invention contained in the Sequence Listing, and most preferably it has about 92%  
to about 99% sequence identity with an amino acid sequence of the invention  
contained in the Sequence Listing; the polypeptide has an amino acid sequence  
essentially the same as an amino acid sequence of the invention contained in the  
Sequence Listing; the polypeptide is at least about 5, 10, 20, 50, 100, or 150 amino  
15 acid residues in length; the polypeptide includes at least about 5, preferably at least  
about 10, more preferably at least about 20, still more preferably at least about 50,  
100, or 150 contiguous amino acid residues of the invention contained in the  
Sequence Listing. In yet another preferred embodiment, the amino acid sequence  
which differs in sequence identity by about 7% to about 8% from the *E. cloacae*  
20 amino acid sequences of the invention contained in the Sequence Listing is also  
encompassed by the invention.

In preferred embodiments: the *E. cloacae* polypeptide is encoded by a nucleic  
acid of the invention contained in the Sequence Listing, or by a nucleic acid having at  
least about 60%, 70%, 80%, 90%, 95%, 98%, or 99% homology with a nucleic acid  
25 of the invention contained in the Sequence Listing.

In a preferred embodiment, the subject *E. cloacae* polypeptide differs in amino acid sequence at about 1, 2, 3, 5, 10 or more residues from a sequence of the invention contained in the Sequence Listing. The differences, however, are such that the *E. cloacae* polypeptide exhibits an *E. cloacae* biological activity, e.g., the *E. cloacae* polypeptide retains a biological activity of a naturally occurring *E. cloacae* enzyme.

In preferred embodiments, the polypeptide includes all or a fragment of an amino acid sequence of the invention contained in the Sequence Listing; fused, in reading frame, to additional amino acid residues, preferably to residues encoded by genomic DNA 5' or 3' to the genomic DNA which encodes a sequence of the invention contained in the Sequence Listing.

In yet other preferred embodiments, the *E. cloacae* polypeptide is a recombinant fusion protein having a first *E. cloacae* polypeptide portion and a second polypeptide portion, e.g., a second polypeptide portion having an amino acid sequence unrelated to *E. cloacae*. The second polypeptide portion can be, e.g., any of glutathione-S-transferase, a DNA binding domain, or a polymerase activating domain. In preferred embodiment the fusion protein can be used in a two-hybrid assay.

Polypeptides of the invention include those which arise as a result of alternative transcription events, alternative RNA splicing events, and alternative translational and postranslational events.

In a preferred embodiment, the encoded *E. cloacae* polypeptide differs (e.g., by amino acid substitution, addition or deletion of at least one amino acid residue) in amino acid sequence at about 1, 2, 3, 5, 10 or more residues, from a sequence of the invention contained in the Sequence Listing. The differences, however, are such that the *E. cloacae* encoded polypeptide exhibits an *E. cloacae* biological activity, e.g., the

encoded *E. cloacae* enzyme retains a biological activity of a naturally occurring *E. cloacae*.

In preferred embodiments, the encoded polypeptide includes all or a fragment of an amino acid sequence of the invention contained in the Sequence Listing; fused, in reading frame, to additional amino acid residues, preferably to residues encoded by genomic DNA 5' or 3' to the genomic DNA which encodes a sequence of the invention contained in the Sequence Listing.

The *E. cloacae* strain, 15842, from which genomic sequences have been sequenced, has been deposited on August 22, 1997, in the American Type Culture Collection and assigned the ATCC designation # 202023.

Included in the invention are: allelic variations; natural mutants; induced mutants; proteins encoded by DNA that hybridize under high or low stringency conditions to a nucleic acid which encodes a polypeptide of the invention contained in the Sequence Listing (for definitions of high and low stringency see Current Protocols in Molecular Biology, John Wiley & Sons, New York, 1989, 6.3.1 - 6.3.6, hereby incorporated by reference); and, polypeptides specifically bound by antisera to *E. cloacae* polypeptides, especially by antisera to an active site or binding domain of *E. cloacae* polypeptide. The invention also includes fragments, preferably biologically active fragments. These and other polypeptides are also referred to herein as *E. cloacae* polypeptide analogs or variants.

The invention further provides nucleic acids, e.g., RNA or DNA, encoding a polypeptide of the invention. This includes double stranded nucleic acids as well as coding and antisense single strands.

In preferred embodiments, the subject *E. cloacae* nucleic acid will include a transcriptional regulatory sequence, e.g., at least one of a transcriptional promoter or transcriptional enhancer sequence, operably linked to the *E. cloacae* gene sequence,



e.g., to render the *E. cloacae* gene sequence suitable for expression in a recombinant host cell.

In yet a further preferred embodiment, the nucleic acid which encodes an *E. cloacae* polypeptide of the invention, hybridizes under stringent conditions to a  
5 nucleic acid probe corresponding to at least about 8 consecutive nucleotides of the invention contained in the Sequence Listing; more preferably to at least about 12 consecutive nucleotides of the invention contained in the Sequence Listing; still more preferably to at least about 20 consecutive nucleotides of the invention contained in the Sequence Listing; most preferably to at least about 40 consecutive nucleotides of  
10 the invention contained in the Sequence Listing.

In another aspect, the invention provides a substantially pure nucleic acid having a nucleotide sequence which encodes an *E. cloacae* polypeptide. In preferred embodiments: the encoded polypeptide has biological activity; the encoded polypeptide has an amino acid sequence at least about 60%, 70%, 80%, 90%, 95%,  
15 98% or 99% homologous to an amino acid sequence of the invention contained in the Sequence Listing; the encoded polypeptide has an amino acid sequence essentially the same as an amino acid sequence of the invention contained in the Sequence Listing; the encoded polypeptide is at least about 5, 10, 20, 50, 100, or 150 amino acids in length; the encoded polypeptide comprises at least about 5, preferably at least about  
20 10, more preferably at least about 20, still more preferably at least about 50, 100, or 150 contiguous amino acids of the invention contained in the Sequence Listing.

In another aspect, the invention encompasses: a vector including a nucleic acid which encodes an *E. cloacae* polypeptide or an *E. cloacae* polypeptide variant as described herein; a host cell transfected with the vector; and a method of producing a  
25 recombinant *E. cloacae* polypeptide or *E. cloacae* polypeptide variant; including

culturing the cell, e.g., in a cell culture medium, and isolating an *E. cloacae* or *E. cloacae* polypeptide variant, e.g., from the cell or from the cell culture medium.

One embodiment of the invention is directed to substantially isolated nucleic acids. Nucleic acids of the invention include sequences comprising at least about 8  
5 nucleotides in length, more preferably at least about 12 nucleotides in length, even more preferably at least about 15-20 nucleotides in length, that correspond to a subsequence of any one of SEQ ID NO: 1 - SEQ ID NO: 5662 or complements thereof. Alternatively, the nucleic acids comprise sequences contained within any ORF (open reading frame), including a complete protein-coding sequence, of which  
10 any of SEQ ID NO: 1 - SEQ ID NO: 5662 forms a part. The invention encompasses sequence-conservative variants and function-conservative variants of these sequences. The nucleic acids may be DNA, RNA, DNA/RNA duplexes, protein-nucleic acid (PNA), or derivatives thereof.

In another aspect, the invention features a purified recombinant nucleic acid  
15 having at least about 50%, 60%, 70%, 80%, 90%, 95%, 98%, or 99% homology with a sequence of the invention contained in the Sequence Listing

The invention also encompasses recombinant DNA (including DNA cloning and expression vectors) comprising these *E. cloacae*-derived sequences; host cells comprising such DNA, including fungal, bacterial, yeast, plant, insect, and mammalian  
20 host cells; and methods for producing expression products comprising RNA and polypeptides encoded by the *E. cloacae* sequences. These methods are carried out by incubating a host cell comprising an *E. cloacae*-derived nucleic acid sequence under conditions in which the sequence is expressed. The host cell may be native or recombinant. The polypeptides can be obtained by (a) harvesting the incubated cells  
25 to produce a cell fraction and a medium fraction; and (b) recovering the *E. cloacae*

polypeptide from the cell fraction, the medium fraction, or both. The polypeptides can also be made by *in vitro* translation.

In another aspect, the invention features nucleic acids capable of binding mRNA of *E. cloacae*. Such nucleic acid is capable of acting as antisense nucleic acid to control the translation of mRNA of *E. cloacae*. A further aspect features a nucleic acid which is capable of binding specifically to an *E. cloacae* nucleic acid. These nucleic acids are also referred to herein as complements and have utility as probes and as capture reagents.

In another aspect, the invention features an expression system comprising an open reading frame corresponding to *E. cloacae* nucleic acid. The nucleic acid further comprises a control sequence compatible with an intended host. The expression system is useful for making polypeptides corresponding to *E. cloacae* nucleic acid.

In another aspect, the invention encompasses: a vector including a nucleic acid which encodes an *E. cloacae* polypeptide or an *E. cloacae* polypeptide variant as described herein; a host cell transfected with the vector; and a method of producing a recombinant *E. cloacae* polypeptide or *E. cloacae* polypeptide variant; including culturing the cell, e.g., in a cell culture medium, and isolating the *E. cloacae* or *E. cloacae* polypeptide variant, e.g., from the cell or from the cell culture medium.

In yet another embodiment of the invention encompasses reagents for detecting bacterial infection, including *E. cloacae* infection, which comprise at least one *E. cloacae*-derived nucleic acid defined by any one of SEQ ID NO: 1 - SEQ ID NO: 5662, or sequence-conservative or function-conservative variants thereof.

Alternatively, the diagnostic reagents comprise nucleotide sequences that are contained within any open reading frames (ORFs), including preferably complete protein-coding sequences, contained within any of SEQ ID NO: 1 - SEQ ID NO: 5662, or polypeptide sequences contained within any of SEQ ID NO: 5663 - SEQ ID NO:

11324, or polypeptides of which any of the above sequences forms a part, or antibodies directed against any of the above peptide sequences or function-conservative variants and/or fragments thereof.

The invention further provides antibodies, preferably monoclonal antibodies, which specifically bind to the polypeptides of the invention. Methods are also provided for producing antibodies in a host animal. The methods of the invention comprise immunizing an animal with at least one *E. cloacae*-derived immunogenic component, wherein the immunogenic component comprises one or more of the polypeptides encoded by any one of SEQ ID NO: 1 - SEQ ID NO: 5662 or sequence-conservative or function-conservative variants thereof; or polypeptides that are contained within any ORFs, including complete protein-coding sequences, of which any<sup>a</sup> of SEQ ID NO: 1 - SEQ ID NO: 5662 forms a part; or polypeptide sequences contained within any of SEQ ID NO: 5663 - SEQ ID NO: 11324; or polypeptides of which any of SEQ ID NO: 5663 - SEQ ID NO: 11324 forms a part. Host animals include any warm blooded animal, including without limitation mammals and birds. Such antibodies have utility as reagents for immunoassays to evaluate the abundance and distribution of *E. cloacae*-specific antigens.

In yet another aspect, the invention provides diagnostic methods for detecting *E. cloacae* antigenic components or anti-*E. cloacae* antibodies in a sample. *E. cloacae* antigenic components may be detected by known processes, including but not limited to detection by a process comprising: (i) contacting a sample suspected to contain a bacterial antigenic component with a bacterial-specific antibody, under conditions in which a stable antigen-antibody complex can form between the antibody and bacterial antigenic components in the sample; and (ii) detecting any antigen-antibody complex formed in step (i), wherein detection of an antigen-antibody complex indicates the presence of at least one bacterial antigenic component in the

sample. In different embodiments of this method, the antibodies used are directed against a sequence encoded by any of SEQ ID NO: 1 - SEQ ID NO: 5662 or sequence-conservative or function-conservative variants thereof, or against a polypeptide sequence contained in any of SEQ ID NO: 5663 - SEQ ID NO: 11324 or  
5 function-conservative variants thereof.

In yet another aspect, the invention provides a method for detecting antibacterial-specific antibodies in a sample, which comprises: (i) contacting a sample suspected to contain antibacterial-specific antibodies with an *E. cloacae* antigenic component, under conditions in which a stable antigen-antibody complex can form  
10 between the *E. cloacae* antigenic component and antibacterial antibodies in the sample; and (ii) detecting any antigen-antibody complex formed in step (i), wherein detection of an antigen-antibody complex indicates the presence of antibacterial antibodies in the sample. In different embodiments of this method, the antigenic component is encoded by a sequence contained in any of SEQ ID NO: 1 - SEQ ID  
15 NO: 5662 or sequence-conservative and function-conservative variants thereof, or is a polypeptide sequence contained in any of SEQ ID NO: 5663 - SEQ ID NO: 11324 or function-conservative variants thereof.

In another aspect, the invention features a method of generating vaccines for immunizing an individual against *E. cloacae*. The method includes: immunizing a  
20 subject with an *E. cloacae* polypeptide, e.g., a surface or secreted polypeptide, or a combination of such peptides or active portion(s) thereof, and a pharmaceutically acceptable carrier. Such vaccines have therapeutic and prophylactic utilities.

In another aspect, the invention features a method of evaluating a compound, e.g., a polypeptide, e.g., a fragment of a host cell polypeptide, for the ability to bind  
25 an *E. cloacae* polypeptide. The method includes contacting the compound to be evaluated with an *E. cloacae* polypeptide and determining if the compound binds or

otherwise interacts with the *E. cloacae* polypeptide. Compounds which bind or otherwise interact with *E. cloacae* polypeptides are candidates as modulators, including activators and inhibitors, of the bacterial life cycle. These assays can be performed *in vitro* or *in vivo*.

- 5           In another aspect, the invention features a method of evaluating a compound, e.g., a polypeptide, e.g., a fragment of a host cell polypeptide, for the ability to bind an *E. cloacae* nucleic acid, e.g., DNA or RNA. The method includes contacting the compound to be evaluated with an *E. cloacae* nucleic acid and determining if the compound binds or otherwise interacts with the *E. cloacae* nucleic acid. Compounds  
10   which bind *E. cloacae* are candidates as modulators, including activators and inhibitors, of the bacterial life cycle. These assays can be performed *in vitro* or *in vivo*.

A particularly preferred embodiment of the invention is directed to a method of screening test compounds for anti-bacterial activity, which method comprises: selecting as a target a bacterial specific sequence, which sequence is essential to the  
15   viability of a bacterial species; contacting a test compound with said target sequence; and selecting those test compounds which bind to said target sequence as potential anti-bacterial candidates. In one embodiment, the target sequence selected is specific to a single species, or even a single strain, such as, for example, the strain *E. cloacae* 15842. In a second embodiment, the target sequence is common to at least two  
20   species of bacteria. In a third embodiment, the target sequence is common to a family of bacteria. The target sequence may be a nucleic acid sequence or a polypeptide sequence. Methods employing sequences common to more than one species of microorganism may be used to screen candidates for broad spectrum anti-bacterial activity.

- 25           The invention also provides methods for preventing or treating disease caused by certain bacteria, including *E. cloacae*, which are carried out by administering to an

animal in need of such treatment, in particular a warm-blooded vertebrate, including but not limited to birds and mammals, a compound that specifically inhibits or interferes with the function of a bacterial polypeptide or nucleic acid. In a particularly preferred embodiment, the mammal to be treated is human.

5

## DETAILED DESCRIPTION OF THE INVENTION

The sequences of the present invention include the specific nucleic acid and amino acid sequences set forth in the Sequence Listing that forms a part of the present specification, and which are designated SEQ ID NO:1 - SEQ ID NO: 11324. Use of  
10 the terms "SEQ ID NO: 1 - SEQ ID NO: 5662 ", " SEQ ID NO: 5663 - SEQ ID NO: 11324, "the sequences depicted in Table 2", etc., is intended, for convenience, to refer to each individual SEQ ID NO *individually*, and is not intended to refer to the genus of these sequences unless such reference would be indicated. In other words, it is a shorthand for listing all of these sequences individually. The invention encompasses  
15 each sequence individually, as well as any combination thereof.

### Definitions

"Nucleic acid" or "polynucleotide" as used herein refers to purine- and pyrimidine-containing polymers of any length, either polyribonucleotides or  
20 polydeoxyribonucleotides or mixed polyribo-polydeoxyribo nucleotides. This includes single- and double-stranded molecules, i.e., DNA-DNA, DNA-RNA and RNA-RNA hybrids, as well as "protein nucleic acids" (PNA) formed by conjugating bases to an amino acid backbone. This also includes nucleic acids containing modified bases.

A nucleic acid or polypeptide sequence that is "derived from" a designated  
25 sequence refers to a sequence that corresponds to a region of the designated sequence. For nucleic acid sequences, this encompasses sequences that are homologous or

complementary to the sequence, as well as "sequence-conservative variants" and "function-conservative variants." For polypeptide sequences, this encompasses "function-conservative variants." Sequence-conservative variants are those in which a change of one or more nucleotides in a given codon position results in no alteration in the amino acid encoded at that position. Function-conservative variants are those in which a given amino acid residue in a polypeptide has been changed without altering the overall conformation and function of the native polypeptide, including, but not limited to, replacement of an amino acid with one having similar physico-chemical properties (such as, for example, acidic, basic, hydrophobic, and the like). "Function-conservative" variants also include any polypeptides that have the ability to elicit antibodies specific to a designated polypeptide.

An "*E. cloacae*-derived" nucleic acid or polypeptide sequence may or may not be present in other bacterial species, and may or may not be present in all *E. cloacae* strains. This term is intended to refer to the source from which the sequence was originally isolated. Thus, an *E. cloacae*-derived polypeptide, as used herein, may be used, e.g., as a target to screen for a broad spectrum antibacterial agent, to search for homologous proteins in other species of bacteria or in eukaryotic organisms such as bacteria humans, etc.

A purified or isolated polypeptide or a substantially pure preparation of a polypeptide are used interchangeably herein and, as used herein, mean a polypeptide that has been separated from other proteins, lipids, and nucleic acids with which it naturally occurs. Preferably, the polypeptide is also separated from substances, e.g., antibodies or gel matrix, e.g., polyacrylamide, which are used to purify it. Preferably, the polypeptide constitutes at least about 10, 20, 50 70, 80 or 95% dry weight of the purified preparation. Preferably, the preparation contains sufficient polypeptide to allow protein sequencing; at least about 1, 10, or preferably 100 mg of polypeptide.



A purified preparation of cells refers to, in the case of plant or animal cells, an *in vitro* preparation of cells and not an entire intact plant or animal. In the case of cultured cells or microbial cells, it consists of a preparation of at least about 10%, more preferably at least about 50%, of the subject cells.

5        A purified or isolated or a substantially pure nucleic acid, e.g., a substantially pure DNA, (are terms used interchangeably herein) is a nucleic acid which is one or both of the following: not immediately contiguous with both of the coding sequences with which it is immediately contiguous (i.e., one at the 5' end and one at the 3' end) in the naturally-occurring genome of the organism from which the nucleic acid is  
10        derived; or which is substantially free of a nucleic acid with which it occurs in the organism from which the nucleic acid is derived. The term includes, for example, a recombinant DNA which is incorporated into a vector, e.g., into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule (e.g., a cDNA or a genomic DNA fragment  
15        produced by PCR or restriction endonuclease treatment) independent of other DNA sequences. Substantially pure DNA also includes a recombinant DNA which is part of a hybrid gene encoding additional *E. cloacae* DNA sequence.

A "contig" as used herein is a nucleic acid representing a continuous stretch of genomic sequence of an organism.

20        An "open reading frame", also referred to herein as ORF, is a region of nucleic acid which encodes a polypeptide. This region may represent a portion of a coding sequence or a total sequence and can be determined from a stop to stop codon or from a start to stop codon.

As used herein, a "coding sequence" is a nucleic acid which is transcribed into  
25        messenger RNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are

determined by a translation start codon at the five prime terminus and a translation stop code at the three prime terminus. A coding sequence can include but is not limited to messenger RNA, synthetic DNA, and recombinant nucleic acid sequences.

A "complement" of a nucleic acid as used herein refers to an anti-parallel or antisense sequence that participates in Watson-Crick base-pairing with the original sequence.

A "gene product" is a protein or structural RNA which is specifically encoded by a gene.

As used herein, the term "probe" refers to a nucleic acid, peptide or other chemical entity which specifically binds to a molecule of interest. Probes are often associated with or capable of associating with a label. A label is a chemical moiety capable of detection. Typical labels comprise dyes, radioisotopes, luminescent and chemiluminescent moieties, fluorophores, enzymes, precipitating agents, amplification sequences, and the like. Similarly, a nucleic acid, peptide or other chemical entity which specifically binds to a molecule of interest and immobilizes such molecule is referred herein as a "capture ligand". Capture ligands are typically associated with or capable of associating with a support such as nitro-cellulose, glass, nylon membranes, beads, particles and the like. The specificity of hybridization is dependent on conditions such as the base pair composition of the nucleotides, and the temperature and salt concentration of the reaction. These conditions are readily discernable to one of ordinary skill in the art using routine experimentation.

"Homologous" refers to the sequence similarity or sequence identity between two polypeptides or between two nucleic acid molecules. When a position in both of the two compared sequences is occupied by the same base or amino acid monomer subunit, e.g., if a position in each of two DNA molecules is occupied by adenine, then the molecules are homologous at that position. The percent of homology between two

sequences is a function of the number of matching or homologous positions shared by the two sequences divided by the number of positions compared x 100. For example, if 6 of 10 of the positions in two sequences are matched or homologous then the two sequences are 60% homologous. By way of example, the DNA sequences ATTGCC-  
5 and TATGGC share 50% homology. Generally, a comparison is made when two sequences are aligned to give maximum homology.

Nucleic acids are hybridizable to each other when at least one strand of a nucleic acid can anneal to the other nucleic acid under defined stringency conditions. Stringency of hybridization is determined by: (a) the temperature at which  
10 hybridization and/or washing is performed; and (b) the ionic strength and polarity of the hybridization and washing solutions. Hybridization requires that the two nucleic acids contain complementary sequences; depending on the stringency of hybridization, however, mismatches may be tolerated. Typically, hybridization of two sequences at high stringency (such as, for example, in a solution of 0.5X SSC, at 65° C) requires  
15 that the sequences be essentially completely homologous. Conditions of intermediate stringency (such as, for example, 2X SSC at 65 ° C) and low stringency (such as, for example 2X SSC at 55° C) require correspondingly less overall complementarity between the hybridizing sequences. (1X SSC is 0.15 M NaCl, 0.015 M Na citrate).

The terms peptides, proteins, and polypeptides are used interchangeably herein.  
20 As used herein, the term "surface protein" refers to all surface accessible proteins, e.g. inner and outer membrane proteins, proteins adhering to the cell wall, and secreted proteins.

A polypeptide has *E. cloacae* biological activity if it has one, two or preferably more of the following properties: (1) if when expressed in the course of an *E.*  
25 *cloacae* infection, it can promote, or mediate the attachment of *E. cloacae* to a cell; (2) it has an enzymatic activity, structural or regulatory function characteristic of an *E.*

*cloacae* protein; (3) the gene which encodes it can rescue a lethal mutation in an *E. cloacae* gene. A polypeptide has biological activity if it is an antagonist, agonist, or super-agonist of a polypeptide having one of the above-listed properties.

A biologically active fragment or analog is one having an *in vivo* or *in vitro* activity which is characteristic of the *E. cloacae* polypeptides of the invention contained in the Sequence Listing, or of other naturally occurring *E. cloacae* polypeptides, e.g., one or more of the biological activities described herein. Especially preferred are fragments which exist *in vivo*, e.g., fragments which arise from post transcriptional processing or which arise from translation of alternatively spliced RNA's. Fragments include those expressed in native or endogenous cells as well as those made in expression systems, e.g., in CHO (Chinese Hamster Ovary) cells. Because peptides such as *E. cloacae* polypeptides often exhibit a range of physiological properties and because such properties may be attributable to different portions of the molecule, a useful *E. cloacae* fragment or *E. cloacae* analog is one which exhibits a biological activity in any biological assay for *E. cloacae* activity. The fragment or analog possesses about 10%, preferably about 40%, more preferably about 60%, 70%, 80% or 90% or greater of the activity of *E. cloacae*, in any *in vivo* or *in vitro* assay.

Analogous can differ from naturally occurring *E. cloacae* polypeptides in amino acid sequence or in ways that do not involve sequence, or both. Non-sequence modifications include changes in acetylation, methylation, phosphorylation, carboxylation, or glycosylation. Preferred analogs include *E. cloacae* polypeptides (or biologically active fragments thereof) whose sequences differ from the wild-type sequence by one or more conservative amino acid substitutions or by one or more non-conservative amino acid substitutions, deletions, or insertions which do not substantially diminish the biological activity of the *E. cloacae* polypeptide.

Conservative substitutions typically include the substitution of one amino acid for another with similar characteristics, e.g., substitutions within the following groups: valine, glycine; glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Other conservative substitutions can be made in view of the table below.

TABLE 1

CONSERVATIVE AMINO ACID REPLACEMENTS

	For Amino Acid	Code	Replace with any of
	Alanine	A	D-Ala, Gly, beta-Ala, L-Cys, D-Cys
10	Arginine	R	D-Arg, Lys, D-Lys, homo-Arg, D-homo-Arg, Met,
	Asparagine	N	D-Asn, Asp, D-Asp, Glu, D-Glu, Gln, D-Gln
	Aspartic Acid	R	D-Asp, D-Asn, Asn, Glu, D-Glu, Gln, D-Gln
	Cysteine	F	D-Cys, S-Me-Cys, Met, D-Met, Thr, D-Thr
	Glutamine	Q	D-Gln, Asn, D-Asn, Glu, D-Glu, Asp, D-Asp
15	Glutamic Acid	L	D-Glu, D-Asp, Asp, Asn, D-Asn, Gln, D-Gln
	Glycine	L	Ala, D-Ala, Pro, D-Pro, $\beta$ -Ala, Acp
	Isoleucine	L	D-Ile, Val, D-Val, Leu, D-Leu, Met, D-Met
	Leucine	L	D-Leu, Val, D-Val, Leu, D-Leu, Met, D-Met
	Lysine	R	D-Lys, Arg, D-Arg, homo-Arg, D-homo-Arg, Met,
20	Methionine	M	D-Met, S-Me-Cys, Ile, D-Ile, Leu, D-Leu, Val, D-
	Phenylalanine	F	D-Phe, Tyr, D-Thr, L-Dopa, His, D-His, Trp, D-
	Proline	P	D-Pro, L-I-thioazolidine-4-carboxylic acid, D-or L-
	Serine	S	D-Ser, Thr, D-Thr, allo-Thr, Met, D-Met, Met(O),
	Threonine	T	D-Thr, Ser, D-Ser, allo-Thr, Met, D-Met, Met(O),
25	Tyrosine	Y	D-Tyr, Phe, D-Phe, L-Dopa, His, D-His
	Valine	V	D-Val, Leu, D-Leu, Ile, D-Ile, Met, D-Met

Other analogs within the invention are those with modifications which increase peptide stability; such analogs may contain, for example, one or more non-peptide bonds (which replace the peptide bonds) in the peptide sequence. Also included are: analogs that include residues other than naturally occurring L-amino acids, e.g., D-amino acids or non-naturally occurring or synthetic amino acids, e.g.,  $\beta$  or  $\gamma$  amino acids; and cyclic analogs.

As used herein, the term "fragment", as applied to an *E. cloacae* analog, will ordinarily be at least about 20 residues, more typically at least about 40 residues, preferably at least about 60 residues in length. Fragments of *E. cloacae* polypeptides can be generated by methods known to those skilled in the art. The ability of an *Enterobacter* fragment to exhibit a biological activity of *E. cloacae* polypeptide can be assessed by methods known to those skilled in the art as described herein. Also included are *E. cloacae* polypeptides containing residues that are not required for biological activity of the peptide or that result from alternative mRNA splicing or alternative protein processing events.

An "immunogenic component" as used herein is a moiety, such as an *E. cloacae* polypeptide, analog or fragment thereof, that is capable of eliciting a humoral and/or cellular immune response in a host animal.

An "antigenic component" as used herein is a moiety, such as an *E. cloacae* polypeptide, analog or fragment thereof, that is capable of binding to a specific antibody with sufficiently high affinity to form a detectable antigen-antibody complex.

The term "antibody" as used herein is intended to include fragments thereof which are specifically reactive with *E. cloacae* polypeptides.

As used herein, the term "cell-specific promoter" means a DNA sequence that serves as a promoter, i.e., regulates expression of a selected DNA sequence operably linked to the promoter, and which effects expression of the selected DNA sequence in

specific cells of a tissue. The term also covers so-called "leaky" promoters, which regulate expression of a selected DNA primarily in one tissue, but cause expression in other tissues as well.

Misexpression, as used herein, refers to a non-wild type pattern of gene expression. It includes: expression at non-wild type levels, i.e., over or under expression; a pattern of expression that differs from wild type in terms of the time or stage at which the gene is expressed, e.g., increased or decreased expression (as compared with wild type) at a predetermined developmental period or stage; a pattern of expression that differs from wild type in terms of increased expression (as compared with wild type) in a predetermined cell type or tissue type; a pattern of expression that differs from wild type in terms of the splicing size, amino acid sequence, post-translational modification, or biological activity of the expressed polypeptide; a pattern of expression that differs from wild type in terms of the effect of an environmental stimulus or extracellular stimulus on expression of the gene, e.g., a pattern of increased or decreased expression (as compared with wild type) in the presence of an increase or decrease in the strength of the stimulus.

As used herein, "host cells" and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refers to cells which can become or have been used as recipients for a recombinant vector or other transfer DNA, and include the progeny of the original cell which has been transfected. It is understood by individuals skilled in the art that the progeny of a single parental cell may not necessarily be completely identical in genomic or total DNA compliment to the original parent, due to accident or deliberate mutation.

As used herein, the term "control sequence" refers to a nucleic acid having a base sequence which is recognized by the host organism to effect the expression of encoded sequences to which they are ligated. The nature of such control sequences

differs depending upon the host organism; in prokaryotes, such control sequences generally include a promoter, ribosomal binding site, terminators, and in some cases operators; in eukaryotes, generally such control sequences include promoters, terminators and in some instances, enhancers. The term control sequence is intended  
5 to include at a minimum, all components whose presence is necessary for expression, and may also include additional components whose presence is advantageous, for example, leader sequences.

As used herein, the term "operably linked" refers to sequences joined or ligated to function in their intended manner. For example, a control sequence is operably  
10 linked to coding sequence by ligation in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequence and host cell.

The "metabolism" of a substance, as used herein, means any aspect of the expression, function, action, or regulation of the substance. The metabolism of a  
15 substance includes modifications, e.g., covalent or non-covalent modifications of the substance. The metabolism of a substance includes modifications, e.g., covalent or non-covalent modification, the substance induces in other substances. The metabolism of a substance also includes changes in the distribution of the substance. The metabolism of a substance includes changes the substance induces in the distribution  
20 of other substances.

A "sample" as used herein refers to a biological sample, such as, for example, tissue or fluid isolated from an individual (including without limitation plasma, serum, cerebrospinal fluid, lymph, tears, saliva and tissue sections) or from *in vitro* cell culture constituents, as well as samples from the environment.

25 Technical and scientific terms used herein have the meanings commonly understood by one of ordinary skill in the art to which the present invention pertains,



unless otherwise defined. Reference is made herein to various methodologies known to those of skill in the art. Publications and other materials setting forth such known methodologies to which reference is made are incorporated herein by reference in their entireties as though set forth in full. The practice of the invention will employ, unless

5 otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. *See e.g.*, Sambrook, Fritsch, and Maniatis, *Molecular Cloning; Laboratory Manual* 2nd ed. (1989); *DNA Cloning*, Volumes I and II (D.N Glover ed. 1985); *Oligonucleotide Synthesis* (M.J. Gait ed,

10 1984); *Nucleic Acid Hybridization* (B.D. Hames & S.J. Higgins eds. 1984); the series, *Methods in Enzymology* (Academic Press, Inc.), particularly Vol. 154 and Vol. 155 (Wu and Grossman, eds.); *PCR-A Practical Approach* (McPherson, Quirke, and Taylor, eds., 1991); *Immunology*, 2d Edition, 1989, Roitt *et al.*, C.V. Mosby Company, and New York; *Advanced Immunology*, 2d Edition, 1991, Male *et al.*,

15 Grower Medical Publishing, New York.; *DNA Cloning: A Practical Approach*, Volumes I and II, 1985 (D.N. Glover ed.); *Oligonucleotide Synthesis*, 1984, (M.L. Gait ed); *Transcription and Translation*, 1984 (Hames and Higgins eds.); *Animal Cell Culture*, 1986 (R.I. Freshney ed.); *Immobilized Cells and Enzymes*, 1986 (IRL Press); Perbal, 1984, *A Practical Guide to Molecular Cloning; Gene Transfer Vectors for*

20 *Mammalian Cells*, 1987 (J. H. Miller and M. P. Calos eds., Cold Spring Harbor Laboratory); Martin J. Bishop, ed., *Guide to Human Genome Computing*, 2d Edition, Academic Press, San Diego, CA. (1998); and Leonard F. Peruski, Jr., and Anne Harwood Peruski, *The Internet and the New Biology: Tools for Genomic and Molecular Research*, American Society for Microbiology, Washington, D.C. (1997).

25 Any suitable materials and/or methods known to those of skill can be utilized in carrying out the present invention; however, preferred materials and/or methods are

described. Materials, reagents and the like to which reference is made in the following description and examples are obtainable from commercial sources, unless otherwise noted.

## 5 *E. cloacae* Genomic Sequence

This invention provides nucleotide sequences of the genome of *E. cloacae* which thus comprises a DNA sequence library of *E. cloacae* genomic DNA. The detailed description that follows provides nucleotide sequences of *E. cloacae*, and also  
10 describes how the sequences were obtained and how ORFs and protein-coding sequences were identified. Also described are compositions and methods of using the disclosed *E. cloacae* sequences in methods including diagnostic and therapeutic applications. Furthermore, the library can be used as a database for identification and comparison of medically important sequences in this and other strains of *E. cloacae*.

15 To determine the genomic sequence of *E. cloacae*, DNA from strain 15842 of *E. cloacae* was isolated after Zymolyase digestion, sodium dodecyl sulfate lysis, potassium acetate precipitation, phenol:chloroform extraction and ethanol precipitation (Soll, D.R., T. Srikantha and S.R. Lockhart: Characterizing Developmentally Regulated Genes in *E. cloacae*. In Microbial Genome Methods. K.W. Adolph, editor.  
20 CRC Press. New York. p 17-37.). DNA was sheared hydrodynamically using an HPLC (Oefner, et. al., 1996) to an insert size of 2000-3000 bp. After size fractionation by gel electrophoresis the fragments were blunt-ended, ligated to adapter oligonucleotides and cloned into the pGTC (Thomann) vector to construct a "shotgun" subclone library.

DNA sequencing was achieved using established ABI sequencing methods on ABI377 automated DNA sequencers. The cloning and sequencing procedures are described in more detail in the Exemplification.

Individual sequence reads were assembled using PHRAP (P. Green, Abstracts of DOE Human Genome Program Contractor-Grantee Workshop V, Jan. 1996, p.157). The average contig length was about 3-4 kb.

All subsequent steps were based on sequencing by ABI377 automated DNA sequencing methods. The cloning and sequencing procedures are described in more detail in the Exemplification.

10 A variety of approaches may be used to order the contigs so as to obtain a continuous sequence representing the entire *E. cloacae* genome. Synthetic oligonucleotides are designed that are complementary to sequences at the end of each contig. These oligonucleotides may be hybridized to libraries of *E. cloacae* genomic DNA in, for example, lambda phage vectors or plasmid vectors to identify clones that  
15 contain sequences corresponding to the junctional regions between individual contigs. Such clones are then used to isolate template DNA and the same oligonucleotides are used as primers in polymerase chain reaction (PCR) to amplify junctional fragments, the nucleotide sequence of which is then determined.

The *E. cloacae* sequences were analyzed for the presence of open reading  
20 frames (ORFs) comprising at least 180 nucleotides. As a result of the analysis of ORFs based on stop-to-stop codon reads, it should be understood that these ORFs may not correspond to the ORF of a naturally-occurring *E. cloacae* polypeptide. These ORFs may contain start codons which indicate the initiation of protein synthesis of a naturally-occurring *E. cloacae* polypeptide. Such start codons within the ORFs  
25 provided herein were identified by those of ordinary skill in the relevant art, and the resulting ORF and the encoded *E. cloacae* polypeptide is within the scope of this

invention. For example, within the ORFs a codon such as AUG or GUG (encoding methionine or valine) which is part of the initiation signal for protein synthesis were identified and the portion of an ORF corresponding to a naturally-occurring *E.*

*cloacae* polypeptide was recognized. The predicted coding regions were defined by  
5 evaluating the coding potential of such sequences with the program GENEMARK™  
(Borodovsky and McIninch, 1993, *Comp. .* 17:123).

Each predicted ORF amino acid sequence was compared with all sequences found in current GENBANK, SWISS-PROT, and PIR databases using the BLAST algorithm. BLAST identifies local alignments occurring by chance between the ORF  
10 sequence and the sequence in the databank (Altschul et al., 1990, *L Mol. Biol.* 215:403-410). Homologous ORFs (probabilities less than  $10^{-5}$  by chance) and ORF's that are probably non-homologous (probabilities greater than  $10^{-5}$  by chance) but have good codon usage were identified. Both homologous, sequences and non-homologous sequences with good codon usage, are likely to encode proteins and are encompassed  
15 by the invention.

#### *E. cloacae* Nucleic Acids

The present invention provides a library of *E. cloacae*-derived nucleic acid  
20 sequences. The libraries provide probes, primers, and markers which are used as markers in epidemiological studies. The present invention also provides a library of *E. cloacae*-derived nucleic acid sequences which comprise or encode targets for therapeutic drugs.

The nucleic acids of this invention may be obtained directly from the DNA of  
25 the above referenced *E. cloacae* strain by using the polymerase chain reaction (PCR). See "*PCR, A Practical Approach*" (McPherson, Quirke, and Taylor, eds., IRL Press,

Oxford, UK, 1991) for details about the PCR. High fidelity PCRs used to ensure a faithful DNA copy prior to expression. In addition, the authenticity of amplified products is verified by conventional sequencing methods. Clones carrying the desired sequences described in this invention may also be obtained by screening the libraries  
5 by means of the PCR or by hybridization of synthetic oligonucleotide probes to filter lifts of the library colonies or plaques as known in the art (see, e.g., Sambrook et al., *Molecular Cloning, A Laboratory Manual* 2nd edition, 1989, Cold Spring Harbor Press, NY).

It is also possible to obtain nucleic acids encoding *E. cloacae* polypeptides  
10 from a cDNA library in accordance with protocols herein described. A cDNA encoding an *E. cloacae* polypeptide can be obtained by isolating total mRNA from an appropriate strain. Double stranded cDNAs can then be prepared from the total mRNA. Subsequently, the cDNAs can be inserted into a suitable plasmid or viral (e.g., bacteriophage) vector using any one of a number of known techniques. Genes  
15 encoding *E. cloacae* polypeptides can also be cloned using established polymerase chain reaction techniques in accordance with the nucleotide sequence information provided by the invention. The nucleic acids of the invention can be DNA or RNA. Preferred nucleic acids of the invention are contained in the Sequence Listing.

The nucleic acids of the invention can also be chemically synthesized using  
20 standard techniques. Various methods of chemically synthesizing polydeoxynucleotides are known, including solid-phase synthesis which, like peptide synthesis, has been fully automated in commercially available DNA synthesizers (See e.g., Itakura et al. U.S. Patent No. 4,598,049; Caruthers et al. U.S. Patent No. 4,458,066; and Itakura U.S. Patent Nos. 4,401,796 and 4,373,071, incorporated by  
25 reference herein).

In another example, DNA can be chemically synthesized using, e.g., the phosphoramidite solid support method of Matteucci *et al.*, 1981, *J. Am. Chem. Soc.* 103:3185, the method of Yoo *et al.*, 1989, *J. Biol. Chem.* 264:17078, or other well known methods. This can be done by sequentially linking a series of oligonucleotide  
5 cassettes comprising pairs of synthetic oligonucleotides, as described below.

Nucleic acids isolated or synthesized in accordance with features of the present invention are useful, by way of example, without limitation, as probes, primers, capture ligands, antisense genes and for developing expression systems for the synthesis of proteins and peptides corresponding to such sequences. As probes,  
10 primers, capture ligands and antisense agents, the nucleic acid normally consists of all or part (approximately twenty or more nucleotides for specificity as well as the ability to form stable hybridization products) of the nucleic acids of the invention contained in the Sequence Listing. These uses are described in further detail below.

### 15       Probes

A nucleic acid isolated or synthesized in accordance with the sequence of the invention contained in the Sequence Listing can be used as a probe to specifically detect *E. cloacae*. With the sequence information set forth in the present application, sequences of twenty or more nucleotides are identified which provide the desired  
20 inclusivity and exclusivity with respect to *E. cloacae*, and extraneous nucleic acids likely to be encountered during hybridization conditions. More preferably, the sequence will comprise at least about twenty to thirty nucleotides to convey stability to the hybridization product formed between the probe and the intended target molecules.

25       Sequences larger than 1000 nucleotides in length are difficult to synthesize but can be generated by recombinant DNA techniques. Individuals skilled in the art will

readily recognize that the nucleic acids, for use as probes, can be provided with a label to facilitate detection of a hybridization product.

Nucleic acid isolated and synthesized in accordance with the sequence of the invention contained in the Sequence Listing can also be useful as probes to detect  
5 homologous regions (especially homologous genes) of other *Enterobacter* species using appropriate stringency hybridization conditions as described herein.

### Capture Ligand

For use as a capture ligand, the nucleic acid selected in the manner described  
10 above with respect to probes, can be readily associated with a support. The manner in which nucleic acid is associated with supports is well known. Nucleic acid having twenty or more nucleotides in a sequence of the invention contained in the Sequence Listing have utility to separate *E. cloacae* nucleic acid from one strain from the nucleic acid of other another strain as well as from other organisms. Nucleic acid  
15 having twenty or more nucleotides in a sequence of the invention contained in the Sequence Listing can also have utility to separate other *Enterobacter* species from each other and from other organisms. Preferably, the sequence will comprise at least about twenty nucleotides to convey stability to the hybridization product formed between the probe and the intended target molecules. Sequences larger than 1000  
20 nucleotides in length are difficult to synthesize but can be generated by recombinant DNA techniques.

### Primers

Nucleic acid isolated or synthesized in accordance with the sequences  
25 described herein have utility as primers for the amplification of *E. cloacae* nucleic acid. These nucleic acids may also have utility as primers for the amplification of

nucleic acids in other *Enterobacter* species. With respect to polymerase chain reaction (PCR) techniques, nucleic acid sequences of  $\geq 10$ -15 nucleotides of the invention contained in the Sequence Listing have utility in conjunction with suitable enzymes and reagents to create copies of *E. cloacae* nucleic acid. More preferably, the sequence will comprise twenty or more nucleotides to convey stability to the hybridization product formed between the primer and the intended target molecules. Binding conditions of primers greater than 100 nucleotides are more difficult to control to obtain specificity. High fidelity PCR can be used to ensure a faithful DNA copy prior to expression. In addition, amplified products can be checked by conventional sequencing methods.

The copies can be used in diagnostic assays to detect specific sequences, including genes from *E. cloacae* and/or other *Enterobacter* species. The copies can also be incorporated into cloning and expression vectors to generate polypeptides corresponding to the nucleic acid synthesized by PCR, as is described in greater detail herein.

The nucleic acids of the present invention find use as templates for the recombinant production of *E. cloacae*-derived peptides or polypeptides

#### Antisense

Nucleic acid or nucleic acid-hybridizing derivatives isolated or synthesized in accordance with the sequences described herein have utility as antisense agents to prevent the expression of *E. cloacae* genes. These sequences also have utility as antisense agents to prevent expression of genes of other *Enterobacter* species.

In one embodiment, nucleic acid or derivatives corresponding to *E. cloacae* nucleic acids is loaded into a suitable carrier such as a liposome or bacteriophage for introduction into bacterial cells. For example, a nucleic acid having twenty or more



nucleotides is capable of binding to bacteria nucleic acid or bacteria messenger RNA. Preferably, the antisense nucleic acid is comprised of 20 or more nucleotides to provide necessary stability of a hybridization product of non-naturally occurring nucleic acid and bacterial nucleic acid and/or bacterial messenger RNA. Nucleic acid  
5 having a sequence greater than 1000 nucleotides in length is difficult to synthesize but can be generated by recombinant DNA techniques. Methods for loading antisense nucleic acid in liposomes is known in the art as exemplified by U.S. Patent 4,241,046 issued December 23, 1980 to Papahadjopoulos et al.

The present invention encompasses isolated polypeptides and nucleic acids  
10 derived from *E. cloacae* that are useful as reagents for diagnosis of bacterial infection, components of effective anti-bacterial vaccines, and/or as targets for anti-bacterial drugs, including anti-*E. cloacae* drugs.

#### Expression of *E. cloacae* Nucleic Acids

15

Table 2, which is appended herewith and which forms part of the present specification, provides a list of open reading frames (ORFs) in both strands and a putative identification of the particular function of a polypeptide which is encoded by each ORF, based on the homology match (determined by the BLAST algorithm) of  
20 the predicted polypeptide with known proteins encoded by ORFs in other organisms. An ORF is a region of nucleic acid which encodes a polypeptide. This region may represent a portion of a coding sequence or a total sequence and was determined from stop to stop codons. The first column contains a designation for the contig from which each ORF was identified (numbered arbitrarily). Each contig represents a  
25 continuous stretch of the genomic sequence of the organism. The second column lists the ORF designation. The third and fourth columns list the SEQ ID numbers for the

nucleic acid and amino acid sequences corresponding to each ORF, respectively. The fifth and sixth columns list the length of the nucleic acid ORF and the length of the amino acid ORF, respectively. The nucleotide sequence corresponding to each ORF begins at the first nucleotide immediately following a stop codon and ends at the

5 nucleotide immediately preceding the next downstream stop codon in the same reading frame. It will be recognized by one skilled in the art that the natural translation initiation sites will correspond to ATG, GTG, or TTG codons located within the ORFs. The natural initiation sites depend not only on the sequence of a start codon but also on the context of the DNA sequence adjacent to the start codon. Usually, a

10 recognizable ribosome binding site is found within 20 nucleotides upstream from the initiation codon. In some cases where genes are translationally coupled and coordinately expressed together in "operons", ribosome binding sites are not present, but the initiation codon of a downstream gene may occur very close to, or overlap, the stop codon of the an upstream gene in the same operon. The correct start codons can

15 be generally identified without undue experimentation because only a few codons need be tested. It is recognized that the translational machinery in bacteria initiates all polypeptide chains with the amino acid methionine, regardless of the sequence of the start codon. In some cases, polypeptides are post-translationally modified, resulting in an N-terminal amino acid other than methionine *in vivo*. The seventh and eighth

20 columns provide metrics for assessing the likelihood of the homology match (determined by the BLASTP2 algorithm), as is known in the art, to the genes indicated in the eleventh column when the designated ORF was compared against a non-redundant comprehensive protein database. Specifically, the seventh column represents the "Blast Score" for the match (a higher score is a better match), and the

25 eighth column represents the "P-value" for the match (the probability that such a match can have occurred by chance; the lower the value, the more likely the match is

valid). If a BLASTP2 score of less than 46 was obtained, no value is reported in the table the "P-value". Column nine provides the name of the organism that was identified as having the closest homology match. The tenth column provides, where available, either a public database accession number or our own sequence name. The  
5 eleventh column provides, where available, the Swissprot accession number (SP),(SP), the locus name (LN), the Organism (OR), Source of variant (SR), E.C. number (EC), the gene name (GN), the product name (PN), the Function Description (FN), Left End (LE), Right End (RE), Coding Direction (DI), and the description (DE) or notes (NT) for each ORF. Information that is not preceded by a code designation  
10 in the eleventh column represents a description of the ORF. This information allows one of ordinary skill in the art to determine a potential use for each identified coding sequence and, as a result, allows to use the polypeptides of the present invention for commercial and industrial purposes.

Using the information provided in SEQ ID NO: 1 - SEQ ID NO: 5662, SEQ  
15 ID NO: 5663 - SEQ ID NO: 11324 and in Table 2 together with routine cloning and sequencing methods, one of ordinary skill in the art will be able to clone and sequence all the nucleic acid fragments of interest including open reading frames (ORFs) encoding a large variety of proteins of *E. cloacae*.

Nucleic acid isolated or synthesized in accordance with the sequences  
20 described herein have utility to generate polypeptides. The nucleic acid of the invention exemplified in SEQ ID NO: 1 - SEQ ID NO: 5662 and in Table 2 or fragments of said nucleic acid encoding active portions of *E. cloacae* polypeptides can be cloned into suitable vectors or used to isolate nucleic acid. The isolated nucleic acid is combined with suitable DNA linkers and cloned into a suitable vector.

25 The function of a specific gene or operon can be ascertained by expression in a bacterial strain under conditions where the activity of the gene product(s) specified by

the gene or operon in question can be specifically measured. Alternatively, a gene product may be produced in large quantities in an expressing strain for use as an antigen, an industrial reagent, for structural studies, etc. This expression can be accomplished in a mutant strain which lacks the activity of the gene to be tested, or in  
5 a strain that does not produce the same gene product(s). This includes, but is not limited to, Eucaryotic species such as the yeast *Saccharomyces cerevisiae*, *Methanobacterium* strains or other Archaea, and Eubacteria such as *E. coli*, *B. Subtilis*, *S. Aureus*, *S. Pneumonia* or *Pseudomonas putida*. In some cases the expression host will utilize the natural *E. cloacae* promoter whereas in others, it will  
10 be necessary to drive the gene with a promoter sequence derived from the expressing organism (e.g., an *E. coli* beta-galactosidase promoter for expression in *E. coli*).

To express a gene product using the natural *E. cloacae* promoter, a procedure such as the following can be used. A restriction fragment containing the gene of interest, together with its associated natural promoter element and regulatory  
15 sequences (identified using the DNA sequence data) is cloned into an appropriate recombinant plasmid containing an origin of replication that functions in the host organism and an appropriate selectable marker. This can be accomplished by a number of procedures known to those skilled in the art. It is most preferably done by cutting the plasmid and the fragment to be cloned with the same restriction enzyme to  
20 produce compatible ends that can be ligated to join the two pieces together. The recombinant plasmid is introduced into the host organism by, for example, electroporation and cells containing the recombinant plasmid are identified by selection for the marker on the plasmid. Expression of the desired gene product is detected using an assay specific for that gene product.

25 In the case of a gene that requires a different promoter, the body of the gene (coding sequence) is specifically excised and cloned into an appropriate expression

plasmid. This subcloning can be done by several methods, but is most easily accomplished by PCR amplification of a specific fragment and ligation into an expression plasmid after treating the PCR product with a restriction enzyme or exonuclease to create suitable ends for cloning.

- 5           A suitable host cell for expression of a gene can be any procaryotic or eucaryotic cell. Suitable methods for transforming host cells can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory Press (1989)), and other laboratory textbooks.

- For example, a host cell transfected with a nucleic acid vector directing  
10       expression of a nucleotide sequence encoding an *E. cloacae* polypeptide can be cultured under appropriate conditions to allow expression of the polypeptide to occur. Suitable media for cell culture are well known in the art. Polypeptides of the invention can be isolated from cell culture medium, host cells, or both using techniques known in the art for purifying proteins including ion-exchange  
15       chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, and immunoaffinity purification with antibodies specific for such polypeptides. Additionally, in many situations, polypeptides can be produced by chemical cleavage of a native protein (e.g., tryptic digestion) and the cleavage products can then be purified by standard techniques.

- 20           In the case of membrane bound proteins, these can be isolated from a host cell by contacting a membrane-associated protein fraction with a detergent forming a solubilized complex, where the membrane-associated protein is no longer entirely embedded in the membrane fraction and is solubilized at least to an extent which allows it to be chromatographically isolated from the membrane fraction.  
25       Chromatographic techniques which can be used in the final purification step are

known in the art and include hydrophobic interaction, lectin affinity, ion exchange, dye affinity and immunoaffinity.

One strategy to maximize recombinant *E. cloacae* peptide expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically  
5 cleave the recombinant protein (Gottesman, S., Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, California (1990) 119-128).

Another strategy would be to alter the nucleic acid encoding an *E. cloacae* peptide to be inserted into an expression vector so that the individual codons for each amino acid would be those preferentially utilized in highly expressed *E. coli* proteins (Wada et al.,  
10 (1992) *Nuc. Acids Res.* 20:2111-2118). Such alteration of nucleic acids of the invention can be carried out by standard DNA synthesis techniques.

The nucleic acids of the invention can also be chemically synthesized using standard techniques. Various methods of chemically synthesizing polydeoxynucleotides are known, including solid-phase synthesis which, like peptide  
15 synthesis, has been fully automated in commercially available DNA synthesizers (See, e.g., Itakura et al. U.S. Patent No. 4,598,049; Caruthers et al. U.S. Patent No. 4,458,066; and Itakura U.S. Patent Nos. 4,401,796 and 4,373,071, incorporated by reference herein).

The present invention provides a library of *E. cloacae*-derived nucleic acid  
20 sequences. The libraries provide probes, primers, and markers which can be used as markers in epidemiological studies. The present invention also provides a library of *E. cloacae*-derived nucleic acid sequences which comprise or encode targets for therapeutic drugs.

Nucleic acids comprising any of the sequences disclosed herein or sub-  
25 sequences thereof can be prepared by standard methods using the nucleic acid sequence information provided in SEQ ID NO: 1 - SEQ ID NO: 5662. For example,

DNA can be chemically synthesized using, e.g., the phosphoramidite solid support method of Matteucci *et al.*, 1981, *J. Am. Chem. Soc.* 103:3185, the method of Yoo *et al.*, 1989, *J. Biol. Chem.* 764:17078, or other well known methods. This can be done by sequentially linking a series of oligonucleotide cassettes comprising pairs of synthetic oligonucleotides, as described below.

Of course, due to the degeneracy of the genetic code, many different nucleotide sequences can encode polypeptides having the amino acid sequences defined by SEQ ID NO: 5663 - SEQ ID NO: 11324 or sub-sequences thereof. The codons can be selected for optimal expression in prokaryotic or eukaryotic systems.

Such degenerate variants are also encompassed by this invention.

Insertion of nucleic acids (typically DNAs) encoding the polypeptides of the invention into a vector is easily accomplished when the termini of both the DNAs and the vector comprise compatible restriction sites. If this cannot be done, it may be necessary to modify the termini of the DNAs and/or vector by digesting back single-stranded DNA overhangs generated by restriction endonuclease cleavage to produce blunt ends, or to achieve the same result by filling in the single-stranded termini with an appropriate DNA polymerase.

Alternatively, any site desired may be produced, e.g., by ligating nucleotide sequences (linkers) onto the termini. Such linkers may comprise specific oligonucleotide sequences that define desired restriction sites. Restriction sites can also be generated by the use of the polymerase chain reaction (PCR). See, e.g., Saiki *et al.*, 1988, *Science* 239:48. The cleaved vector and the DNA fragments may also be modified if required by homopolymeric tailing.

The nucleic acids of the invention may be isolated directly from cells. Alternatively, the polymerase chain reaction (PCR) method can be used to produce the nucleic acids of the invention, using either chemically synthesized strands or genomic

material as templates. Primers used for PCR can be synthesized using the sequence information provided herein and can further be designed to introduce appropriate new restriction sites, if desirable, to facilitate incorporation into a given vector for recombinant expression.

- 5           The nucleic acids of the present invention may be flanked by natural *E. cloacae* regulatory sequences, or may be associated with heterologous sequences, including promoters, enhancers, response elements, signal sequences, polyadenylation sequences, introns, 5'- and 3'- noncoding regions, and the like. The nucleic acids may also be modified by many means known in the art. Non-limiting examples of such
- 10       modifications include methylation, "caps", substitution of one or more of the naturally occurring nucleotides with an analog; internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoroamidates, carbamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.). Nucleic acids may contain one or more
- 15       additional covalently linked moieties, such as, for example, proteins (e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), intercalators (e.g., acridine, psoralen, etc.), chelators (e.g., metals, radioactive metals, iron, oxidative metals, etc.), and alkylators. PNAs are also included. The nucleic acid may be derivatized by formation of a methyl or ethyl phosphotriester or an alkyl phosphoramidate linkage.
- 20       Furthermore, the nucleic acid sequences of the present invention may also be modified with a label capable of providing a detectable signal, either directly or indirectly. Exemplary labels include radioisotopes, fluorescent molecules, biotin, and the like.

          The invention also provides nucleic acid vectors comprising the disclosed *E. cloacae*-derived sequences or derivatives or fragments thereof. A large number of

25       vectors, including plasmid and bacterial vectors, have been described for replication



and/or expression in a variety of eukaryotic and prokaryotic hosts, and may be used for cloning or protein expression.

The encoded *E. cloacae* polypeptides may be expressed by using many known vectors, such as pUC plasmids, pET plasmids (Novagen, Inc., Madison, WI), or  
5 pRSET or pREP (Invitrogen, San Diego, CA), and many appropriate host cells, using methods disclosed or cited herein or otherwise known to those skilled in the relevant art. The particular choice of vector/host is not critical to the practice of the invention.

Recombinant cloning vectors will often include one or more replication systems for cloning or expression, one or more markers for selection in the host, e.g.  
10 antibiotic resistance, and one or more expression cassettes. The inserted *E. cloacae* coding sequences may be synthesized by standard methods, isolated from natural sources, or prepared as hybrids, etc. Ligation of the *E. cloacae* coding sequences to transcriptional regulatory elements and/or to other amino acid coding sequences may be achieved by known methods. Suitable host cells may be  
15 transformed/transfected/infected as appropriate by any suitable method including electroporation, CaCl<sub>2</sub> mediated DNA uptake, bacterial infection, microinjection, microprojectile, or other established methods.

Appropriate host cells include bacteria, archebacteria, fungi, especially yeast, and plant and animal cells, especially mammalian cells. Of particular interest are *E.*  
20 *cloacae*, *E. coli*, *B. Subtilis*, *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, *Schizosaccharomyces pombe*, SF9 cells, C129 cells, 293 cells, *Neurospora*, and CHO cells, COS cells, HeLa cells, and immortalized mammalian myeloid and lymphoid cell lines. Preferred replication systems include M13, ColE1, SV40, baculovirus, lambda, adenovirus, and the like. A large number of transcription initiation and termination  
25 regulatory regions have been isolated and shown to be effective in the transcription and translation of heterologous proteins in the various hosts. Examples of these

regions, methods of isolation, manner of manipulation, etc. are known in the art. Under appropriate expression conditions, host cells can be used as a source of recombinantly produced *E. cloacae*-derived peptides and polypeptides.

Advantageously, vectors may also include a transcription regulatory element (i.e., a promoter) operably linked to the *E. cloacae* portion. The promoter may optionally contain operator portions and/or ribosome binding sites. Non-limiting examples of bacterial promoters compatible with *E. coli* include: b-lactamase (penicillinase) promoter; lactose promoter; tryptophan (trp) promoter; araBAD (arabinose) operon promoter; lambda-derived P<sub>l</sub> promoter and N gene ribosome binding site; and the hybrid tac promoter derived from sequences of the trp and lac UV5 promoters. Non-limiting examples of yeast promoters include 3-phosphoglycerate kinase promoter, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) promoter, galactokinase (GAL1) promoter, galactose epimerase promoter, and alcohol dehydrogenase (ADH) promoter. Suitable promoters for mammalian cells include without limitation viral promoters such as that from Simian Virus 40 (SV40), Rous sarcoma virus (RSV), adenovirus (ADV), and bovine papilloma virus (BPV). Mammalian cells may also require terminator sequences, polyA addition sequences and enhancer sequences to increase expression. Sequences which cause amplification of the gene may also be desirable. Furthermore, sequences that facilitate secretion of the recombinant product from cells, including, but not limited to, bacteria, yeast, and animal cells, such as secretory signal sequences and/or prohormone pro region sequences, may also be included. These sequences are well described in the art.

Nucleic acids encoding wild-type or variant *E. cloacae*-derived polypeptides may also be introduced into cells by recombination events. For example, such a sequence can be introduced into a cell, and thereby effect homologous recombination at the site of an endogenous gene or a sequence with substantial identity to the gene.

Other recombination-based methods such as nonhomologous recombinations or deletion of endogenous genes by homologous recombination may also be used.

The nucleic acids of the present invention find use as templates for the recombinant production of *E. cloacae*-derived peptides or polypeptides.

5

#### Identification and Use of *E. cloacae* Nucleic Acid Sequences

The disclosed *E. cloacae* polypeptide and nucleic acid sequences, or other sequences that are contained within ORFs, including complete protein-coding  
10 sequences, of which any of the disclosed *E. cloacae*-specific sequences forms a part, are useful as target components for diagnosis and/or treatment of *E. cloacae*- caused infection

It will be understood that the sequence of an entire protein-coding sequence of which each disclosed nucleic acid sequence forms a part can be isolated and identified  
15 based on each disclosed sequence. This can be achieved, for example, by using an isolated nucleic acid encoding the disclosed sequence, or fragments thereof, to prime a sequencing reaction with genomic *E. cloacae* DNA as template; this is followed by sequencing the amplified product. The isolated nucleic acid encoding the disclosed sequence, or fragments thereof, can also be hybridized to *E. cloacae* genomic libraries  
20 to identify clones containing additional complete segments of the protein-coding sequence of which the shorter sequence forms a part. Then, the entire protein-coding sequence, or fragments thereof, or nucleic acids encoding all or part of the sequence, or sequence-conservative or function-conservative variants thereof, may be employed in practicing the present invention.

25 Preferred sequences are those that are useful in diagnostic and/or therapeutic applications. Diagnostic applications include without limitation nucleic-acid-based and

antibody-based methods for detecting bacterial infection. Therapeutic applications include without limitation vaccines, passive immunotherapy, and drug treatments directed against gene products that are both unique to bacteria and essential for growth and/or replication of bacteria.

5

Identification of Nucleic Acids Encoding Vaccine Components and Targets for Agents Effective Against *E. cloacae*

The disclosed *E. cloacae* genome sequence includes segments that direct the  
10 synthesis of ribonucleic acids and polypeptides, as well as origins of replication, promoters, other types of regulatory sequences, and intergenic nucleic acids. The invention encompasses nucleic acids encoding immunogenic components of vaccines and targets for agents effective against *E. cloacae*. Identification of said immunogenic components involved in the determination of the function of the disclosed sequences,  
15 which can be achieved using a variety of approaches. Non-limiting examples of these approaches are described briefly below.

Homology to known sequences:

Computer-assisted comparison of the disclosed *E. cloacae* sequences with  
20 previously reported sequences present in publicly available databases is useful for identifying functional *E. cloacae* nucleic acid and polypeptide sequences. It will be understood that protein-coding sequences, for example, may be compared as a whole, and that a high degree of sequence homology between two proteins (such as, for example, >80-90%) at the amino acid level indicates that the two proteins also possess  
25 some degree of functional homology, such as, for example, among enzymes involved in metabolism, DNA synthesis, or cell wall synthesis, and proteins involved in

transport, cell division, etc. In addition, many structural features of particular protein classes have been identified and correlate with specific consensus sequences, such as, for example, binding domains for nucleotides, DNA, metal ions, and other small molecules; sites for covalent modifications such as phosphorylation, acylation, and the like; sites of protein:protein interactions, etc. These consensus sequences may be quite short and thus may represent only a fraction of the entire protein-coding sequence. Identification of such a feature in an *E. cloacae* sequence is therefore useful in determining the function of the encoded protein and identifying useful targets of antibacterial drugs.

Of particular relevance to the present invention are structural features that are common to secretory, transmembrane, and surface proteins, including secretion signal peptides and hydrophobic transmembrane domains. *E. cloacae* proteins identified as containing putative signal sequences and/or transmembrane domains are useful as immunogenic components of vaccines.

Targets for therapeutic drugs according to the invention include, but are not limited to, polypeptides of the invention, whether unique to *E. cloacae* or not, that are essential for growth and/or viability of *E. cloacae* under at least one growth condition. Polypeptides essential for growth and/or viability can be determined by examining the effect of deleting and/or disrupting the genes, i.e., by so-called gene "knockout".

Alternatively, genetic footprinting can be used (Smith *et al.*, 1995, *Proc. Natl. Acad. Sci. USA* 92:5479-6433; Published International Application WO 94/26933; U.S. Patent No. 5,612,180). Still other methods for assessing essentiality includes the ability to isolate conditional lethal mutations in the specific gene (e.g., temperature sensitive mutations). Other useful targets for therapeutic drugs, which include polypeptides that are not essential for growth or viability *per se* but lead to loss of viability of the cell, can be used to target therapeutic agents to cells.

### Strain-specific sequences:

Because of the evolutionary relationship between different *E. cloacae* strains, it is believed that the presently disclosed *E. cloacae* sequences are useful for identifying, and/or discriminating between, previously known and new *E. cloacae* strains. It is  
5 believed that other *E. cloacae* strains will exhibit at least about 70% sequence homology with the presently disclosed sequence. Systematic and routine analyses of DNA sequences derived from samples containing *E. cloacae* strains, and comparison with the present sequence allows for the identification of sequences that can be used to discriminate between strains, as well as those that are common to all *E. cloacae*  
10 strains. In one embodiment, the invention provides nucleic acids, including probes, and peptide and polypeptide sequences that discriminate between different strains of *E. cloacae*. Strain-specific components can also be identified functionally by their ability to elicit or react with antibodies that selectively recognize one or more *E. cloacae* strains.

15 In another embodiment, the invention provides nucleic acids, including probes, and peptide and polypeptide sequences that are common to all *E. cloacae* strains but are not found in other bacterial species.

### *E. cloacae* Polypeptides

20

This invention encompasses isolated *E. cloacae* polypeptides encoded by the disclosed *E. cloacae* genomic sequences, including the polypeptides of the invention contained in the Sequence Listing. Polypeptides of the invention are preferably at least about 5 amino acid residues in length. Using the DNA sequence information  
25 provided herein, the amino acid sequences of the polypeptides encompassed by the invention can be deduced using methods well-known in the art. It will be understood

that the sequence of an entire nucleic acid encoding an *E. cloacae* polypeptide can be isolated and identified based on an ORF that encodes only a fragment of the cognate protein-coding region. This can be achieved, for example, by using the isolated nucleic acid encoding the ORF, or fragments thereof, to prime a polymerase chain  
5 reaction with genomic *E. cloacae* DNA as template; this is followed by sequencing the amplified product.

The polypeptides of the present invention, including function-conservative variants of the disclosed ORFs, may be isolated from wild-type or mutant *E. cloacae* cells, or from heterologous organisms or cells (including, but not limited to, bacteria,  
10 fungi, insect, plant, and mammalian cells) including *E. cloacae* into which an *E. cloacae*-derived protein-coding sequence has been introduced and expressed. Furthermore, the polypeptides may be part of recombinant fusion proteins.

*E. cloacae* polypeptides of the invention can be chemically synthesized using commercially automated procedures such as those referenced herein , including,  
15 without limitation, exclusive solid phase synthesis, partial solid phase methods, fragment condensation or classical solution synthesis. The polypeptides are preferably prepared by solid phase peptide synthesis as described by Merrifield, 1963, *J. Am. Chem. Soc.* 85:2149. The synthesis is carried out with amino acids that are protected at the alpha-amino terminus. Trifunctional amino acids with labile side-chains are  
20 also protected with suitable groups to prevent undesired chemical reactions from occurring during the assembly of the polypeptides. The alpha-amino protecting group is selectively removed to allow subsequent reaction to take place at the amino-terminus. The conditions for the removal of the alpha-amino protecting group do not remove the side-chain protecting groups.

25 Methods for polypeptide purification are well-known in the art, including, without limitation, preparative disc-gel electrophoresis, isoelectric focusing, HPLC,

reversed-phase HPLC, gel filtration, ion exchange and partition chromatography, and countercurrent distribution. For some purposes, it is preferable to produce the polypeptide in a recombinant system in which the *E. cloacae* protein contains an additional sequence tag that facilitates purification, such as, but not limited to, a polyhistidine sequence. The polypeptide can then be purified from a crude lysate of the host cell by chromatography on an appropriate solid-phase matrix. Alternatively, antibodies produced against an *E. cloacae* protein or against peptides derived therefrom can be used as purification reagents. Other purification methods are possible.

10       The present invention also encompasses derivatives and homologues of *E. cloacae*-encoded polypeptides. For some purposes, nucleic acid sequences encoding the peptides may be altered by substitutions, additions, or deletions that provide for functionally equivalent molecules, i.e., function-conservative variants. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of similar properties, such as, for example, positively charged amino acids (arginine, lysine, and histidine); negatively charged amino acids (aspartate and glutamate); polar neutral amino acids; and non-polar amino acids.

20       The isolated polypeptides may be modified by, for example, phosphorylation, sulfation, acylation, or other protein modifications. They may also be modified with a label capable of providing a detectable signal, either directly or indirectly, including, but not limited to, radioisotopes and fluorescent compounds.

      To identify *E. cloacae*-derived polypeptides for use in the present invention, essentially the complete genomic sequence of a virulent, methicillin-resistant isolate of *Enterobacter cloacae* isolate was analyzed. While, in very rare instances, a nucleic acid sequencing error may be revealed, resolving a rare sequencing error is well



within the art, and such an occurrence will not prevent one skilled in the art from practicing the invention.

Also encompassed are any *E. cloacae* polypeptide sequences that are contained within the open reading frames (ORFs), including complete protein-coding sequences, of which any of SEQ ID NO: 1 - SEQ ID NO: 5662 forms a part. Table 2, which is appended herewith and which forms part of the present specification, provides a putative identification of the particular function of a polypeptide which is encoded by each ORF, based on the homology match (determined by the BLAST algorithm) of the predicted polypeptide with known proteins encoded by ORFs in other organisms.

As a result, one skilled in the art can use the polypeptides of the present invention for commercial and industrial purposes consistent with the type of putative identification of the polypeptide.

The present invention provides a library of *E. cloacae*-derived polypeptide sequences, and a corresponding library of nucleic acid sequences encoding the polypeptides, wherein the polypeptides themselves, or polypeptides contained within ORFs of which they form a part, comprise sequences that are contemplated for use as components of vaccines. Non-limiting examples of such sequences are listed by SEQ ID NO in Table 2, which is appended herewith and which forms part of the present specification.

The present invention also provides a library of *E. cloacae*-derived polypeptide sequences, and a corresponding library of nucleic acid sequences encoding the polypeptides, wherein the polypeptides themselves, or polypeptides contained within ORFs of which they form a part, comprise sequences lacking homology to any known prokaryotic or eukaryotic sequences. Such libraries provide probes, primers, and markers which can be used to diagnose *E. cloacae* infection, including use as markers

in epidemiological studies. Non-limiting examples of such sequences are listed by SEQ ID NO in Table 2, which is appended

The present invention also provides a library of *E. cloacae*-derived polypeptide sequences, and a corresponding library of nucleic acid sequences encoding the  
5 polypeptides, wherein the polypeptides themselves, or polypeptides contained within ORFs of which they form a part, comprise targets for therapeutic drugs.

Specific Example: Determination Of *Enterobacter* Protein Antigens For Antibody  
And Vaccine Development

10

The selection of *Enterobacter* protein antigens for vaccine development can be derived from the nucleic acids encoding *E. cloacae* polypeptides. First, the ORF's can be analyzed for homology to other known exported or membrane proteins and analyzed using the discriminant analysis described by Klein, et al. (Klein, P.,  
15 Kanehsia, M., and DeLisi, C. (1985) *Biochimica et Biophysica Acta* 815, 468-476) for predicting exported and membrane proteins.

Homology searches can be performed using the BLAST algorithm contained in the Wisconsin Sequence Analysis Package (Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711) to compare each predicted  
20 ORF amino acid sequence with all sequences found in the current GenBank, SWISS-PROT and PIR databases. BLAST searches for local alignments between the ORF and the databank sequences and reports a probability score which indicates the probability of finding this sequence by chance in the database. ORF's with significant homology (e.g. probabilities lower than  $1 \times 10^{-6}$  that the homology is only due to  
25 random chance) to membrane or exported proteins represent protein antigens for

vaccine development. Possible functions can be provided to *E. cloacae* genes based on sequence homology to genes cloned in other organisms.

Discriminant analysis (Klein, et al. supra) can be used to examine the ORF amino acid sequences. This algorithm uses the intrinsic information contained in the ORF amino acid sequence and compares it to information derived from the properties of known membrane and exported proteins. This comparison predicts which proteins will be exported, membrane associated or cytoplasmic. ORF amino acid sequences identified as exported or membrane associated by this algorithm are likely protein antigens for vaccine development.

10

#### Production of Fragments and Analogs of *E. cloacae* Nucleic Acids and Polypeptides

Based on the discovery of the *E. cloacae* gene products of the invention provided in the Sequence Listing, one skilled in the art can alter the disclosed structure of *E. cloacae* genes, e.g., by producing fragments or analogs, and test the newly produced structures for activity. Examples of techniques known to those skilled in the relevant art which allow the production and testing of fragments and analogs are discussed below. These, or analogous methods can be used to make and screen libraries of polypeptides, e.g., libraries of random peptides or libraries of fragments or analogs of cellular proteins for the ability to bind *E. cloacae* polypeptides. Such screens are useful for the identification of inhibitors of *E. cloacae*.

20

#### Generation of Fragments

Fragments of a protein can be produced in several ways, e.g., recombinantly, by proteolytic digestion, or by chemical synthesis. Internal or terminal fragments of a polypeptide can be generated by removing one or more nucleotides from one end (for

a terminal fragment) or both ends (for an internal fragment) of a nucleic acid which encodes the polypeptide. Expression of the mutagenized DNA produces polypeptide fragments. Digestion with "end-nibbling" endonucleases can thus generate DNAs which encode an array of fragments. DNAs which encode fragments of a protein can  
5 also be generated by random shearing, restriction digestion or a combination of the above-discussed methods.

Fragments can also be chemically synthesized using techniques known in the art such as conventional Merrifield solid phase f-Moc or t-Boc chemistry. For example, peptides of the present invention may be arbitrarily divided into fragments of  
10 desired length with no overlap of the fragments, or divided into overlapping fragments of a desired length.

#### Alteration of Nucleic Acids and Polypeptides: Random Methods

Amino acid sequence variants of a protein can be prepared by random  
15 mutagenesis of DNA which encodes a protein or a particular domain or region of a protein. Useful methods include PCR mutagenesis and saturation mutagenesis. A library of random amino acid sequence variants can also be generated by the synthesis of a set of degenerate oligonucleotide sequences. (Methods for screening proteins in a library of variants are elsewhere herein).

20

#### PCR Mutagenesis

In PCR mutagenesis, reduced Taq polymerase fidelity is used to introduce random mutations into a cloned fragment of DNA (Leung et al., 1989, *Technique* 1:11-15). The DNA region to be mutagenized is amplified using the polymerase  
25 chain reaction (PCR) under conditions that reduce the fidelity of DNA synthesis by Taq DNA polymerase, e.g., by using a dGTP/dATP ratio of five and adding  $Mn^{2+}$  to

the PCR reaction. The pool of amplified DNA fragments are inserted into appropriate cloning vectors to provide random mutant libraries.

### Saturation Mutagenesis

5 Saturation mutagenesis allows for the rapid introduction of a large number of single base substitutions into cloned DNA fragments (Mayers et al., 1985, *Science* 229:242). This technique includes generation of mutations, e.g., by chemical treatment or irradiation of single-stranded DNA *in vitro*, and synthesis of a complimentary DNA strand. The mutation frequency can be modulated by  
10 modulating the severity of the treatment, and essentially all possible base substitutions can be obtained. Because this procedure does not involve a genetic selection for mutant fragments both neutral substitutions, as well as those that alter function, are obtained. The distribution of point mutations is not biased toward conserved sequence elements.

15

### Degenerate Oligonucleotides

A library of homologs can also be generated from a set of degenerate oligonucleotide sequences. Chemical synthesis of a degenerate sequences can be carried out in an automatic DNA synthesizer, and the synthetic genes then ligated into  
20 an appropriate expression vector. The synthesis of degenerate oligonucleotides is known in the art (see for example, Narang, SA (1983) *Tetrahedron* 39:3; Itakura et al. (1981) *Recombinant DNA, Proc 3rd Cleveland Sympos. Macromolecules*, ed. AG Walton, Amsterdam: Elsevier pp273-289; Itakura et al. (1984) *Annu. Rev. Biochem.* 53:323; Itakura et al. (1984) *Science* 198:1056; Ike et al. (1983) *Nucleic Acid Res.*  
25 11:477. Such techniques have been employed in the directed evolution of other proteins (see, for example, Scott et al. (1990) *Science* 249:386-390; Roberts et al.

(1992) *PNAS* 89:2429-2433; Devlin et al. (1990) *Science* 249: 404-406; Cwirla et al. (1990) *PNAS* 87: 6378-6382; as well as U.S. Patents Nos. 5,223,409, 5,198,346, and 5,096,815).

5    Alteration of Nucleic Acids and Polypeptides: Methods for Directed Mutagenesis

Non-random or directed, mutagenesis techniques can be used to provide specific sequences or mutations in specific regions. These techniques can be used to create variants which include, e.g., deletions, insertions, or substitutions, of residues of the known amino acid sequence of a protein. The sites for mutation can be modified individually or in series, e.g., by (1) substituting first with conserved amino acids and then with more radical choices depending upon results achieved, (2) deleting the target residue, or (3) inserting residues of the same or a different class adjacent to the located site, or combinations of options 1-3.

15    Alanine Scanning Mutagenesis

Alanine scanning mutagenesis is a useful method for identification of certain residues or regions of the desired protein that are preferred locations or domains for mutagenesis, Cunningham and Wells (*Science* 244:1081-1085, 1989). In alanine scanning, a residue or group of target residues are identified (e.g., charged residues such as Arg, Asp, His, Lys, and Glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine). Replacement of an amino acid can affect the interaction of the amino acids with the surrounding aqueous environment in or outside the cell. Those domains demonstrating functional sensitivity to the substitutions are then refined by introducing further or other variants at or for the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation per se need not be

predetermined. For example, to optimize the performance of a mutation at a given site, alanine scanning or random mutagenesis may be conducted at the target codon or region and the expressed desired protein subunit variants are screened for the optimal combination of desired activity.

5

#### Oligonucleotide-Mediated Mutagenesis

Oligonucleotide-mediated mutagenesis is a useful method for preparing substitution, deletion, and insertion variants of DNA, see, e.g., Adelman et al., (*DNA* 2:183, 1983). Briefly, the desired DNA is altered by hybridizing an oligonucleotide  
10 encoding a mutation to a DNA template, where the template is the single-stranded form of a plasmid or bacteriophage containing the unaltered or native DNA sequence of the desired protein. After hybridization, a DNA polymerase is used to synthesize an entire second complementary strand of the template that will thus incorporate the oligonucleotide primer, and will code for the selected alteration in the desired protein  
15 DNA. Generally, oligonucleotides of at least about 25 nucleotides in length are used. An optimal oligonucleotide will have 12 to 15 nucleotides that are completely complementary to the template on either side of the nucleotide(s) coding for the mutation. This ensures that the oligonucleotide will hybridize properly to the single-stranded DNA template molecule. The oligonucleotides are readily synthesized using  
20 techniques known in the art such as that described by Crea et al. (*Proc. Natl. Acad. Sci. USA*, 75: 5765[1978]).

#### Cassette Mutagenesis

Another method for preparing variants, cassette mutagenesis, is based on the  
25 technique described by Wells et al. (*Gene*, 34:315[1985]). The starting material is a plasmid (or other vector) which includes the protein subunit DNA to be mutated. The

codon(s) in the protein subunit DNA to be mutated are identified. There must be a unique restriction endonuclease site on each side of the identified mutation site(s). If no such restriction sites exist, they may be generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at appropriate

- 5 locations in the desired protein subunit DNA. After the restriction sites have been introduced into the plasmid, the plasmid is cut at these sites to linearize it. A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the desired mutation(s) is synthesized using standard procedures. The two strands are synthesized separately and then hybridized together using standard
- 10 techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 3' and 5' ends that are comparable with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated desired protein subunit DNA sequence.

## 15 Combinatorial Mutagenesis

- Combinatorial mutagenesis can also be used to generate mutants (Ladner et al., WO 88/06630). In this method, the amino acid sequences for a group of homologs or other related proteins are aligned, preferably to promote the highest homology possible. All of the amino acids which appear at a given position of the aligned
- 20 sequences can be selected to create a degenerate set of combinatorial sequences. The variegated library of variants is generated by combinatorial mutagenesis at the nucleic acid level, and is encoded by a variegated gene library. For example, a mixture of synthetic oligonucleotides can be enzymatically ligated into gene sequences such that the degenerate set of potential sequences are expressible as individual peptides, or
- 25 alternatively, as a set of larger fusion proteins containing the set of degenerate sequences.



### Other Modifications of *E. cloacae* Nucleic Acids and Polypeptides

It is possible to modify the structure of an *E. cloacae* polypeptide for such purposes as increasing solubility, enhancing stability (e.g., shelf life *ex vivo* and  
5 resistance to proteolytic degradation *in vivo*). A modified *E. cloacae* protein or peptide can be produced in which the amino acid sequence has been altered, such as by amino acid substitution, deletion, or addition as described herein.

An *E. cloacae* peptide can also be modified by substitution of cysteine residues preferably with alanine, serine, threonine, leucine or glutamic acid residues to  
10 minimize dimerization via disulfide linkages. In addition, amino acid side chains of fragments of the protein of the invention can be chemically modified. Another modification is cyclization of the peptide.

In order to enhance stability and/or reactivity, an *E. cloacae* polypeptide can be modified to incorporate one or more polymorphisms in the amino acid sequence of the  
15 protein resulting from any natural allelic variation. Additionally, D-amino acids, non-natural amino acids, or non-amino acid analogs can be substituted or added to produce a modified protein within the scope of this invention. Furthermore, an *E. cloacae* polypeptide can be modified using polyethylene glycol (PEG) according to the method of A. Schon and co-workers (Wie et al., *supra*) to produce a protein conjugated with  
20 PEG. In addition, PEG can be added during chemical synthesis of the protein. Other modifications of *E. cloacae* proteins include reduction/alkylation (Tarr, *Methods of Protein Microcharacterization*, J. E. Silver ed., Humana Press, Clifton NJ 155-194 (1986)); acylation (Tarr, *supra*); chemical coupling to an appropriate carrier (Mishell and Shiigi, eds, *Selected Methods in Cellular Immunology*, WH Freeman, San  
25 Francisco, CA (1980), U.S. Patent 4,939,239; or mild formalin treatment (Marsh, (1971) *Int. Arch. of Allergy and Appl. Immunol.*, 41: 199 - 215).

To facilitate purification and potentially increase solubility of an *E. cloacae* protein or peptide, it is possible to add an amino acid fusion moiety to the peptide backbone. For example, hexa-histidine can be added to the protein for purification by immobilized metal ion affinity chromatography (Hochuli, E. et al., (1988)

- 5 *Bio/Technology*, 6: 1321 - 1325). In addition, to facilitate isolation of peptides free of irrelevant sequences, specific endoprotease cleavage sites can be introduced between the sequences of the fusion moiety and the peptide.

- To potentially aid proper antigen processing of epitopes within an *E. cloacae* polypeptide, canonical protease sensitive sites can be engineered between regions,  
10 each comprising at least one epitope via recombinant or synthetic methods. For example, charged amino acid pairs, such as KK or RR, can be introduced between regions within a protein or fragment during recombinant construction thereof. The resulting peptide can be rendered sensitive to cleavage by cathepsin and/or other trypsin-like enzymes which would generate portions of the protein containing one or  
15 more epitopes. In addition, such charged amino acid residues can result in an increase in the solubility of the peptide.

#### Primary Methods for Screening Polypeptides and Analogs

- Various techniques are known in the art for screening generated mutant gene  
20 products. Techniques for screening large gene libraries often include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the genes under conditions in which detection of a desired activity, e.g., in this case, binding to *E. cloacae* polypeptide or an interacting protein, facilitates relatively easy isolation of the vector encoding the  
25 gene whose product was detected. Each of the techniques described below is

amenable to high through-put analysis for screening large numbers of sequences created, e.g., by random mutagenesis techniques.

### Two Hybrid Systems

- 5 Two hybrid assays such as the system described below (as with the other screening methods described herein), can be used to identify polypeptides, e.g., fragments or analogs of a naturally-occurring *E. cloacae* polypeptide, e.g., of cellular proteins, or of randomly generated polypeptides which bind to an *E. cloacae* protein. (The *E. cloacae* domain is used as the bait protein and the library of variants are
- 10 expressed as prey fusion proteins.) In an analogous fashion, a two hybrid assay (as with the other screening methods described herein), can be used to find polypeptides which bind an *E. cloacae* polypeptide.

### Display Libraries

- 15 In one approach to screening assays, the Enterobacter peptides are displayed on the surface of a cell or viral particle, and the ability of particular cells or viral particles to bind an appropriate receptor protein via the displayed product is detected in a "panning assay". For example, the gene library can be cloned into the gene for a surface membrane protein of a bacterial cell, and the resulting fusion protein detected
- 20 by panning (Ladner et al., WO 88/06630; Fuchs et al. (1991) *Bio/Technology* 9:1370-1371; and Goward et al. (1992) *TIBS* 18:136-140). In a similar fashion, a detectably labeled ligand can be used to score for potentially functional peptide homologs. Fluorescently labeled ligands, e.g., receptors, can be used to detect homologs which retain ligand-binding activity. The use of fluorescently labeled ligands, allows cells to
- 25 be visually inspected and separated under a fluorescence microscope, or, where the

morphology of the cell permits, to be separated by a fluorescence-activated cell sorter.

A gene library can be expressed as a fusion protein on the surface of a viral particle. For instance, in the filamentous phage system, foreign peptide sequences can be expressed on the surface of infectious phage, thereby conferring two significant benefits. First, since these phage can be applied to affinity matrices at concentrations well over  $10^{13}$  phage per milliliter, a large number of phage can be screened at one time. Second, since each infectious phage displays a gene product on its surface, if a particular phage is recovered from an affinity matrix in low yield, the phage can be amplified by another round of infection. The group of almost identical *E. coli* filamentous phages, M13, fd., and f1, are most often used in phage display libraries. Either of the phage gIII or gVIII coat proteins can be used to generate fusion proteins without disrupting the ultimate packaging of the viral particle. Foreign epitopes can be expressed at the NH<sub>2</sub>-terminal end of pIII and phage bearing such epitopes recovered from a large excess of phage lacking this epitope (Ladner et al. PCT publication WO 90/02909; Garrard et al., PCT publication WO 92/09690; Marks et al. (1992) *J. Biol. Chem.* 267:16007-16010; Griffiths et al. (1993) *EMBO J* 12:725-734; Clackson et al. (1991) *Nature* 352:624-628; and Barbas et al. (1992) *PNAS* 89:4457-4461).

A common approach uses the maltose receptor of *E. coli* (the outer membrane protein, LamB) as a peptide fusion partner (Charbit et al. (1986) *EMBO* 5, 3029-3037). Oligonucleotides have been inserted into plasmids encoding the LamB gene to produce peptides fused into one of the extracellular loops of the protein. These peptides are available for binding to ligands, e.g., to antibodies, and can elicit an immune response when the cells are administered to animals. Other cell surface proteins, e.g., OmpA (Schorr et al. (1991) *Vaccines* 91, pp. 387-392), PhoE

(Agterberg, et al. (1990) *Gene* 88, 37-45), and PAL (Fuchs et al. (1991) *Bio/Tech* 9, 1369-1372), as well as large bacterial surface structures have served as vehicles for peptide display. Peptides can be fused to pilin, a protein which polymerizes to form the pilus-a conduit for interbacterial exchange of genetic information (Thiry et al.

5 (1989) *Appl. Environ. Microbiol.* 55, 984-993). Because of its role in interacting with other cells, the pilus provides a useful support for the presentation of peptides to the extracellular environment. Another large surface structure used for peptide display is the bacterial motive organ, the flagellum. Fusion of peptides to the subunit protein flagellin offers a dense array of many peptide copies on the host cells (Kuwajima et  
10 al. (1988) *Bio/Tech.* 6, 1080-1083). Surface proteins of other bacterial species have also served as peptide fusion partners. Examples include the *Staphylococcus* protein A and the outer membrane IgA protease of *Neisseria* (Hansson et al. (1992) *J. Bacteriol.* 174, 4239-4245 and Klauser et al. (1990) *EMBO J.* 9, 1991-1999).

In the filamentous phage systems and the LamB system described above, the  
15 physical link between the peptide and its encoding DNA occurs by the containment of the DNA within a particle (cell or phage) that carries the peptide on its surface. Capturing the peptide captures the particle and the DNA within. An alternative scheme uses the DNA-binding protein LacI to form a link between peptide and DNA (Cull et al. (1992) *PNAS USA* 89:1865-1869). This system uses a plasmid containing  
20 the LacI gene with an oligonucleotide cloning site at its 3'-end. Under the controlled induction by arabinose, a LacI-peptide fusion protein is produced. This fusion retains the natural ability of LacI to bind to a short DNA sequence known as LacO operator (LacO). By installing two copies of LacO on the expression plasmid, the LacI-peptide fusion binds tightly to the plasmid that encoded it. Because the plasmids in each cell  
25 contain only a single oligonucleotide sequence and each cell expresses only a single peptide sequence, the peptides become specifically and stably associated with the

DNA sequence that directed its synthesis. The cells of the library are gently lysed and the peptide-DNA complexes are exposed to a matrix of immobilized receptor to recover the complexes containing active peptides. The associated plasmid DNA is then reintroduced into cells for amplification and DNA sequencing to determine the identity of the peptide ligands. As a demonstration of the practical utility of the method, a large random library of dodecapeptides was made and selected on a monoclonal antibody raised against the opioid peptide dynorphin B. A cohort of peptides was recovered, all related by a consensus sequence corresponding to a six-residue portion of dynorphin B. (Cull et al. (1992) *Proc. Natl. Acad. Sci. U.S.A.* 89-1869)

This scheme, sometimes referred to as peptides-on-plasmids, differs in two important ways from the phage display methods. First, the peptides are attached to the C-terminus of the fusion protein, resulting in the display of the library members as peptides having free carboxy termini. Both of the filamentous phage coat proteins, pIII and pVIII, are anchored to the phage through their C-termini, and the guest peptides are placed into the outward-extending N-terminal domains. In some designs, the phage-displayed peptides are presented right at the amino terminus of the fusion protein. (Cwirla, et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.* 87, 6378-6382) A second difference is the set of biological biases affecting the population of peptides actually present in the libraries. The LacI fusion molecules are confined to the cytoplasm of the host cells. The phage coat fusions are exposed briefly to the cytoplasm during translation but are rapidly secreted through the inner membrane into the periplasmic compartment, remaining anchored in the membrane by their C-terminal hydrophobic domains, with the N-termini, containing the peptides, protruding into the periplasm while awaiting assembly into phage particles. The peptides in the LacI and phage libraries may differ significantly as a result of their exposure to different proteolytic

activities. The phage coat proteins require transport across the inner membrane and signal peptidase processing as a prelude to incorporation into phage. Certain peptides exert a deleterious effect on these processes and are underrepresented in the libraries (Gallop et al. (1994) *J. Med. Chem.* 37(9):1233-1251). These particular biases are not  
5 a factor in the LacI display system.

The number of small peptides available in recombinant random libraries is enormous. Libraries of  $10^7$ - $10^9$  independent clones are routinely prepared. Libraries as large as  $10^{11}$  recombinants have been created, but this size approaches the practical limit for clone libraries. This limitation in library size occurs at the step of  
10 transforming the DNA containing randomized segments into the host bacterial cells. To circumvent this limitation, an *in vitro* system based on the display of nascent peptides in polysome complexes has recently been developed. This display library method has the potential of producing libraries 3-6 orders of magnitude larger than the currently available phage/phagemid or plasmid libraries. Furthermore, the construction  
15 of the libraries, expression of the peptides, and screening, is done in an entirely cell-free format.

In one application of this method (Gallop et al. (1994) *J. Med. Chem.* 37(9):1233-1251), a molecular DNA library encoding  $10^{12}$  decapeptides was constructed and the library expressed in an *E. coli* S30 *in vitro* coupled  
20 transcription/translation system. Conditions were chosen to stall the ribosomes on the mRNA, causing the accumulation of a substantial proportion of the RNA in polysomes and yielding complexes containing nascent peptides still linked to their encoding RNA. The polysomes are sufficiently robust to be affinity purified on immobilized receptors in much the same way as the more conventional recombinant  
25 peptide display libraries are screened. RNA from the bound complexes is recovered, converted to cDNA, and amplified by PCR to produce a template for the next round

of synthesis and screening. The polysome display method can be coupled to the phage display system. Following several rounds of screening, cDNA from the enriched pool of polysomes was cloned into a phagemid vector. This vector serves as both a peptide expression vector, displaying peptides fused to the coat proteins, and as  
5 a DNA sequencing vector for peptide identification. By expressing the polysome-derived peptides on phage, one can either continue the affinity selection procedure in this format or assay the peptides on individual clones for binding activity in a phage ELISA, or for binding specificity in a completion phage ELISA (Barret, et al. (1992) *Anal. Biochem* 204,357-364). To identify the sequences of the active peptides one  
10 sequences the DNA produced by the phagemid host.

#### Secondary Screening of Polypeptides and Analogs

The high through-put assays described above can be followed by secondary screens in order to identify further biological activities which will, e.g., allow one  
15 skilled in the art to differentiate agonists from antagonists. The type of a secondary screen used will depend on the desired activity that needs to be tested. For example, an assay can be developed in which the ability to inhibit an interaction between a protein of interest and its respective ligand can be used to identify antagonists from a group of peptide fragments isolated though one of the primary screens described  
20 above.

Therefore, methods for generating fragments and analogs and testing them for activity are known in the art. Once the core sequence of interest is identified, it is routine for one skilled in the art to obtain analogs and fragments.



### Peptide Mimetics of *E. cloacae* Polypeptides

The invention also provides for reduction of the protein binding domains of the subject *E. cloacae* polypeptides to generate mimetics, e.g. peptide or non-peptide agents. The peptide mimetics are able to disrupt binding of a polypeptide to its counter ligand, e.g., in the case of an *E. cloacae* polypeptide binding to a naturally occurring ligand. The critical residues of a subject *E. cloacae* polypeptide which are involved in molecular recognition of a polypeptide can be determined and used to generate *E. cloacae*-derived peptidomimetics which competitively or noncompetitively inhibit binding of the *E. cloacae* polypeptide with an interacting polypeptide (see, for example, European patent applications EP-412,762A and EP-B31,080A).

For example, scanning mutagenesis can be used to map the amino acid residues of a particular *E. cloacae* polypeptide involved in binding an interacting polypeptide, peptidomimetic compounds (e.g. diazepam or isoquinoline derivatives) can be generated which mimic those residues in binding to an interacting polypeptide, and which therefore can inhibit binding of an *E. cloacae* polypeptide to an interacting polypeptide and thereby interfere with the function of *E. cloacae* polypeptide. For instance, non-hydrolyzable peptide analogs of such residues can be generated using benzodiazepine (e.g., see Freidinger et al. in *Peptides: Chemistry and Biology*, G.R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), azepine (e.g., see Huffman et al. in *Peptides: Chemistry and Biology*, G.R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), substituted gamma lactam rings (Garvey et al. in *Peptides: Chemistry and Biology*, G.R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), keto-methylene pseudopeptides (Ewenson et al. (1986) *J Med Chem* 29:295; and Ewenson et al. in *Peptides: Structure and Function* (Proceedings of the 9th American Peptide Symposium) Pierce Chemical Co. Rockland, IL, 1985), b-turn dipeptide cores (Nagai et al. (1985) *Tetrahedron Lett* 26:647; and Sato et al.

(1986) *J Chem Soc Perkin Trans* 1:1231), and b-aminoalcohols (Gordon et al. (1985) *Biochem Biophys Res Commun* 126:419; and et al. (1986) *Biochem Biophys Res Commun* 134:71).

5 Vaccine Formulations for *E. cloacae* Nucleic Acids and Polypeptides

This invention also features vaccine compositions for protection against infection by *E. cloacae* or for treatment of *E. cloacae* infection. In one embodiment, the vaccine compositions contain one or more immunogenic components such as a surface protein from *E. cloacae*, or portion thereof, and a pharmaceutically acceptable  
10 carrier. Nucleic acids within the scope of the invention are exemplified by the nucleic acids of the invention contained in the Sequence Listing which encode *E. cloacae* surface proteins. Any nucleic acid encoding an immunogenic *E. cloacae* protein, or portion thereof, which is capable of expression in a cell, can be used in the present invention. These vaccines have therapeutic and prophylactic utilities.

15 One aspect of the invention provides a vaccine composition for protection against infection by *E. cloacae* which contains at least one immunogenic fragment of an *E. cloacae* protein and a pharmaceutically acceptable carrier. Preferred fragments include peptides of at least about 10 amino acid residues in length, preferably about 10-20 amino acid residues in length, and more preferably about 12-16 amino acid  
20 residues in length.

Immunogenic components of the invention can be obtained, for example, by screening polypeptides recombinantly produced from the corresponding fragment of the nucleic acid encoding the full-length *E. cloacae* protein. In addition, fragments can be chemically synthesized using techniques known in the art such as conventional  
25 Merrifield solid phase f-Moc or t-Boc chemistry.

In one embodiment, immunogenic components are identified by the ability of the peptide to stimulate T cells. Peptides which stimulate T cells, as determined by, for example, T cell proliferation or cytokine secretion are defined herein as comprising at least one T cell epitope. T cell epitopes are believed to be involved in initiation and perpetuation of the immune response to the protein allergen which is responsible for the clinical symptoms of allergy. These T cell epitopes are thought to trigger early events at the level of the T helper cell by binding to an appropriate HLA molecule on the surface of an antigen presenting cell, thereby stimulating the T cell subpopulation with the relevant T cell receptor for the epitope. These events lead to T cell proliferation, lymphokine secretion, local inflammatory reactions, recruitment of additional immune cells to the site of antigen/T cell interaction, and activation of the B cell cascade, leading to the production of antibodies. A T cell epitope is the basic element, or smallest unit of recognition by a T cell receptor, where the epitope comprises amino acids essential to receptor recognition (e.g., approximately 6 or 7 amino acid residues). Amino acid sequences which mimic those of the T cell epitopes are within the scope of this invention.

Screening immunogenic components can be accomplished using one or more of several different assays. For example, *in vitro*, peptide T cell stimulatory activity is assayed by contacting a peptide known or suspected of being immunogenic with an antigen presenting cell which presents appropriate MHC molecules in a T cell culture. Presentation of an immunogenic *E. cloacae* peptide in association with appropriate MHC molecules to T cells in conjunction with the necessary co-stimulation has the effect of transmitting a signal to the T cell that induces the production of increased levels of cytokines, particularly of interleukin-2 and interleukin-4. The culture supernatant can be obtained and assayed for interleukin-2 or other known cytokines. For example, any one of several conventional assays for interleukin-2 can be

employed, such as the assay described in *Proc. Natl. Acad. Sci USA*, 86: 1333 (1989) the pertinent portions of which are incorporated herein by reference. A kit for an assay for the production of interferon is also available from Genzyme Corporation (Cambridge, MA).

- 5           Alternatively, a common assay for T cell proliferation entails measuring tritiated thymidine incorporation. The proliferation of T cells can be measured *in vitro* by determining the amount of <sup>3</sup>H-labeled thymidine incorporated into the replicating DNA of cultured cells. Therefore, the rate of DNA synthesis and, in turn, the rate of cell division can be quantified.
- 10           Vaccine compositions of the invention containing immunogenic components (e.g., *E. cloacae* polypeptide or fragment thereof or nucleic acid encoding an *E. cloacae* polypeptide or fragment thereof) preferably include a pharmaceutically acceptable carrier. The term "pharmaceutically acceptable carrier" refers to a carrier that does not cause an allergic reaction or other untoward effect in patients to whom it
- 15 is administered. Suitable pharmaceutically acceptable carriers include, for example, one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. Pharmaceutically acceptable carriers may further comprise minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or
- 20 effectiveness of the antibody. For vaccines of the invention containing *E. cloacae* polypeptides, the polypeptide is co-administered with a suitable adjuvant.

It will be apparent to those of skill in the art that the therapeutically effective amount of DNA or protein of this invention will depend, *inter alia*, upon the administration schedule, the unit dose of antibody administered, whether the protein or

25 DNA is administered in combination with other therapeutic agents, the immune status and health of the patient, and the therapeutic activity of the particular protein or DNA.

Vaccine compositions are conventionally administered parenterally, e.g., by injection, either subcutaneously or intramuscularly. Methods for intramuscular immunization are described by Wolff et al. (1990) *Science* 247: 1465-1468 and by Sedegah et al. (1994) *Immunology* 91: 9866-9870. Other modes of administration

5 include oral and pulmonary formulations, suppositories, and transdermal applications. Oral immunization is preferred over parenteral methods for inducing protection against infection by *E. cloacae*. Cain et. al. (1993) *Vaccine* 11: 637-642. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose,

10 magnesium carbonate, and the like.

The vaccine compositions of the invention can include an adjuvant, including, but not limited to aluminum hydroxide; N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP); N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP); N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-

15 sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE); RIBI, which contains three components from bacteria; monophosphoryl lipid A; trehalose dimycolate; cell wall skeleton (MPL + TDM + CWS) in a 2% squalene/Tween 80 emulsion; and cholera toxin. Others which may be used are non-toxic derivatives of cholera toxin, including its B subunit, and/or conjugates or

20 genetically engineered fusions of the *E. cloacae* polypeptide with cholera toxin or its B subunit, procholeraenoid, fungal polysaccharides, including schizophyllan, muramyl dipeptide, muramyl dipeptide derivatives, phorbol esters, labile toxin of *E. coli*, non-*E. cloacae* bacterial lysates, block polymers or saponins.

Other suitable delivery methods include biodegradable microcapsules or

25 immuno-stimulating complexes (ISCOMs), cochleates, or liposomes, genetically engineered attenuated live vectors such as viruses or bacteria, and recombinant

(chimeric) virus-like particles, e.g., bluetongue. The amount of adjuvant employed will depend on the type of adjuvant used. For example, when the mucosal adjuvant is cholera toxin, it is suitably used in an amount of 5 mg to 50 mg, for example 10 mg to 35 mg. When used in the form of microcapsules, the amount used will depend on  
5 the amount employed in the matrix of the microcapsule to achieve the desired dosage. The determination of this amount is within the skill of a person of ordinary skill in the art.

Carrier systems in humans may include enteric release capsules protecting the antigen from the acidic environment of the stomach, and including *E. cloacae*  
10 polypeptide in an insoluble form as fusion proteins. Suitable carriers for the vaccines of the invention are enteric coated capsules and polylactide-glycolide microspheres. Suitable diluents are 0.2 N NaHCO<sub>3</sub> and/or saline.

Vaccines of the invention can be administered as a primary prophylactic agent in adults or in children, as a secondary prevention, after successful eradication of *E.*  
15 *cloacae* in an infected host, or as a therapeutic agent in the aim to induce an immune response in a susceptible host to prevent infection by *E. cloacae*. The vaccines of the invention are administered in amounts readily determined by persons of ordinary skill in the art. Thus, for adults a suitable dosage will be in the range of 10 mg to 10 g, preferably 10 mg to 100 mg. A suitable dosage for adults will also be in the range  
20 of 5 mg to 500 mg. Similar dosage ranges will be applicable for children. Those skilled in the art will recognize that the optimal dose may be more or less depending upon the patient's body weight, disease, the route of administration, and other factors. Those skilled in the art will also recognize that appropriate dosage levels can be obtained based on results with known oral vaccines such as, for example, a vaccine  
25 based on an *E. coli* lysate (6 mg dose daily up to total of 540 mg) and with an enterotoxigenic *E. coli* purified antigen (4 doses of 1 mg) (Schulman et al., *J. Urol.*

150:917-921 (1993); Boedecker et al., *American Gastroenterological Assoc.* 999:A-222 (1993)). The number of doses will depend upon the disease, the formulation, and efficacy data from clinical trials. Without intending any limitation as to the course of treatment, the treatment can be administered over 3 to 8 doses for a primary  
5 immunization schedule over 1 month (Boedeker, *American Gastroenterological Assoc.* 888:A-222 (1993)).

In a preferred embodiment, a vaccine composition of the invention can be based on a killed whole *E. coli* preparation with an immunogenic fragment of an *E. cloacae* protein of the invention expressed on its surface or it can be based on an *E.*  
10 *coli* lysate, wherein the killed *E. coli* acts as a carrier or an adjuvant.

It will be apparent to those skilled in the art that some of the vaccine compositions of the invention are useful only for preventing *E. cloacae* infection, some are useful only for treating *E. cloacae* infection, and some are useful for both preventing and treating *E. cloacae* infection. In a preferred embodiment, the vaccine  
15 composition of the invention provides protection against *E. cloacae* infection by stimulating humoral and/or cell-mediated immunity against *E. cloacae*. It should be understood that amelioration of any of the symptoms of *E. cloacae* infection is a desirable clinical goal, including a lessening of the dosage of medication used to treat *E. cloacae*-caused disease, or an increase in the production of antibodies in the serum  
20 or mucous of patients.

#### Antibodies Reactive With *E. cloacae* Polypeptides

The invention also includes antibodies specifically reactive with the subject *E. cloacae* polypeptide. Anti-protein/anti-peptide antisera or monoclonal antibodies can  
25 be made by standard protocols (See, for example, *Antibodies: A Laboratory Manual* ed. by Harlow and Lane (Cold Spring Harbor Press: 1988)). A mammal such as a

mouse, a hamster or rabbit can be immunized with an immunogenic form of the peptide. Techniques for conferring immunogenicity on a protein or peptide include conjugation to carriers or other techniques well known in the art. An immunogenic portion of the subject *E. cloacae* polypeptide can be administered in the presence of  
5 adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassays can be used with the immunogen as antigen to assess the levels of antibodies.

In a preferred embodiment, the subject antibodies are immunospecific for antigenic determinants of the *E. cloacae* polypeptides of the invention, e.g. antigenic  
10 determinants of a polypeptide of the invention contained in the Sequence Listing, or a closely related human or non-human mammalian homolog (e.g., 90% homologous, more preferably at least about 95% homologous). In yet a further preferred embodiment of the invention, the anti-*E. cloacae* antibodies do not substantially cross react (i.e., react specifically) with a protein which is for example, less than 80%  
15 percent homologous to a sequence of the invention contained in the Sequence Listing. By "not substantially cross react", it is meant that the antibody has a binding affinity for a non-homologous protein which is less than 10 percent, more preferably less than 5 percent, and even more preferably less than 1 percent, of the binding affinity for a protein of the invention contained in the Sequence Listing. In a most preferred  
20 embodiment, there is no cross-reactivity between bacterial and mammalian antigens.

The term antibody as used herein is intended to include fragments thereof which are also specifically reactive with *E. cloacae* polypeptides. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for whole antibodies. For example, F(ab')<sub>2</sub>  
25 fragments can be generated by treating antibody with pepsin. The resulting F(ab')<sub>2</sub> fragment can be treated to reduce disulfide bridges to produce Fab' fragments. The



antibody of the invention is further intended to include bispecific and chimeric molecules having an anti-*E. cloacae* portion.

Both monoclonal and polyclonal antibodies (Ab) directed against *E. cloacae* polypeptides or *E. cloacae* polypeptide variants, and antibody fragments such as Fab`  
5 and F(ab`)<sub>2</sub>, can be used to block the action of *E. cloacae* polypeptide and allow the study of the role of a particular *E. cloacae* polypeptide of the invention in aberrant or unwanted intracellular signaling, as well as the normal cellular function of the *E. cloacae* and by microinjection of anti-*E. cloacae* polypeptide antibodies of the present invention.

10 Antibodies which specifically bind *E. cloacae* epitopes can also be used in immunohistochemical staining of tissue samples in order to evaluate the abundance and pattern of expression of *E. cloacae* antigens. Anti-*E. cloacae* polypeptide antibodies can be used diagnostically in immuno-precipitation and immuno-blotting to detect and evaluate *E. cloacae* levels in tissue or bodily fluid as part of a clinical  
15 testing procedure. Likewise, the ability to monitor *E. cloacae* polypeptide levels in an individual can allow determination of the efficacy of a given treatment regimen for an individual afflicted with such a disorder. The level of an *E. cloacae* polypeptide can be measured in cells found in bodily fluid, such as in urine samples or can be measured in tissue, such as produced by gastric biopsy. Diagnostic assays using anti-  
20 *E. cloacae* antibodies can include, for example, immunoassays designed to aid in early diagnosis of *E. cloacae* infections. The present invention can also be used as a method of detecting antibodies contained in samples from individuals infected by this bacterium using specific *E. cloacae* antigens.

Another application of anti-*E. cloacae* polypeptide antibodies of the invention  
25 is in the immunological screening of cDNA libraries constructed in expression vectors such as  $\lambda$ gt11,  $\lambda$ gt18-23,  $\lambda$ ZAP, and  $\lambda$ ORF8. Messenger libraries of this type,

having coding sequences inserted in the correct reading frame and orientation, can produce fusion proteins. For instance,  $\lambda$ gt11 will produce fusion proteins whose amino termini consist of  $\beta$ -galactosidase amino acid sequences and whose carboxy termini consist of a foreign polypeptide. Antigenic epitopes of a subject *E. cloacae* polypeptide can then be detected with antibodies, as, for example, reacting nitrocellulose filters lifted from infected plates with anti-*E. cloacae* polypeptide antibodies. Phage, scored by this assay, can then be isolated from the infected plate. Thus, the presence of *E. cloacae* gene homologs can be detected and cloned from other species, and alternate isoforms (including splicing variants) can be detected and cloned.

#### Kits Containing Nucleic Acids, Polypeptides or Antibodies of the Invention

The nucleic acid, polypeptides and antibodies of the invention can be combined with other reagents and articles to form kits. Kits for diagnostic purposes typically comprise the nucleic acid, polypeptides or antibodies in vials or other suitable vessels. Kits typically comprise other reagents for performing hybridization reactions, polymerase chain reactions (PCR), or for reconstitution of lyophilized components, such as aqueous media, salts, buffers, and the like. Kits may also comprise reagents for sample processing such as detergents, chaotropic salts and the like. Kits may also comprise immobilization means such as particles, supports, wells, dipsticks and the like. Kits may also comprise labeling means such as dyes, developing reagents, radioisotopes, fluorescent agents, luminescent or chemiluminescent agents, enzymes, intercalating agents and the like. With the nucleic acid and amino acid sequence information provided herein, individuals skilled in art can readily assemble kits to serve their particular purpose. Kits further can include instructions for use.

## Bio chip Technology

The nucleic acid sequence of the present invention may be used to detect *E. cloacae* or other species of *Enterobacter* acid sequence using bio chip technology. Bio chips containing arrays of nucleic acid sequence can also be used to measure expression of genes of *E. cloacae* or other species of *Enterobacter*. For example, to diagnose a patient with a *E. cloacae* or other *Enterobacter* infection, a sample from a human or animal can be used as a probe on a bio chip containing an array of nucleic acid sequence from the present invention. In addition, a sample from a disease state can be compared to a sample from a non-disease state which would help identify a gene that is up-regulated or expressed in the disease state. This would provide valuable insight as to the mechanism by which the disease manifests. Changes in gene expression can also be used to identify critical pathways involved in drug transport or metabolism, and may enable the identification of novel targets involved in virulence or host cell interactions involved in maintenance of an infection. Procedures using such techniques have been described by Brown *et al.*, 1995, *Science* 270: 467-470.

Bio chips can also be used to monitor the genetic changes of potential therapeutic compounds including, deletions, insertions or mismatches. Once the therapeutic is added to the patient, changes to the genetic sequence can be evaluated for its efficacy. In addition, the nucleic acid sequence of the present invention can be used to determine essential genes in cell cycling. As described in Iyer *et al.*, 1999 (*Science*, 283:83-87 ) genes essential in the cell cycle can be identified using bio chips. Furthermore, the present invention provides nucleic acid sequence which can be used with bio chip technology to understand regulatory networks in bacteria, measure the response to environmental signals or drugs as in drug screening, and

study virulence induction. (Mons *et al.*, 1998, *Nature Biotechnology*, 16: 45-48.

Patents teaching this technology include U.S. Patents 5445934, 5744305, and 5800992.

#### Drug Screening Assays Using *E. cloacae* Polypeptides

- 5 By making available purified and recombinant *E. cloacae* polypeptides, the present invention provides assays which can be used to screen for drugs which are either agonists or antagonists of the normal cellular function, in this case, of the subject *E. cloacae* polypeptides, or of their role in intracellular signaling. Such inhibitors or potentiators may be useful as new therapeutic agents to combat *E.*
- 10 *cloacae* infections in humans. A variety of assay formats will suffice and, in light of the present inventions, will be comprehended by the person skilled in the art.

In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in

15 cell-free systems, such as may be derived with purified or semi-purified proteins, are often preferred as "primary" screens in that they can be generated to permit rapid development and relatively easy detection of an alteration in a molecular target which is mediated by a test compound. Moreover, the effects of cellular toxicity and/or bioavailability of the test compound can be generally ignored in the *in vitro* system,

20 the assay instead being focused primarily on the effect of the drug on the molecular target as may be manifest in an alteration of binding affinity with other proteins or change in enzymatic properties of the molecular target. Accordingly, in an exemplary screening assay of the present invention, the compound of interest is contacted with an isolated and purified *E. cloacae* polypeptide.

- 25 Screening assays can be constructed *in vitro* with a purified *E. cloacae* polypeptide or fragment thereof, such as an *E. cloacae* polypeptide having enzymatic

activity, such that the activity of the polypeptide produces a detectable reaction product. The efficacy of the compound can be assessed by generating dose response curves from data obtained using various concentrations of the test compound.

Moreover, a control assay can also be performed to provide a baseline for comparison.

- 5 Suitable products include those with distinctive absorption, fluorescence, or chemiluminescence properties, for example, because detection may be easily automated. A variety of synthetic or naturally occurring compounds can be tested in the assay to identify those which inhibit or potentiate the activity of the *E. cloacae* polypeptide. Some of these active compounds may directly, or with chemical alterations to promote
- 10 membrane permeability or solubility, also inhibit or potentiate the same activity (e.g., enzymatic activity) in whole, live *E. cloacae* cells.

### Overexpression Assays

- Overexpression assays are based on the premise that overproduction of a
- 15 protein would lead to a higher level of resistance to compounds that selectively interfere with the function of that protein. Overexpression assays may be used to identify compounds that interfere with the function of virtually any type of protein, including without limitation enzymes, receptors, DNA- or RNA-binding proteins, or any proteins that are directly or indirectly involved in regulating cell growth.
- 20 Typically, two bacterial strains are constructed. One contains a single copy of the gene of interest, and a second contains several copies of the same gene. Identification of useful inhibitory compounds of this type of assay is based on a comparison of the activity of a test compound in inhibiting growth and/or viability of the two strains. The method involves constructing a nucleic acid vector that directs
- 25 high level expression of a particular target nucleic acid. The vectors are then transformed into host cells in single or multiple copies to produce strains that express

low to moderate and high levels of protein encoding by the target sequence (strain A and B, respectively). Nucleic acid comprising sequences encoding the target gene can, of course, be directly integrated into the host cell.

Large numbers of compounds (or crude substances which may contain active compounds) are screened for their effect on the growth of the two strains. Agents which interfere with an unrelated target equally inhibit the growth of both strains. Agents which interfere with the function of the target at high concentration should inhibit the growth of both strains. It should be possible, however, to titrate out the inhibitory effect of the compound in the overexpressing strain. That is, if the compound is affecting the particular target that is being tested, it should be possible to inhibit the growth of strain A at a concentration of the compound that allows strain B to grow.

Alternatively, a bacterial strain is constructed that contains the gene of interest under the control of an inducible promoter. Identification of useful inhibitory agents using this type of assay is based on a comparison of the activity of a test compound in inhibiting growth and/or viability of this strain under both inducing and non-inducing conditions. The method involves constructing a nucleic acid vector that directs high-level expression of a particular target nucleic acid. The vector is then transformed into host cells that are grown under both non-inducing and inducing conditions (conditions A and B, respectively).

Large numbers of compounds (or crude substances which may contain active compounds) are screened for their effect on growth under these two conditions. Agents that interfere with the function of the target should inhibit growth under both conditions. It should be possible, however, to titrate out the inhibitory effect of the compound in the overexpressing strain. That is, if the compound is affecting the

particular target that is being tested, it should be possible to inhibit growth under condition A at a concentration that allows the strain to grow under condition B.

### Ligand-binding Assays

5           Many of the targets according to the invention have functions that have not yet been identified. Ligand-binding assays are useful to identify inhibitor compounds that interfere with the function of a particular target, even when that function is unknown. These assays are designed to detect binding of test compounds to particular targets. The detection may involve direct measurement of binding. Alternatively, indirect  
10 indications of binding may involve stabilization of protein structure or disruption of a biological function. Non-limiting examples of useful ligand-binding assays are detailed below.

A useful method for the detection and isolation of binding proteins is the Biomolecular Interaction Assay (BIAcore) system developed by Pharmacia Biosensor  
15 and described in the manufacturer's protocol (LKB Pharmacia, Sweden). The BIAcore system uses an affinity purified anti-GST antibody to immobilize GST-fusion proteins onto a sensor chip. The sensor utilizes surface plasmon resonance which is an optical phenomenon that detects changes in refractive indices. In accordance with the practice of the invention, a protein of interest is coated onto a chip and test  
20 compounds are passed over the chip. Binding is detected by a change in the refractive index (surface plasmon resonance).

A different type of ligand-binding assay involves scintillation proximity assays (SPA, described in U.S. Patent No. 4,568,649).

Another type of ligand binding assay, also undergoing development, is based  
25 on the fact that proteins containing mitochondrial targeting signals are imported into isolated mitochondria *in vitro* (Hurt *et al.*, 1985, *Embo J.* 4:2061-2068; Eilers and

Schatz, *Nature*, 1986, 322:228-231). In a mitochondrial import assay, expression vectors are constructed in which nucleic acids encoding particular target proteins are inserted downstream of sequences encoding mitochondrial import signals. The chimeric proteins are synthesized and tested for their ability to be imported into  
5 isolated mitochondria in the absence and presence of test compounds. A test compound that binds to the target protein should inhibit its uptake into isolated mitochondria *in vitro*.

Another ligand-binding assay is the yeast two-hybrid system (Fields and Song, 1989, *Nature* 340:245-246). The yeast two-hybrid system takes advantage of the  
10 properties of the GAL4 protein of the yeast *Saccharomyces cerevisiae*. The GAL4 protein is a transcriptional activator required for the expression of genes encoding enzymes of galactose utilization. This protein consists of two separable and functionally essential domains: an N-terminal domain which binds to specific DNA sequences (UAS<sub>G</sub>); and a C-terminal domain containing acidic regions, which is  
15 necessary to activate transcription. The native GAL4 protein, containing both domains, is a potent activator of transcription when yeast are grown on galactose media. The N-terminal domain binds to DNA in a sequence-specific manner but is unable to activate transcription. The C-terminal domain contains the activating regions but cannot activate transcription because it fails to be localized to UAS<sub>G</sub>. In  
20 the two-hybrid system, a system of two hybrid proteins containing parts of GAL4: (1) a GAL4 DNA-binding domain fused to a protein 'X' and (2) a GAL4 activation region, fused to a protein 'Y'. If X and Y can form a protein-protein complex and reconstitute proximity of the GAL4 domains, transcription of a gene regulated by UAS<sub>G</sub> occurs. Creation of two hybrid proteins, each containing one of the interacting proteins X and  
25 Y, allows the activation region of UAS<sub>G</sub> to be brought to its normal site of action.



The binding assay described in Fodor *et al.*, 1991, *Science* 251:767-773, which involves testing the binding affinity of test compounds for a plurality of defined polymers synthesized on a solid substrate, may also be useful.

Compounds which bind to the polypeptides of the invention are potentially  
5 useful as antibacterial agents for use in therapeutic compositions.

Pharmaceutical formulations suitable for antibacterial therapy comprise the antibacterial agent in conjunction with one or more biologically acceptable carriers. Suitable biologically acceptable carriers include, but are not limited to, phosphate-buffered saline, saline, deionized water, or the like. Preferred biologically acceptable  
10 carriers are physiologically or pharmaceutically acceptable carriers.

The antibacterial compositions include an antibacterial effective amount of active agent. Antibacterial effective amounts are those quantities of the antibacterial agents of the present invention that afford prophylactic protection against bacterial infections or which result in amelioration or cure of an existing bacterial infection.  
15 This antibacterial effective amount will depend upon the agent, the location and nature of the infection, and the particular host. The amount can be determined by experimentation known in the art, such as by establishing a matrix of dosages and frequencies and comparing a group of experimental units or subjects to each point in the matrix.

20 The antibacterial active agents or compositions can be formed into dosage unit forms, such as for example, creams, ointments, lotions, powders, liquids, tablets, capsules, suppositories, sprays, aerosols or the like. If the antibacterial composition is formulated into a dosage unit form, the dosage unit form may contain an antibacterial effective amount of active agent. Alternatively, the dosage unit form may include less  
25 than such an amount if multiple dosage unit forms or multiple dosages are to be used to administer a total dosage of the active agent. Dosage unit forms can include, in

addition, one or more excipient(s), diluent(s), disintegrant(s), lubricant(s), plasticizer(s), colorant(s), dosage vehicle(s), absorption enhancer(s), stabilizer(s), bactericide(s), or the like.

For general information concerning formulations, see, e.g., Gilman et al. (eds.), 1990, *Goodman and Gilman's: The Pharmacological Basis of Therapeutics*, 8th ed., Pergamon Press; and *Remington's Pharmaceutical Sciences*, 17th ed., 1990, Mack Publishing Co., Easton, PA; Avis et al. (eds.), 1993, *Pharmaceutical Dosage Forms: Parenteral Medications*, Dekker, New York; Lieberman et al (eds.), 1990, *Pharmaceutical Dosage Forms: Disperse Systems*, Dekker, New York.

10       The antibacterial agents and compositions of the present invention are useful for preventing or treating *E. cloacae* infections. Infection prevention methods incorporate a prophylactically effective amount of an antibacterial agent or composition. A prophylactically effective amount is an amount effective to prevent *E. cloacae* infection and will depend upon the specific bacterial strain, the agent, and the  
15       host. These amounts can be determined experimentally by methods known in the art and as described above.

*E. cloacae* infection treatment methods incorporate a therapeutically effective amount of an antibacterial agent or composition. A therapeutically effective amount is an amount sufficient to ameliorate or eliminate the infection. The prophylactically  
20       and/or therapeutically effective amounts can be administered in one administration or over repeated administrations. Therapeutic administration can be followed by prophylactic administration, once the initial bacterial infection has been resolved.

The antibacterial agents and compositions can be administered topically or systemically. Topical application is typically achieved by administration of creams,  
25       ointments, lotions, or sprays as described above. Systemic administration includes both oral and parental routes. Parental routes include, without limitation,

subcutaneous, intramuscular, intraperitoneal, intravenous, transdermal, inhalation and intranasal administration.

### EXEMPLIFICATION

5

#### Cloning and Sequencing *E. cloacae* Genomic Sequence

This invention provides nucleotide sequences of the genome of *E. cloacae* which thus comprises a DNA sequence library of *E. cloacae* genomic DNA. The detailed description that follows provides nucleotide sequences of *E. cloacae*, and also describes how the sequences were obtained and how ORFs (Open Reading Frames) and protein-coding sequences can be identified. Also described are methods of using the disclosed *E. cloacae* sequences in methods including diagnostic and therapeutic applications. Furthermore, the library can be used as a database for identification and comparison of medically important sequences in this and other strains of *E. cloacae* as well as other species of *Enterobacter*.

Chromosomal DNA from strain 15842 of *E. cloacae* was isolated after Zymolyase digestion, sodium dodecyl sulfate lysis, potassium acetate precipitation, phenol:chloroform extraction and ethanol precipitation (Soll, D.R., T. Srikantha and S.R. Lockhart: Characterizing Developmentally Regulated Genes in *E. cloacae*. In Microbial Genome Methods. K.W. Adolph, editor. CRC Press. New York. p 17-37.). Genomic *E. cloacae* DNA was hydrodynamically sheared in an HPLC and then separated on a standard 1% agarose gel. Fractions corresponding to 2500-3000 bp in length were excised from the gel and purified by the GeneClean procedure (Bio101, Inc.).

The purified DNA fragments were then blunt-ended using T4 DNA polymerase. The healed DNA was then ligated to unique *Bst*XI-linker adapters (5'-GTCTTCACCA<sup>(SEQ ID NO: 11325)</sup>CGGGG-3' and 5'-GTGGTGAAGAC<sup>(SEQ ID NO: 11326)</sup>-3' in 100-1000 fold molar

excess). These linkers are complimentary to the *Bst*XI-cut pGTC vector, while the overhang is not self-complimentary. Therefore, the linkers will not concatermerize nor will the cut-vector religate itself easily. The linker-adapted inserts were separated from the unincorporated linkers on a 1% agarose gel and purified using GeneClean. The linker-adapted inserts were then ligated to *Bst*XI-cut vector to construct a "shotgun" subclone libraries.

Only major modifications to the protocols are highlighted. Briefly, the library was then transformed into DH5 $\alpha$  competent cells (Gibco/BRL, DH5 $\alpha$  transformation protocol). It was assessed by plating onto antibiotic plates containing ampicillin and IPTG/Xgal. The plates were incubated overnight at 37°C. Transformants were then used for plating of clones and picking for sequencing. The cultures were grown overnight at 37°C. DNA was purified using a silica bead DNA preparation (Engelstein, 1996) method. In this manner, 25  $\mu$ g of DNA was obtained per clone.

These purified DNA samples were then sequenced using primarily ABI dye-terminator chemistry. All subsequent steps were based on sequencing by ABI377 automated DNA sequencing methods. The ABI dye terminator sequence reads were run on ABI377 machines and the data was transferred to UNIX machines following lane tracking of the gels. Base calls and quality scores were determined using the program PHRED (Ewing et al., 1998, Genome Res. 8: 175-185; Ewing and Green, 1998, Genome Res. 8: 685-734). Reads were assembled using PHRAP (P. Green, Abstracts of DOE Human Genome Program Contractor-Grantee Workshop V, Jan. 1996, p.157) with default program parameters and quality scores. The initial assembly was done at 6-fold coverage and yielded 513 contigs.

Finishing can follow the initial assembly. Missing mates (sequences from clones that only gave reads from one end of the *Enterobacter* DNA inserted in the plasmid) can be identified and sequenced with ABI technology to allow the identification of additional overlapping contigs.

5                   End-sequencing of randomly picked genomic lambda was also performed. Sequencing on a both sides was done for all lambda sequences. The lambda library backbone helped to verify the integrity of the assembly and allowed closure of some of the physical gaps. Primers for walking off the ends of contigs would be selected using pick\_primer (a GTC program) near the ends of the clones to  
10 facilitate gap closure. These walks can be sequenced using the selected clones and primers. These data are then reassembled with PHRAP. Additional sequencing using PCR-generated templates and screened and/or unscreened lambda templates can be done in addition.

To identify *E. cloacae* polypeptides the complete genomic sequence of *E.*  
15 *cloacae* were analyzed essentially as follows: First, all possible stop-to- stop open reading frames (ORFs) greater than 180 nucleotides in all six reading frames were translated into amino acid sequences. Second, the identified ORFs were analyzed for homology to known (archeabacter, prokaryotic and eukaryotic) protein sequences. Third, the coding potential of non-homologous sequences were evaluated with the  
20 program GENEMARKTM (Borodovsky and McIninch, 1993, Comp. Chem. 17:123).

#### Identification, Cloning and Expression of *E.cloacae* Nucleic Acids

Expression and purification of the *E.cloacae* polypeptides of the invention can  
25 be performed essentially as outlined below.

To facilitate the cloning, expression and purification of membrane and secreted proteins from *E.cloacae*, a gene expression system, such as the pET System (Novagen), for cloning and expression of recombinant proteins in *E. coli*, is selected. Also, a DNA sequence encoding a peptide tag, the His-Tag, is fused to the 3' end of DNA sequences of interest in order to facilitate purification of the recombinant protein products. The 3' end is selected for fusion in order to avoid alteration of any 5' terminal signal sequence.

#### PCR Amplification and Cloning of Nucleic Acids Containing ORF's Encoding

#### 10 Enzymes

Nucleic acids chosen (for example, from the nucleic acids set forth in SEQ ID NO: 1 - SEQ ID NO: 5662 for cloning from the 15842 strain of *E.cloacae* are prepared for amplification cloning by polymerase chain reaction (PCR). Synthetic oligonucleotide primers specific for the 5' and 3' ends of open reading frames (ORFs) are designed and purchased from GibcoBRL Life Technologies (Gaithersburg, MD, USA). All forward primers (specific for the 5' end of the sequence) are designed to include an NcoI cloning site at the extreme 5' terminus. These primers are designed to permit initiation of protein translation at a methionine residue followed by a valine residue and the coding sequence for the remainder of the native *E.cloacae* DNA sequence. All reverse primers (specific for the 3' end of any *E.cloacae* ORF) include a EcoRI site at the extreme 5' terminus to permit cloning of each *E.cloacae* sequence into the reading frame of the pET-28b. The pET-28b vector provides sequence encoding an additional 20 carboxy-terminal amino acids including six histidine residues (at the extreme C-terminus), which comprise the His-Tag.

Genomic DNA prepared from the 15842 strain of *E.cloacae* is used as the source of template DNA for PCR amplification reactions (Current Protocols in Molecular Biology, John Wiley and Sons, Inc., F. Ausubel et al., eds., 1994). To amplify a DNA sequence containing an *E.cloacae* ORF, genomic DNA (50  
5 nanograms) is introduced into a reaction vial containing 2 mM MgCl<sub>2</sub>, 1 micromolar synthetic oligonucleotide primers (forward and reverse primers) complementary to and flanking a defined *E.cloacae* ORF, 0.2 mM of each deoxynucleotide triphosphate; dATP, dGTP, dCTP, dTTP and 2.5 units of heat stable DNA polymerase (Amplitaq, Roche Molecular Systems, Inc., Branchburg, NJ, USA) in a final volume of 100  
10 microliters.

Upon completion of thermal cycling reactions, each sample of amplified DNA is washed and purified using the Qiaquick Spin PCR purification kit (Qiagen, Gaithersburg, MD, USA). All amplified DNA samples are subjected to digestion with the restriction endonucleases, e.g., NcoI and EcoRI (New England BioLabs, Beverly,  
15 MA, USA)(Current Protocols in Molecular Biology, John Wiley and Sons, Inc., F. Ausubel et al., eds., 1994). DNA samples are then subjected to electrophoresis on 1.0 % NuSeive (FMC BioProducts, Rockland, ME USA) agarose gels. DNA is visualized by exposure to ethidium bromide and long wave uv irradiation. DNA contained in slices isolated from the agarose gel is purified using the Bio 101 GeneClean Kit  
20 protocol (Bio 101 Vista, CA, USA).

#### Cloning of *E. cloacae* Nucleic Acids Into an Expression Vector

The pET-28b vector is prepared for cloning by digestion with restriction  
25 endonucleases, e.g., NcoI and EcoRI (Current Protocols in Molecular Biology, John Wiley and Sons, Inc., F. Ausubel et al., eds., 1994). The pET-28a vector, which

encodes a His-Tag that can be fused to the 5' end of an inserted gene, is prepared by digestion with appropriate restriction endonucleases.

Following digestion, DNA inserts are cloned (Current Protocols in Molecular Biology, John Wiley and Sons, Inc., F. Ausubel et al., eds., 1994) into the previously  
5 digested pET-28b expression vector. Products of the ligation reaction are then used to transform the BL21 strain of *E. coli* (Current Protocols in Molecular Biology, John Wiley and Sons, Inc., F. Ausubel et al., eds., 1994) as described below.

#### Transformation Of Competent Bacteria With Recombinant Plasmids

10

Competent bacteria, *E. coli* strain BL21 or *E. coli* strain BL21(DE3), are transformed with recombinant pET expression plasmids carrying the cloned *E. cloacae* sequences according to standard methods (Current Protocols in Molecular, John Wiley and Sons, Inc., F. Ausubel et al., eds., 1994). Briefly, 1 microliter of ligation reaction  
15 is mixed with 50 microliters of electrocompetent cells and subjected to a high voltage pulse, after which, samples are incubated in 0.45 milliliters SOC medium (0.5% yeast extract, 2.0 % tryptone, 10 mM NaCl, 2.5 mM KCl, 10 mM MgCl<sub>2</sub>, 10 mM MgSO<sub>4</sub> and 20, mM glucose) at 37°C with shaking for 1 hour. Samples are then spread on LB agar plates containing 25 microgram/ml kanamycin sulfate for growth overnight.  
20 Transformed colonies of BL21 are then picked and analyzed to evaluate cloned inserts as described below.

#### Identification Of Recombinant Expression Vectors With *E. cloacae* Nucleic Acids

25

Individual BL21 clones transformed with recombinant pET-28b *E. cloacae* ORFs are analyzed by PCR amplification of the cloned inserts using the same forward



and reverse primers, specific for each *E. cloacae* sequence, that were used in the original PCR amplification cloning reactions. Successful amplification verifies the integration of the *E. cloacae* sequences in the expression vector (Current Protocols in Molecular Biology, John Wiley and Sons, Inc., F. Ausubel et al., eds., 1994).

5

#### Isolation and Preparation of Nucleic Acids From Transformants

Individual clones of recombinant pET-28b vectors carrying properly cloned *E. cloacae* ORFs are picked and incubated in 5 mls of LB broth plus 25 microgram/ml  
10 kanamycin sulfate overnight. The following day plasmid DNA is isolated and purified using the Qiagen plasmid purification protocol (Qiagen Inc., Chatsworth, CA, USA).

#### Expression Of Recombinant *E. cloacae* Sequences In *E. coli*

15 The pET vector can be propagated in any *E. coli* K-12 strain e.g. HMS174, HB101, JM109, DH5, etc. for the purpose of cloning or plasmid preparation. Hosts for expression include *E. coli* strains containing a chromosomal copy of the gene for T7 RNA polymerase. These hosts are lysogens of bacteriophage DE3, a lambda  
20 derivative that carries the lacI gene, the lacUV5 promoter and the gene for T7 RNA polymerase. T7 RNA polymerase is induced by addition of isopropyl-B-D-thiogalactoside (IPTG), and the T7 RNA polymerase transcribes any target plasmid, such as pET-28b, carrying its gene of interest. Strains used include: BL21(DE3) (Studier, F.W., Rosenberg, A.H., Dunn, J.J., and Dubendorff, J.W. (1990) Meth. Enzymol. 185, 60-89).

25 To express recombinant *E. cloacae* sequences, 50 nanograms of plasmid DNA isolated as described above is used to transform competent BL21(DE3) bacteria as

described above (provided by Novagen as part of the pET expression system kit).

The lacZ gene (beta-galactosidase) is expressed in the pET-System as described for the *E. cloacae* recombinant constructions. Transformed cells are cultured in SOC medium for 1 hour, and the culture is then plated on LB plates containing 25

5 micrograms/ml kanamycin sulfate. The following day, bacterial colonies are pooled and grown in LB medium containing kanamycin sulfate (25 micrograms/ml) to an optical density at 600 nm of 0.5 to 1.0 O.D. units, at which point, 1 millimolar IPTG was added to the culture for 3 hours to induce gene expression of the *E. cloacae* recombinant DNA constructions .

10 After induction of gene expression with IPTG, bacteria are pelleted by centrifugation in a Sorvall RC-3B centrifuge at 3500 x g for 15 minutes at 4°C. Pellets are resuspended in 50 milliliters of cold 10 mM Tris-HCl, pH 8.0, 0.1 M NaCl and 0.1 mM EDTA (STE buffer). Cells are then centrifuged at 2000 x g for 20 min at 4°C. Wet pellets are weighed and frozen at -80°C until ready for protein  
15 purification.

A variety of methodologies known in the art can be utilized to purify the isolated proteins. (Current Protocols in Protein Science, John Wiley and Sons, Inc., J. E. Coligan et al., eds., 1995). For example, the frozen cells may be thawed, resuspended in buffer and ruptured by several passages through a small volume  
20 microfluidizer (Model M-110S, Microfluidics International Corporation, Newton, MA). The resultant homogenate may be centrifuged to yield a clear supernatant (crude extract) and following filtration the crude extract may be fractionated over columns. Fractions may be monitored by absorbance at OD<sub>280</sub> nm. and peak fractions may analyzed by SDS-PAGE

25 The concentrations of purified protein preparations may be quantified spectrophotometrically using absorbance coefficients calculated from amino acid

content (Perkins, S.J. 1986 Eur. J. Biochem. 157, 169-180). Protein concentrations are also measured by the method of Bradford, M.M. (1976) Anal. Biochem. 72, 248-254, and Lowry, O.H., Rosebrough, N., Farr, A.L. & Randall, R.J. (1951) J. Biol. Chem. 193, pages 265-275, using bovine serum albumin as a standard.

- 5        SDS-polyacrylamide gels of various concentrations may be purchased from BioRad (Hercules, CA, USA), and stained with Coomassie blue. Molecular weight markers may include rabbit skeletal muscle myosin (200 kDa), *E. coli* (-galactosidase (116 kDa), rabbit muscle phosphorylase B (97.4 kDa), bovine serum albumin (66.2 kDa), ovalbumin (45 kDa), bovine carbonic anhydrase (31 kDa), soybean trypsin
- 10   inhibitor (21.5 kDa), egg white lysozyme (14.4 kDa) and bovine aprotinin (6.5 kDa).

## EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more  
5 than routine experimentation, many equivalents to the specific embodiments and  
methods described herein. The specific embodiments described herein are offered by  
way of example only, and the invention is to limited only by the terms of the  
appended claims, along with the full scope of equivalents to which such claims are  
entitled.

10

TABLE 2

Contig	Orf	nt ID	aa ID	nt Length	aa Length	Blast Score	Blast Probability	Subject Taxonomy	Subject Name	Subject Description
CONTIG1	31892515_f2_2	1	5663	432	144	264	6.2(10)-23	Salmonella typhimurium	U94729	[de:salmonella typhimurium oxd-6 operon, putative substrate-binding protein (oxd-6a), putative transmembrane protein (oxd-6), putative transmembrane protein (oxd-6c), putative atpase (oxd-6d), and putative atpase (oxd-6e)]
CONTIG1	32542792_c2_6	2	5664	231	77	92	0.0016	Plasmodium falciparum	P09346	knob-associated histidine-rich protein precursor (kahrp)
CONTIG10	22066424_c1_4	3	5665	490	163	262	3.2(10)-22	Escherichia coli	b1006	[pn:hypothetical protein] [gn:yodg]
CONTIG100	30752167_c1_6	4	5666	267	89	390	2.7(10)-36	Escherichia coli	b4162	[pn:hypothetical 23.5 kd protein in psd-amib intergenic region] [gn:yjer]
CONTIG100	15735381_c3_7	5	5667	393	131	542	2.2(10)-52	Escherichia coli	b4162	[pn:hypothetical 23.5 kd protein in psd-amib intergenic region] [gn:yjer]
CONTIG101	3222031_f1_1	6	5668	675	225	797	2.1(10)-79	Escherichia coli	b1047	[pn:hypothetical protein]
CONTIG101	6261469_c2_6	7	5669	647	215	621	9.3(10)-61	Escherichia coli	b1045	[pn:hypothetical protein]
CONTIG102	4535155_c2_13	8	5670	924	308	1137	1.8(10)-115	Escherichia coli	b0433	[pn:ampg protein] [gn:ampg]
CONTIG103	19719827_f1_1	9	5671	1005	335	120	0.00079	Human herpesvirus 6 (strain Uganda-1102)	AF015297	[de:human herpesvirus 6 (strain uganda-1102) ie2hom mrna, complete cds.] [pn:ie2hom] [gn:ie2hom] [nt:similar to the immediate-early 2 protein of human]
CONTIG105	24245287_c1_5	10	5672	1170	390	1142	5.7(10)-116	Escherichia coli	b4261	[pn:hypothetical 40.4 kd protein in pepa-gntv intergenic region] [gn:yjgp]
CONTIG106	15104766_f3_7	11	5673	756	252	518	7.7(10)-50	Escherichia coli	P06615	resolvase (protein d)
CONTIG108	35269762_f2_2	12	5674	951	317	131	4.0(10)-6	Bacillus subtilis	yojN	[pn:hypothetical protein]

CONTIG109	7240778_f2_2	13	5675	519	173	224	1.1(10)-18	Haemophilus influenzae	HI1415	[pn:hypothetical protein]
CONTIG11	29877090_c2_8	14	5676	345	115	191	1.2(10)-14	Cloning vector pCMVLacI	U64448	or:cloning vector pcmlaci pn:lac repressor gn:laci le:2685 re:3803 di:direct
CONTIG11	3328900_c3_9	15	5677	342	114	203	1.8(10)-16	Haemophilus influenzae	HI0522	[pn:sp]
CONTIG110	4084392_f2_1	16	5678	1110	370	154	2.1(10)-8	Escherichia coli	P07620	plasmid partition par a protein.
CONTIG111	25803317_f2_3	17	5679	1056	352	1446	3.5(10)-148	Escherichia coli	b2392	[pn:hypothetical protein]
CONTIG113	34192257_f2_1	18	5680	1071	357	805	3.0(10)-80	Escherichia coli	b3589	[pn:hypothetical 40.2 kd protein in avta-selb intergenic region] [gn:yiaY]
CONTIG115	26750793_f3_3	19	5681	537	179	799	1.3(10)-79	Escherichia coli	b3279	[pn:hypothetical protein] [gn:yrdA]
CONTIG115	16175299_c1_5	20	5682	199	66	275	4.2(10)-24	Escherichia coli	b3281	[pn:shikimate dehydrogenase] [gn:aroE]
CONTIG115	22938215_c3_7	21	5683	357	119	311	6.5(10)-28	Escherichia coli	b3280	[pn:hypothetical 10.0 kd protein in rmd-aroE intergenic region]
CONTIG116	19713405_c3_8	22	5684	198	66	180	5.0(10)-14	Escherichia coli	b1972	[pn:hypothetical protein]
CONTIG116	24033587_c3_9	23	5685	522	174	479	1.0(10)-45	Haemophilus influenzae	HI0970	[pn:3-dehydroquinate dehydratase] [gn:aroQ]
CONTIG117	13852211_f1_1	24	5686	591	197	647	1.6(10)-63	Escherichia coli	b1988	[pn:nitrogen assimilation regulatory protein] [gn:nac]
CONTIG117	21690876_f2_2	25	5687	240	80	224	1.1(10)-18	Escherichia coli	b1990	[pn:31.6 kd protein in cobT 3' region precursor] [gn:erfK]
CONTIG117	29942556_f2_4	26	5688	342	114	250	1.8(10)-21	Escherichia coli	b1988	[pn:nitrogen assimilation regulatory protein] [gn:nac]
CONTIG118	14225431_c1_13	27	5689	492	164	558	4.4(10)-54	Erwinia carotovora subsp. carotovora	JC4729	mob protein c - erwinia carotovora subsp. carotovora a cis-acting locus, orit and trans-acting locus, mob are involved in mobilization of pec3, a non-self-transmissible multiple-copy plasmid. the mob consists of five proteins.
CONTIG118	5163340_c2_16	28	5690	696	232	938	2.3(10)-94	Plasmid pSW200	L42525	or:plasmid psw200 gn:moba le:1657 re:3156 di:direct sr:plasmid psw200 dna

CONTIG119	36072216_c2_6	29	5691	1392	464	2041	3.1(10)-211	Escherichia coli	b0930	[pn:asparaginyl-trna synthetase] [gn:asns]
CONTIG120	2394707_f1_1	30	5692	567	189	629	1.3(10)-61	Escherichia coli	b3983	[pn:50s ribosomal subunit protein 111] [gn:rplk]
CONTIG120	23572188_f1_2	31	5693	708	236	959	1.3(10)-96	Escherichia coli	b3984	[pn:50s ribosomal subunit protein 11] [gn:rpla]
CONTIG120	31651642_f2_4	32	5694	315	105	295	3.2(10)-26	Escherichia coli	b3985	[pn:50s ribosomal subunit protein 110] [gn:rplj]
CONTIG121	23552216_f2_1	33	5695	384	128	612	8.4(10)-60	Escherichia coli	b3310	[pn:50s ribosomal subunit protein 114] [gn:rpln]
CONTIG121	20117711_f2_2	34	5696	600	200	779	1.7(10)-77	Escherichia coli	b3308	[pn:50s ribosomal subunit protein 15] [gn:rple]
CONTIG121	2037562_f3_3	35	5697	357	119	512	3.2(10)-49	Escherichia coli	b3309	[pn:50s ribosomal subunit protein 124] [gn:rplx]
CONTIG122	16411425_f1_2	36	5698	1055	352	1187	9.8(10)-121	Escherichia coli	b2965	[pn:ornithine decarboxylase, constitutive] [gn:spec]
CONTIG122	24229836_c2_9	37	5699	531	177	571	1.8(10)-55	Escherichia coli	b2966	[pn:hypothetical protein] [gn:yqga]
CONTIG123	34484627_f2_2	38	5700	516	172	188	7.0(10)-15	Escherichia coli	b3335	[pn:type 4 prepilin-like protein specific leader peptidase] [gn:hofd]
CONTIG123	32032906_c2_6	39	5701	519	173	732	1.6(10)-72	Escherichia coli	b3336	[pn:bacterioferritin] [gn:bfr]
CONTIG125	24335430_c2_7	40	5702	1173	391	1749	2.7(10)-180	Escherichia coli	b0094	[pn:cell division protein ftsa] [gn:fts]
CONTIG125	10020833_c3_8	41	5703	255	85	309	1.1(10)-27	Escherichia coli	b0093	[pn:cell division protein ftsq] [gn:fts]
CONTIG126	26582552_f1_1	42	5704	657	219	1051	2.5(10)-106	Escherichia coli	b3229	[pn:stringent starvation protein] [gn:sspa]
CONTIG126	31338512_f3_4	43	5705	216	72	243	1.1(10)-20	Escherichia coli	b3228	[pn:stringent starvation protein b] [gn:sspb]
CONTIG127	4891510_f1_1	44	5706	333	111	387	5.7(10)-36	Escherichia coli	b1060	[pn:hypothetical protein]
CONTIG127	33255311_f2_4	45	5707	699	233	774	5.7(10)-77	Escherichia coli	b1059	[pn:hypothetical protein] [gn:sola]
CONTIG128	25781411_f2_1	46	5708	474	158	646	2.1(10)-63	Escherichia coli	b3342	[pn:30s ribosomal subunit protein s12] [gn:rpsl]
CONTIG128	23714561_f2_2	47	5709	411	137	637	1.8(10)-62	Escherichia coli	b3341	[pn:30s ribosomal subunit protein s7] [gn:rpsg]

CONTIG128	46907_c3_4	48	5710	339	113	158	1.1(10)-11	Eikenella corrodens	P35648	hemagglutinin 2.
CONTIG129	16178312_fl_2	49	5711	555	185	706	9.9(10)-70	Escherichia coli	P08504	transposase for transposon tn2501.
CONTIG129	22386375_c1_6	50	5712	636	212	691	3.6(10)-68	Escherichia coli	b1374	[pn:hypothetical protein]
CONTIG130	16016405_fl_1	51	5713	402	134	349	6.2(10)-32	Escherichia coli	b0339	[pn:cyanate permease] [gn:cyn]
CONTIG130	1150468_c2_7	52	5714	909	303	970	9.6(10)-98	Escherichia coli	b1668	[pn:hypothetical protein]
CONTIG131	395662_fl_2	53	5715	380	127	208	5.4(10)-17	Haemophilus influenzae	HI1053	[pn:gb]
CONTIG131	35806526_fl_3	54	5716	357	119	91	0.00013	Mycoplasma genitalium	MG427	[pn:hypothetical protein mg427]
CONTIG131	23601436_fl_4	55	5717	624	208	134	2.7(10)-8	Escherichia coli	b1649	[pn:hypothetical protein]
CONTIG132	22667830_fl_1	56	5718	786	262	331	5.0(10)-30	Bacillus subtilis	pdhD	[pn:diacylglycerol kinase dehydrogenase e3 subunit of both pyruvate dehydrogenase and 2- oxoglutarate dehydrogenase]
CONTIG132	34410768_fl_2	57	5719	249	83	301	7.5(10)-27	Escherichia coli	b1994	[pn:insertion element is5 hypothetical 39.3 kd protein]
CONTIG134	10651717_c1_6	58	5720	498	166	777	2.7(10)-77	Plasmid pSW200	L42525	or:plasmid psw200 gn:mob le:2345 re:2830 di:direct sr:plasmid psw200 dna
CONTIG134	26681536_c1_7	59	5721	219	73	328	1.0(10)-29	Erwinia carotovora subsp. carotovora	JC4730	mob protein d - erwinia carotovora subsp. carotovora a cis-acting locus, orit and trans-acting locus, mob are involved in mobilization of pec3, a non-self-transmissible multiple-copy plasmid. the mob consists of five proteins.
CONTIG134	12897806_c3_9	60	5722	861	287	1188	7.7(10)-121	Plasmid pSW200	L42525	or:plasmid psw200 gn:mob le:1657 re:3156 di:direct sr:plasmid psw200 dna
CONTIG135	964590_fl_2	61	5723	627	209	824	2.8(10)-82	Escherichia coli	b0434	[pn:hypothetical lipoprotein in ampg 5' region] [gn:yaig]
CONTIG135	30722826_fl_3	62	5724	372	124	458	1.7(10)-43	Escherichia coli	b0433	[pn:ampg protein] [gn:ampg]



CONTIG135	34626891_c1_5	63	5725	237	79	316	1.8(10)-28	Escherichia coli	b0435	[pn:bola protein] [gn:bola]
CONTIG136	5954207_f1_1	64	5726	765	255	828	1.1(10)-82	Escherichia coli	b2511	[pn:hypothetical protein]
CONTIG136	32500050_f3_3	65	5727	606	202	891	2.2(10)-89	Escherichia coli	b2512	[pn:hypothetical protein]
CONTIG137	14506503_f1_1	66	5728	462	154	166	2.8(10)-12	Escherichia coli	b4396	[pn:right origin-binding protein] [gn:rob]
CONTIG137	30667937_f3_3	67	5729	192	64	121	2.8(10)-7	Escherichia coli	b1790	[pn:hypothetical protein]
CONTIG137	1189430_f3_4	68	5730	360	120	100	1.5(10)-5	Escherichia coli	b1112	[pn:hypothetical protein]
CONTIG137	14743832_c3_6	69	5731	216	72	228	4.0(10)-19	Escherichia coli	b2106	[pn:hypothetical protein]
CONTIG138	25392331_f1_1	70	5732	312	104	378	5.2(10)-35	Escherichia coli	b1852	[pn:glucose 6-phosphate 1 dehydrogenase] [gn:zwf]
CONTIG138	16695877_f3_6	71	5733	885	295	1346	1.3(10)-137	Escherichia coli	b1851	[pn:phosphogluconate dehydratase] [gn:edd]
CONTIG139	32314061_f2_4	72	5734	387	129	130	1.0(10)-8	Escherichia coli	b1569	[pn:repressor protein of division inhibition gene dicb] [gn:dicc]
CONTIG139	14470637_c2_8	73	5735	423	141	401	1.8(10)-37	Escherichia coli	b1570	[pn:repressor protein of division inhibition gene dicb] [gn:dica]
CONTIG140	35949137_f3_4	74	5736	1260	420	211	1.5(10)-14	Pseudomonas putida	S27611	agglutination protein - pseudomonas putida
CONTIG142	35835092_f1_1	75	5737	1197	399	1652	5.2(10)-170	Escherichia coli	b3544	[pn:periplasmic dipeptide transport protein precursor] [gn:dppa]
CONTIG144	16054202_f1_4	76	5738	336	112	528	6.7(10)-51	Escherichia coli	b1827	[pn:hypothetical protein]
CONTIG144	10314637_c2_10	77	5739	987	329	1049	4.0(10)-106	Escherichia coli	b1828	[pn:hypothetical protein]
CONTIG145	33239137_f2_2	78	5740	711	237	1012	3.3(10)-102	Escherichia coli	b0571	[pn:hypothetical protein] [gn:yica]
CONTIG145	11117780_f3_4	79	5741	342	114	242	6.0(10)-20	Escherichia coli	b0570	[pn:hypothetical protein] [gn:ybcz]
CONTIG145	35569027_c3_10	80	5742	603	201	287	4.5(10)-25	Escherichia coli	b0572	[pn:hypothetical protein] [gn:yicb]
CONTIG146	3906875_f3_4	81	5743	1296	432	1799	1.3(10)-185	Escherichia coli	b3702	[pn:chromosomal replication initiator protein dnaa] [gn:dnaa]

CONTIG147	32610875_f3_3	82	5744	288	96	278	2.1(10)-24	Escherichia coli	b1916	[pn:regulatory protein] [gn:sdia]
CONTIG147	16413907_f3_5	83	5745	461	154	500	6.2(10)-48	Escherichia coli	b1914	[pn:23.9 kd protein in uvrc-sdia intergenic region] [gn:uvry]
CONTIG147	12614702_c1_7	84	5746	225	75	299	1.2(10)-26	Escherichia coli	b1915	[pn:yecf]
CONTIG148	22746011_f1_1	85	5747	366	122	199	4.9(10)-16	Escherichia coli	b1166	[pn:hypothetical protein]
CONTIG148	5250378_c2_7	86	5748	888	296	464	4.0(10)-44	Klebsiella pneumoniae	L23111	or:klebsiella pneumoniae pn:fimbrial adhesin gn:fimk le:1139 re:2380 di:direct sr:klebsiella pneumoniae
CONTIG149	3954682_c2_4	87	5749	558	186	555	9.1(10)-54	Bacteriophage lambda	A14086	repressor protein ci - phage lambda repressor protein ci allows phage lambda to reside inactively in the chromosome of its host bacterium. this lysogenic state is maintained by binding of regulatory protein ci to the or and ol operators, preventing
CONTIG15	36406937_f1_1	88	5750	467	156	168	9.9(10)-12	Escherichia coli	b3066	[pn:dna primase] [gn:dnag]
CONTIG15	9784375_f2_2	89	5751	330	110	113	8.3(10)-6	Haemophilus influenzae	H10532	[pn:dna primase] [gn:dnag]
CONTIG150	24485818_f1_1	90	5752	195	65	160	4.0(10)-11	Escherichia coli	b1184	[pn:umuc protein] [gn:umuc]
CONTIG150	24042892_c1_5	91	5753	258	86	104	0.00018	Methanococcus jannaschii	MJ1643	[pn:chromosome segretation protein]
CONTIG151	4485693_c1_4	92	5754	825	275	1160	7.0(10)-118	Escherichia coli	b0197	[pn:hypothetical 29.4 kd lipoprotein in rcsf-rmh intergenic region]
CONTIG152	32285024_c1_7	93	5755	931	310	1152	5.0(10)-117	Escherichia coli	A25937	arsenical pump-driving atpase (ec 3.6.1.-) - escherichia coli plasmid r773 this anion-transporting atpase catalyzes the extrusion of the oxyanions arsenite, antimonic, and arsenate, thus lowering the intracellular concentration of these toxic oxyanions.
CONTIG153	14258412_f1_1	94	5756	183	61	101	1.2(10)-5	Haemophilus influenzae	H10598	[pn:hypothetical protein]

CONTIG153	1461540_f1_2	95	5757	945	315	633	5.0(10)-62	Escherichia coli	b0626	[pn:hypothetical protein] [gn:ybem]
CONTIG153	14850461_c2_8	96	5758	396	132	588	2.8(10)-57	Escherichia coli	b0624	[pn:hypothetical 13.8 kd protein in cspe-lipa intergenic region] [gn:ybei]
CONTIG154	24645175_c1_1	97	5759	456	152	183	2.3(10)-14	Escherichia coli	S70162	,
CONTIG155	29394584_c2_9	98	5760	839	279	1123	5.9(10)-114	Escherichia coli	b0031	[pn:dihydrodipicolinate reductase] [gn:dapb]
CONTIG155	23632260_c3_10	99	5761	390	130	567	4.9(10)-55	Escherichia coli	b0032	[pn:carbamoyl-phosphate synthase small chain] [gn:cara]
CONTIG156	1384665_f2_1	100	5762	1269	423	1956	3.2(10)-202	Escherichia coli	b4260	[pn:aminopeptidase a/1] [gn:pepa]
CONTIG157	16303956_f2_2	101	5763	2103	701	2013	2.8(10)-208	Escherichia coli	b1102	[pn:outer-membrane receptor for fe] [gn:fhue]
CONTIG158	10563465_f3_3	102	5764	492	164	102	9.3(10)-6	Escherichia coli	b2861	[pn:insertion element is2 hypothetical 13.4 kd protein]
CONTIG159	12991252_f1_1	103	5765	726	242	313	4.0(10)-28	Escherichia coli	b0752	[pn:hypothetical protein] [gn:ybgr]
CONTIG159	23938415_f3_4	104	5766	390	130	102	9.3(10)-6	Bacillus subtilis	yoza	[pn:hypothetical protein]
CONTIG159	24660337_c3_8	105	5767	285	95	375	1.1(10)-34	Escherichia coli	b1994	[pn:insertion element is5 hypothetical 39.3 kd protein]
CONTIG16	4019806_c2_5	106	5768	435	145	108	2.1(10)-6	Escherichia coli	b1219	[pn:hypothetical 12.7 kd protein in chac-nar1 intergenic region] [gn:ychn]
CONTIG160	23728327_f1_1	107	5769	510	170	244	8.3(10)-21	Bacillus subtilis	glcR	[pn:transcriptional regulator] [gn:ywpi]
CONTIG160	35253527_f2_3	108	5770	1143	381	824	2.8(10)-82	Escherichia coli	b1624	[pn:hypothetical protein]
CONTIG161	12694182_c1_9	109	5771	762	254	426	4.2(10)-40	Escherichia coli	b0897	[pn:hypothetical 23.1 kd protein in dmcs 3''' region] [gn:ycac]
CONTIG161	26616641_c3_11	110	5772	582	194	222	1.8(10)-18	Homo sapiens	Y07867	or:homo sapiens pn:pirin le:205 re:1077 di:direct sr:human
CONTIG162	7282255_f1_1	111	5773	372	124	450	1.2(10)-42	Escherichia coli	b0946	[pn:hypothetical protein]
CONTIG162	21741657_c2_12	112	5774	972	324	1313	4.4(10)-134	Escherichia coli	b0947	[pn:hypothetical protein]
CONTIG163	22949037_c1_12	113	5775	792	264	1273	7.5(10)-130	Escherichia coli	b2568	[pn:signal peptidase i] [gn:lepb]

CONTIG163	31880432_c1_13	114	5776	531	177	776	3.5(10)-77	Escherichia coli	b2566	[pn:gtp-binding protein] [gn:era]
CONTIG163	12203387_c2_15	115	5777	801	267	1059	3.6(10)-107	Escherichia coli	b2567	[pn:ribonuclease iii] [gn:rmc]
CONTIG165	20508557_f1_1	116	5778	1307	436	1915	7.0(10)-198	Escherichia coli	b4208	[pn:d-serine/d-alanine/glycine transporter] [gn:cyca]
CONTIG165	26386588_f2_2	117	5779	636	212	821	6.0(10)-82	Escherichia coli	b4207	[pn:probable fbp-type 22 kd peptidyl-prolyl cis- trans isomerase]
CONTIG167	31367263_f1_2	118	5780	1200	400	452	7.5(10)-43	Escherichia coli	AF005044	[PN:TraU] [GN:traU] [DE:Escherichia coli plasmid R100-1 TraV (traV), TraR (traR), OrfG1(orfG1), OrfH (orfH), OrfI (orfI), TraC (traC), TrbI (trbI), TraW(traW), TraU (traU), TrbC (trbC), TraN (traN), TrbE (trbE) and TraF(traF) genes, c
CONTIG168	34615936_c1_7	119	5781	879	293	1005	1.8(10)-101	Escherichia coli	b4174	[pn:hflk protein] [gn:hflk]
CONTIG168	26601532_c2_8	120	5782	333	111	516	1.2(10)-49	Escherichia coli	b4173	[pn:gtp-binding protein hflx] [gn:hflx]
CONTIG169	21657693_f1_2	121	5783	597	199	707	7.2(10)-70	Escherichia coli	b3639	[pn:dfp protein] [gn:dfp]
CONTIG169	32246000_f2_5	122	5784	204	68	151	4.0(10)-10	Escherichia coli	b3639	[pn:dfp protein] [gn:dfp]
CONTIG169	5953150_c3_14	123	5785	852	284	801	7.7(10)-80	Escherichia coli	b3638	[pn:dna repair protein radc] [gn:radc]
CONTIG170	1204702_f1_1	124	5786	336	112	523	2.2(10)-50	Escherichia coli	b0969	[pn:hypothetical protein in held-sert intergenic region] [gn:ycek]
CONTIG170	31462776_f2_4	125	5787	663	221	634	3.8(10)-62	Escherichia coli	b0970	[pn:hypothetical 23.4 kd protein in sert 5'''region] [gn:ycca]
CONTIG170	1210765_c2_13	126	5788	282	94	307	1.7(10)-27	Escherichia coli	b0968	[pn:hypothetical protein]
CONTIG170	32428439_c3_17	127	5789	1152	384	965	3.2(10)-97	Escherichia coli	b1243	[pn:periplasmic oligopeptide-binding protein precursor] [gn:oppa]
CONTIG171	11737962_f3_4	128	5790	948	316	104	0.01	Streptococcus phage phi7201	U89246	[de:streptococcus phage phi7201 orf and orf unknown protein genes,partial cds.] [pn:unknown] [nt:orfy]

CONTIG173	26742882_f2_2	129	5791	555	185	144	1.5(10)-9	Pseudomonas sp.	P18896	increased glyphosate resistance protein.
CONTIG173	4164015_c3_8	130	5792	1047	349	1016	1.3(10)-102	Escherichia coli	b3001	[pn:hypothetical protein]
CONTIG174	5270268_f1_1	131	5793	1392	464	1809	1.2(10)-186	Escherichia coli	b0436	[pn:trigger factor] [gn:tig]
CONTIG175	22050143_f1_1	132	5794	1092	364	463	5.2(10)-44	Escherichia coli	b4287	[pn:ironiii dicitrate transport atp-binding protein fece] [gn:fece]
CONTIG175	4579201_f2_2	133	5795	558	186	304	3.6(10)-27	Bacillus subtilis	yvrB	[pn:hypothetical protein]
CONTIG175	35567462_c2_12	134	5796	402	134	324	8.5(10)-29	Escherichia coli	b1243	[pn:periplasmic oligopeptide-binding protein precursor] [gn:oppa]
CONTIG176	25938376_f2_2	135	5797	576	192	659	8.8(10)-65	Escherichia coli	b0199	[pn:abc] [gn:abc]
CONTIG176	23473562_f2_3	136	5798	483	161	767	3.1(10)-76	Escherichia coli	b0199	[pn:abc] [gn:abc]
CONTIG176	7270092_f3_8	137	5799	614	205	777	2.7(10)-77	Escherichia coli	b0198	[pn:hypothetical abc transporter permease protein yae] [gn:yae]
CONTIG176	29433308_c2_12	138	5800	630	210	892	1.8(10)-89	Escherichia coli	b0200	[pn:hypothetical 21.3 kd protein in abc-rsh intergenic region] [gn:yaed]
CONTIG177	24848941_f2_2	139	5801	525	175	573	3.8(10)-55	Escherichia coli	b2216	[pn:probable sensor protein yojn] [gn:yojn]
CONTIG178	24233325_f1_1	140	5802	429	143	141	6.7(10)-10	Escherichia coli	b3097	[pn:hypothetical 14.5 kd protein in exur-tdcc intergenic region]
CONTIG180	13066567_f1_1	141	5803	654	218	509	6.9(10)-49	Escherichia coli	b4121	[pn:hypothetical 23.4 kd protein in melb-fumb intergenic region] [gn:yjdf]
CONTIG180	30672151_f3_4	142	5804	189	63	270	1.5(10)-23	Escherichia coli	b4121	[pn:hypothetical 23.4 kd protein in melb-fumb intergenic region] [gn:yjdf]
CONTIG182	11814452_f1_1	143	5805	369	123	397	5.0(10)-37	Escherichia coli	b4372	[pn:dna polymerase iii psi subunit] [gn:hold]
CONTIG182	21991462_f2_3	144	5806	492	164	521	3.7(10)-50	Escherichia coli	b4373	[pn:ribosomal-protein-alanine acetyltransferase] [gn:rimi]
CONTIG182	5086063_f2_4	145	5807	351	117	461	8.4(10)-44	Escherichia coli	b4375	[pn:peptide-chain-release factor 3] [gn:prfc]
CONTIG182	34491258_f3_6	146	5808	786	262	999	8.1(10)-101	Escherichia coli	b4374	[pn:hypothetical 22.2 kd protein in rimi-prfc intergenic region] [gn:yjig]

CONTIG184	25969562_c1_6	147	5809	696	232	1038	6.0(10)-105	Escherichia coli	b3346	[pn:hypothetical protein] [gn:yheo]
CONTIG184	4897691_c2_7	148	5810	684	228	837	1.2(10)-83	Escherichia coli	b3347	[pn:fkbp-type peptidyl-prolyl cis-trans isomerase] [gn:fkpa]
CONTIG185	30566284_c1_4	149	5811	1331	443	690	4.5(10)-68	Escherichia coli	b0544	[pn:hypothetical protein] [gn:ybck]
CONTIG186	4416343_f1_3	150	5812	669	223	1018	7.9(10)-103	Escherichia coli	b1180	[pn:hypothetical protein]
CONTIG186	2625761_f2_6	151	5813	402	134	443	6.7(10)-42	Escherichia coli	b1179	[pn:hypothetical protein]
CONTIG186	9805436_c1_12	152	5814	726	242	1038	6.0(10)-105	Escherichia coli	b1175	[pn:cell division inhibitor mind] [gn:mind]
CONTIG186	15681716_c3_17	153	5815	807	269	908	3.6(10)-91	Escherichia coli	b1176	[pn:cell division inhibitor minc] [gn:minc]
CONTIG187	962900_f3_2	154	5816	1848	616	1429	2.2(10)-146	Escherichia coli	b2395	[pn:yfea]
CONTIG188	676041_f2_3	155	5817	1506	502	1903	1.3(10)-196	Escherichia coli	b1603	[pn:pyridine nucleotide transhydrogenase subunit- alpha] [gn:pnta]
CONTIG189	24508336_f3_4	156	5818	579	193	573	1.1(10)-55	Escherichia coli	b1622	[pn:maly protein] [gn:maly]
CONTIG189	34569806_f3_5	157	5819	1011	337	1346	1.3(10)-137	Escherichia coli	b1623	[pn:adenosine deaminase] [gn:add]
CONTIG19	32204791_f1_1	158	5820	399	133	622	7.2(10)-61	Enterobacteriaceae	S07447	[pn:hypothetical protein, 13.1k]
CONTIG19	25522168_f2_3	159	5821	297	99	165	2.0(10)-12	Shigella flexneri	P04337	hypothetical mercuric resistance protein merc.
CONTIG19	12995781_f3_5	160	5822	339	113	114	5.0(10)-7	Shigella flexneri	P04336	mercuric transport protein.
CONTIG19	9901712_c1_6	161	5823	663	221	722	1.8(10)-71	Shigella flexneri	P07044	mercuric resistance operon regulatory protein.
CONTIG190	15651516_f1_1	162	5824	960	320	1514	2.2(10)-155	Escherichia coli	b1415	[pn:lactaldehyde dehydrogenase a] [gn:alda]
CONTIG190	12242930_f1_2	163	5825	444	148	521	3.7(10)-50	Escherichia coli	b1418	[gn:cybb]
CONTIG190	35662765_f2_5	164	5826	204	68	138	1.3(10)-9	Escherichia coli	b1418	[gn:cybb]
CONTIG190	35348277_c1_11	165	5827	345	115	111	1.3(10)-6	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:<1 re:588 di:direct

CONTIG190	16095807_c2_13	166	5828	390	130	135	2.8(10)-9	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:59 re:580 di:direct nt:orf2
CONTIG191	25508265_c2_11	167	5829	735	245	920	1.8(10)-92	Escherichia coli	b1913	[pn:excinuclease abc subunit c] [gn:uvrc]
CONTIG191	6760811_c2_13	168	5830	240	80	316	1.8(10)-28	Escherichia coli	b1912	[pn:phosphotidylglycerophosphate synthetase] [gn:pgsa]
CONTIG191	30128300_c3_15	169	5831	1203	401	1726	7.5(10)-178	Escherichia coli	b1913	[pn:excinuclease abc subunit c] [gn:uvrc]
CONTIG192	29711561_c2_11	170	5832	522	174	322	4.5(10)-29	Escherichia coli	b1043	[pn:hypothetical protein] [gn:csgc]
CONTIG192	35548807_c3_12	171	5833	561	187	589	2.2(10)-57	Escherichia coli	b1041	[pn:nucleation component of curlin monomers] [gn:csgb]
CONTIG192	30331555_c3_13	172	5834	477	159	507	1.1(10)-48	Escherichia coli	b1042	[pn:major curlin subunit precursor] [gn:csga]
CONTIG193	3939087_fl_1	173	5835	891	297	1376	9.1(10)-141	Escherichia coli	b1829	[pn:heat shock protein htpx] [gn:htpx]
CONTIG193	33848575_f2_2	174	5836	1608	536	2343	3.1(10)-243	Escherichia coli	b1830	[pn:tail-specific protease precursor] [gn:prc]
CONTIG194	16019651_fl_2	175	5837	926	309	1434	6.5(10)-147	Escherichia coli	b1370	[pn:insertion element is5 hypothetical protein] [gn:yi52_5]
CONTIG194	30157255_f3_6	176	5838	279	93	94	0.00042	Escherichia coli	U95365	transposase,,is5b,
CONTIG194	21775383_c1_8	177	5839	300	100	447	2.6(10)-42	Escherichia coli	D90774	or:escherichia coli gn:is5 le:13994 re:14362 di:direct sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:orf_id:o263#20; similar to [swissprot accession [pn:hypothetical protein]
CONTIG194	22458580_c1_9	178	5840	348	116	484	3.1(10)-46	Escherichia coli	b1371	
CONTIG195	16500153_c1_7	179	5841	1560	520	1825	2.3(10)-188	Escherichia coli	b2216	[pn:probable sensor protein yojn] [gn:yojn]
CONTIG196	34647280_f2_4	180	5842	414	138	558	4.4(10)-54	Escherichia coli	b4200	[pn:30s ribosomal subunit protein s6] [gn:rpsf]
CONTIG196	14884500_f2_5	181	5843	216	72	345	1.6(10)-31	Escherichia coli	b4201	[pn:primosomal replication protein n] [gn:prib]
CONTIG196	10632182_c1_8	182	5844	399	133	307	1.7(10)-27	Escherichia coli	b4199	[pn:hypothetical 10.1 kd protein in aidb-rpsf intergenic region] [gn:yfy]
CONTIG197	1308262_f2_2	183	5845	564	188	655	2.2(10)-64	Escherichia coli	b3856	[pn:hypothetical protein]

CONTIG198	23687750_c1_6	184	5846	729	243	757	3.6(10)-75	Escherichia coli	b0629	[pn:hypothetical transcriptional regulator in lipa- lipb intergenic region] [gn:ybef]
CONTIG198	3242252_c1_7	185	5847	1011	337	1650	8.5(10)-170	Escherichia coli	b0628	[pn:liipoic acid synthetase] [gn:lipa]
CONTIG199	29723842_c1_10	186	5848	561	187	102	0.0028	Haemophilus influenzae	HI0015	[pn:signal peptidase i] [gn:lepb]
CONTIG199	54140_c2_11	187	5849	813	271	93	0.027	Escherichia coli	AF005044	[PN:TrbC] [GN:trbC] [DE:Escherichia coli plasmid R100-1 TraV (traV), TraR (traR), OrfG1(orfG1), OrfH (orfH), OrfI (orfI), TraC (traC), TrbI (trbI), TraW(traW), TraU (traU), TrbC (trbC), TraN (traN), TrbE (trbE) and TraF(traF) genes, c
CONTIG199	32032827_c3_12	188	5850	741	247	93	0.20999	Vibrio cholerae	Y00557	[PN:protease] [GN:prtV] [DE:Vibrio cholerae DNA for hlyA, hlyB, lipA, lipB and prtV genes.] [LE:7537] [RE:10296] [DI:complement]
CONTIG2	16218928_c2_6	189	5851	417	139	129	1.3(10)-8	Escherichia coli	b1228	[pn:hypothetical protein]
CONTIG2	829150_c3_7	190	5852	336	112	115	3.8(10)-7	Escherichia coli	b1228	[pn:hypothetical protein]
CONTIG20	16620400_c2_3	191	5853	519	173	153	1.2(10)-10	Escherichia coli	b3816	[pn:magnesium and cobalt transport protein cora] [gn:cora]
CONTIG200	31875006_f1_1	192	5854	810	270	966	2.6(10)-97	Escherichia coli	b0761	[pn:molybdenum transport protein mode] [gn:mode]
CONTIG200	10634661_f2_2	193	5855	1164	388	1413	1.1(10)-144	Escherichia coli	b0760	[pn:putative molybdenum transport ap-binding protein modf] [gn:modf]
CONTIG201	5198443_c3_13	194	5856	792	264	490	7.0(10)-47	Bacillus subtilis	ymaE	[pn:hypothetical protein]
CONTIG202	4394533_c3_5	195	5857	831	277	993	3.5(10)-100	Escherichia coli	b0732	[pn:ybgg] [gn:ybgg]
CONTIG203	2375193_c1_4	196	5858	204	68	102	8.9(10)-5	Methanobacterium thermoautotrophicum	MTH104	[pn:multidrug transporter homolog]



CONTIG203	5164657_c1_5	197	5859	741	247	921	1.5(10)-92	Methanobacterium thermoautotrophicum	MTH104	[pn:multidrug transporter homolog]
CONTIG204	1962750_c1_5	198	5860	750	250	962	6.7(10)-97	Escherichia coli	b2055	[pn:hypothetical protein] [gn:wcae]
CONTIG204	4739068_c2_9	199	5861	351	117	558	4.4(10)-54	Escherichia coli	b2054	[pn:hypothetical protein] [gn:wcaf]
CONTIG204	159627_c3_10	200	5862	228	76	355	1.3(10)-32	Escherichia coli	b2056	[pn:hypothetical protein] [gn:wcad]
CONTIG205	26694807_c1_5	201	5863	372	124	594	6.7(10)-58	Yersinia pestis	AF053945	[de:yersinia pestis plasmid ppep1, complete plasmid sequence.]
CONTIG206	21683457_f1_1	202	5864	2127	709	3388	0	Escherichia coli	b3340	[pn:transposase]
CONTIG206	1069655_f1_2	203	5865	378	126	406	5.7(10)-38	Escherichia coli	b3339	[pn:fusa] [gn:fusa]
CONTIG206	510290_c1_4	204	5866	705	235	390	4.0(10)-36	Escherichia coli	P35649	[pn:elongation factor tu] [gn:tufa]
CONTIG206	394215_c1_5	205	5867	1113	371	575	7.0(10)-56	Eikenella corrodens	P35649	hypothetical 66.3 kd protein in hag2 5'region.
CONTIG206	22332280_c2_10	206	5868	189	63	105	5.7(10)-5	Eikenella corrodens	P35649	hypothetical 66.3 kd protein in hag2 5'region.
CONTIG207	2376381_f1_4	207	5869	825	275	274	1.8(10)-33	Escherichia coli	b2106	hypothetical 66.3 kd protein in hag2 5'region.
CONTIG207	21484653_f3_8	208	5870	447	149	168	9.4(10)-13	Escherichia coli	D90774	[pn:hypothetical protein]
CONTIG207	34410768_c2_13	209	5871	987	329	1308	1.5(10)-133	Escherichia coli	b1994	or:escherichia coli gn:is5 le:13994 re:14362 di:direct sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:orf_id:o263#20; similar to [swissprot accession]
CONTIG209	16506441_f1_1	210	5872	1032	344	1346	1.3(10)-137	Escherichia coli	b4175	[pn:insertion element is5 hypothetical 39.3 kd protein]
CONTIG209	25839843_f2_2	211	5873	504	168	450	1.2(10)-42	Escherichia coli	b4174	[pn:hflc protein] [gn:hflc]
CONTIG21	32445803_f2_1	212	5874	909	303	1213	1.7(10)-123	Escherichia coli	P03008	[pn:hflk protein] [gn:hflk]
										transposase for transposon tn3.

CONTIG210	34188287_f2_4	213	5875	2229	743	1836	1.7(10)-189	Pseudomonas aeruginosa	X99514	or:pseudomonas aeruginosa gn:mexf le:1439 re:4627 di:direct
CONTIG210	4867075_f2_5	214	5876	279	93	92	0.00011	Saccharomyces cerevisiae	YNL259C	[pn:antioxidant protein and metal homeostasis factor] [gn:atx1]
CONTIG211	5944581_c2_13	215	5877	1152	384	545	1.1(10)-52	Escherichia coli	AF044506	[de:escherichia coli strain ec50 hop gene, partial cds; rhsg accessory genetic element vrg protein gene, complete cds; and core proteingene, partial cds.] [pn:vrg protein]
CONTIG211	36147260_c2_14	216	5878	1092	364	100	0.12	Mycobacterium tuberculosis	AL021246	[de:mycobacterium tuberculosis sequence v008.] [pn:pgrs-family] [gn:mtv008.46c] [nt:mtv008.46c, member of the m. tuberculosis]
CONTIG212	13880212_f2_2	217	5879	390	130	462	6.5(10)-44	Escherichia coli	b3501	[pn:arsenical resistance operon arsef repressor] [gn:arsr]
CONTIG212	16600136_f2_3	218	5880	393	131	463	5.2(10)-44	Escherichia coli	A25937	arsenical pump-driving atpase (ec 3.6.1.-) - escherichia coli plasmid r773 this anion-transporting atpase catalyzes the extrusion of the oxyanions arsenite, antimonite, and arsenate, thus lowering the intracellular concentration of these toxic oxyanions.
CONTIG212	31284383_c2_8	219	5881	333	111	494	2.7(10)-47	Enterobacter aerogenes	U67194	or:enterobacter aerogenes pn:tnpa gn:tnpa le:10303 re:11307 di:complement
CONTIG213	4494000_f1_1	220	5882	954	318	90	0.23	Drosophila melanogaster	M19537	or:drosophila melanogaster gn:rp115 le:join(<1 re:133,379 di:direct sr:d.melanogaster dna, clone p4.1 nt:description: rna polymerase ii 215kd subunit; rna

CONTIG213	24478907_f1_2	221	5883	1770	590	150	2.5(10)-15	Escherichia coli	AF005044	[PN:TraC] [GN:traC] [DE:Escherichia coli plasmid R100-1 TraV (traV), TraR (traR), OrfG1(orfG1), OrfH (orfH), OrfI (orfI), TraC (traC), TrbI (trbI), TraW(traw), TraU (traU), TrbC (trbC), TraN (traN), TrbE (trbE) and TraF(traf) genes, c
CONTIG214	22770005_f3_5	222	5884	1548	516	2262	1.2(10)-234	Escherichia coli	b2149	[pn:galactoside transport atp-binding protein mglA] [gn:mglA]
CONTIG214	3417013_f3_6	223	5885	1026	342	1311	7.0(10)-134	Escherichia coli	b2148	[pn:galactoside transport system permease protein mglC] [gn:mglC]
CONTIG215	5078300_f3_6	224	5886	876	292	1456	3.1(10)-149	Escherichia coli	b4192	[pn:hypothetical 40.3 kd protein in aidb-rpsf intergenic region] [gn:yjfr]
CONTIG215	22367036_c1_8	225	5887	804	268	1175	1.8(10)-119	Escherichia coli	b4193	[pn:hypothetical 52.9 kd protein in aidb-rpsf intergenic region] [gn:yjfs]
CONTIG216	10411407_f1_3	226	5888	388	130	195	1.3(10)-15	Rhizobium sp.	S34667	hypothetical protein 140 - rhizobium sp.
CONTIG216	10563465_f2_6	227	5889	420	140	102	9.3(10)-6	Escherichia coli	b2861	[pn:insertion element is2 hypothetical 13.4 kd protein]
CONTIG216	6383505_c3_17	228	5890	633	211	232	1.6(10)-19	Bacillus subtilis	soj	[pn:hypothetical protein]
CONTIG217	4301907_f1_1	229	5891	1212	404	1634	4.2(10)-168	Escherichia coli	b2393	[pn:nucleoside permease nupC] [gn:nupC]
CONTIG217	4900012_c1_11	230	5892	651	217	505	4.5(10)-48	Escherichia coli	b2395	[pn:yfea]
CONTIG218	21697135_f2_2	231	5893	873	291	1293	5.7(10)-132	Escherichia coli	b4396	[pn:right origin-binding protein] [gn:rob]
CONTIG22	34587776_c1_4	232	5894	201	67	315	2.5(10)-28	Escherichia coli	b3232	[pn:hypothetical 43.1 kd protein in rplm-hhoA intergenic region] [gn:yhcm]
CONTIG220	13775768_f2_1	233	5895	432	144	675	1.8(10)-66	Enterobacter aerogenes	U67194	or:enterobacter aerogenes pn:tnpa gn:tnpa le:10303 re:11307 di:complement
CONTIG220	1977312_c1_7	234	5896	882	294	91	0.031	Bacillus subtilis	ydcM	[pn:hypothetical protein]
CONTIG221	23609515_f1_1	235	5897	294	98	419	2.3(10)-39	Enterobacter agglomerans	A38965	hypothetical protein a (insertion sequence is1222) -enterobacter agglomerans

CONTIG221	9896067_f3_4	236	5898	185	62	185	1.5(10)-14	Enterobacter agglomerans	B38965	hypothetical protein b (insertion sequence is1222) -enterobacter agglomerans
CONTIG221	190787_c1_6	237	5899	513	171	143	4.2(10)-10	Rhizobium sp.	P55623	hypothetical 18.8 kd protein y4qb,
CONTIG222	29744052_f3_7	238	5900	1122	374	165	6.0(10)-10	Methanococcus jannaschii	MJ1187	[pn:dinitrogenase reductase activating glycohydrolase] [gn:drag]
CONTIG223	34486000_f1_1	239	5901	249	83	341	4.4(10)-31	Escherichia coli	b2833	[pn:hypothetical protein]
CONTIG223	12699062_f1_2	240	5902	459	153	704	1.5(10)-69	Escherichia coli	b2834	[pn:hypothetical protein]
CONTIG223	15898537_f1_3	241	5903	324	108	320	7.2(10)-29	Escherichia coli	b2834	[pn:hypothetical protein]
CONTIG223	10425262_f2_5	242	5904	729	243	954	4.7(10)-96	Escherichia coli	b2831	[pn:muth] [gn:muth]
CONTIG223	24720916_f2_8	243	5905	375	125	504	2.2(10)-48	Escherichia coli	b2834	[pn:hypothetical protein]
CONTIG223	32550407_c3_17	244	5906	354	118	393	1.3(10)-36	Escherichia coli	b2835	[pn:hypothetical protein in muth-aas intergenic region] [gn:yged]
CONTIG224	35447642_c1_15	245	5907	915	305	1424	7.5(10)-146	Escherichia coli	S70160	,
CONTIG224	4110291_c1_16	246	5908	420	140	639	1.2(10)-62	Escherichia coli	S70161	,
CONTIG224	3907697_c2_17	247	5909	1473	491	2437	3.3(10)-253	Escherichia coli	S70159	,
CONTIG224	3939017_c2_20	248	5910	273	91	434	6.0(10)-41	Escherichia coli	S70163	,
CONTIG225	24301557_f1_2	249	5911	1362	454	1668	1.0(10)-171	Escherichia coli	b3502	[pn:arsenical pump membrane protein] [gn:arsb]
CONTIG225	4110263_f1_3	250	5912	279	93	416	4.9(10)-39	Escherichia coli	b3503	[pn:arsenate reductase] [gn:arsc]
CONTIG225	36135812_f3_6	251	5913	336	112	431	1.3(10)-40	Escherichia coli	A25937	arsenical pump-driving atpase (ec 3.6.1.-) - escherichia coli plasmid r773 this anion-transporting atpase catalyzes the extrusion of the oxyanions arsenite, antimonic, and arsenate, thus lowering the intracellular concentration of these toxic oxyanions.

CONTIG225	15661513_c1_10	252	5914	639	213	359	1.1(10)-32	Rhizobium sp.	P50360	hypothetical 29.3 kD protein in region 2 of sym plasmid (no1265).
CONTIG226	22760817_f1_1	253	5915	426	142	472	5.7(10)-45	Escherichia coli	b4206	[pn:hypothetical protein] [gn:ytfb]
CONTIG226	7220463_c1_13	254	5916	462	154	536	9.5(10)-52	Escherichia coli	b4203	[pn:50s ribosomal subunit protein 19] [gn:rplj]
CONTIG226	15056377_c2_16	255	5917	246	82	385	9.5(10)-36	Escherichia coli	b4202	[pn:30s ribosomal subunit protein s18] [gn:rpsf]
CONTIG226	38537_c3_22	256	5918	957	319	151	9.0(10)-9	Escherichia coli	b1533	[pn:hypothetical 28.7 kD protein in marb-dcp intergenic region] [gn:yded]
CONTIG227	3382959_c2_10	257	5919	287	95	231	2.0(10)-19	Escherichia coli	b1099	[pn:dna polymerase iii, delta"" subunit] [gn:holb]
CONTIG227	19800956_c3_11	258	5920	873	291	1259	2.2(10)-128	Escherichia coli	b1100	[pn:hypothetical protein in holb-ptsg intergenic region] [gn:yctf]
CONTIG227	24010937_c3_12	259	5921	1467	489	2174	2.5(10)-225	Escherichia coli	b1101	[pn:pts system, glucose-specific iibc component] [gn:ptsg]
CONTIG228	15814663_c2_11	260	5922	531	177	144	9.0(10)-10	Methanobacterium thermoautotrophicum	MTH806	[pn:protease iv]
CONTIG228	26598393_c3_13	261	5923	738	246	95	0.024	Methanobacterium thermoautotrophicum	MTH466	[pn:unknown]
CONTIG229	12927341_f2_3	262	5924	825	275	1127	2.2(10)-114	Escherichia coli	b2741	[pn:rna polymerase sigma subunit rpos] [gn:rpos]
CONTIG229	15673942_f3_4	263	5925	1095	365	1130	1.1(10)-114	Escherichia coli	b2742	[pn:lipoprotein nlpd precursor] [gn:nlpd]
CONTIG23	24634555_c1_3	264	5926	390	130	100	1.5(10)-5	Escherichia coli	b4283	[pn:insertion element is911 hypothetical 11.6 kD protein] [gn:y91]
CONTIG230	9860003_f2_4	265	5927	1815	605	277	3.3(10)-21	Bacteriophage T7	P03692	dna primase, chains a and b (ec 2.7.7.-)
CONTIG230	32505283_f3_6	266	5928	1029	343	272	2.2(10)-23	Escherichia coli	b1149	[pn:hypothetical protein]
CONTIG231	11963255_f2_3	267	5929	546	182	382	2.0(10)-35	Escherichia coli	b0633	[pn:rare lipoprotein a precursor] [gn:ripa]

CONTIG231	12363162_f2_5	268	5930	318	106	371	2.8(10)-34	Escherichia coli	b0631	[pn:hypothetical 9.8 kd protein in lipb-daca intergenic region]
CONTIG231	15723956_f3_6	269	5931	270	90	204	3.7(10)-16	Escherichia coli	b0633	[pn:rare lipoprotein a precursor] [gn:r pa]
CONTIG231	25563161_f3_7	270	5932	1305	435	1920	2.1(10)-198	Escherichia coli	b0632	[pn:d-alanine carboxypeptidase] [gn:daca]
CONTIG231	32516510_f3_9	271	5933	687	229	858	7.2(10)-86	Escherichia coli	b0630	[pn:lipoate-protein ligase b] [gn:lipb]
CONTIG232	25839831_fl_1	272	5934	1161	387	1782	8.6(10)-184	Escherichia coli	b0677	[pn:n-acetylglucosamine-6-phosphate deacetylase] [gn:naga]
CONTIG232	3009677_f3_5	273	5935	804	268	1338	9.8(10)-137	Escherichia coli	b0678	[pn:glucosamine-6-phosphate isomerase] [gn:nagb]
CONTIG232	34174053_f3_7	274	5936	1197	399	1782	8.6(10)-184	Escherichia coli	b0676	[pn:n-acetylglucosamine repressor] [gn:nagc]
CONTIG233	26445952_c1_13	275	5937	345	115	285	3.7(10)-25	Escherichia coli	b0551	[pn:hypothetical protein] [gn:ybcq]
CONTIG233	36366038_c3_18	276	5938	636	212	159	8.4(10)-12	Escherichia coli	b0798	[pn:hypothetical 18.7 kd protein in rhle-ding/rarb intergenic region] [gn:ybia]
CONTIG236	4166305_c1_17	277	5939	651	217	174	2.2(10)-13	Escherichia coli	b2846	[pn:hypothetical protein]
CONTIG236	5353400_c3_22	278	5940	2436	812	3750	0	Escherichia coli	b0221	[pn:hypothetical protein]
CONTIG236	4191068_c3_23	279	5941	789	263	1171	4.9(10)-119	Escherichia coli	b0219	[pn:hypothetical protein] [gn:yafv]
CONTIG237	21517313_fl_1	280	5942	498	166	483	3.8(10)-46	Escherichia coli	b4365	[pn:hypothetical 27.0 kd protein in dnat-hold intergenic region]
CONTIG237	24705188_f2_10	281	5943	1149	383	478	1.3(10)-45	Bacillus subtilis	litC	[pn:phosphotransferase system] [gn:celb]
CONTIG237	30557000_f2_11	282	5944	213	71	93	0.00079	Borrelia burgdorferi	AE000792	[de:borrelia burgdorferi plasmid cp26, complete plasmid sequence.] [pn:pts system, cellobiose-specific iic component] [gn:bbb04] [nt:similar to gb]
CONTIG237	24431568_f3_14	283	5945	690	230	494	2.7(10)-47	Escherichia coli	b4366	[pn:hypothetical 25.6 kd protein in dnat-hold intergenic region] [gn:yjir]
CONTIG237	14237692_c1_17	284	5946	801	267	964	4.2(10)-97	Escherichia coli	b4367	[pn:hypothetical 30.1 kd protein in dnat-hold intergenic region] [gn:yjis]

CONTIG237	22765961_c1_20	285	5947	492	164	335	1.8(10)-30	Bacillus subtilis	ywhH	[pn:hypothetical protein]
CONTIG238	2382277_f3_9	286	5948	1005	335	1289	1.5(10)-131	Escherichia coli	b3862	[pn:hypothetical 36.3 kd protein in pola 5'''region] [gn:yihg]
CONTIG238	21488165_f3_12	287	5949	555	185	552	1.8(10)-53	Escherichia coli	b3857	[pn:molybdopterin-guanine dinucleotide biosynthesis protein a] [gn:moba]
CONTIG238	2034766_c1_13	288	5950	300	100	442	8.6(10)-42	Escherichia coli	b3858	[pn:hypothetical 10.3 kd protein in moba 3'''region] [gn:yihd]
CONTIG238	25603875_c1_14	289	5951	1002	334	1428	2.7(10)-146	Escherichia coli	b3859	[pn:hypothetical 38.1 kd protein in dsba 5'''region] [gn:yihe]
CONTIG238	12614200_c1_15	290	5952	651	217	867	8.0(10)-87	Escherichia coli	b3860	[pn:dsba] [gn:dsba]
CONTIG239	13063513_f1_3	291	5953	264	88	339	7.0(10)-31	Escherichia coli	b1203	[pn:hypothetical gtp-binding protein in pth 3''' region] [gn:yichf]
CONTIG239	4948290_f1_6	292	5954	249	83	154	1.3(10)-10	Escherichia coli	b1203	[pn:hypothetical gtp-binding protein in pth 3''' region] [gn:yichf]
CONTIG239	4948291_f2_7	293	5955	306	102	186	3.7(10)-14	Escherichia coli	b1203	[pn:hypothetical gtp-binding protein in pth 3''' region] [gn:yichf]
CONTIG239	3395628_c1_19	294	5956	1332	444	253	2.2(10)-28	Escherichia coli	b0703	[pn:hypothetical protein]
CONTIG239	5273307_c2_22	295	5957	804	268	126	6.4(10)-8	coliphage T4	P13309	hypothetical 17.5 kd protein in tk-vs intergenic region.
CONTIG240	26753130_c1_15	296	5958	204	68	298	1.6(10)-26	Escherichia coli	b2696	[pn:csra] [gn:csra]
CONTIG240	35554527_c2_16	297	5959	267	89	218	8.1(10)-17	Escherichia coli	b2697	[pn:alanyl-trna synthetase] [gn:alas]
CONTIG240	36611683_c3_17	298	5960	2121	707	2566	7.2(10)-267	Escherichia coli	b2697	[pn:alanyl-trna synthetase] [gn:alas]
CONTIG241	3224078_f1_2	299	5961	897	299	135	8.0(10)-9	Escherichia coli	b2358	[pn:hypothetical protein] [gn:yfdo]
CONTIG241	6367327_f1_3	300	5962	657	219	157	1.3(10)-11	Escherichia coli	b1361	[pn:hypothetical protein]
CONTIG241	4900300_f2_4	301	5963	417	139	350	4.9(10)-32	Escherichia coli	b1358	[pn:hypothetical protein]
CONTIG241	36207628_f2_6	302	5964	378	126	127	2.1(10)-8	Synechocystis sp.	S74697	[PN:hypothetical protein sll1193] [OR:Synechocystis sp.] [SR:PCC 6803,, PCC 6803] [SR:PCC 6803,,]

CONTIG241	35745317_c1_14	303	5965	615	205	1117	0.00012	Helicobacter pylori	HP0966	[pn:conserved hypothetical protein]
CONTIG242	4491678_f2_5	304	5966	726	242	1200	4.0(10)-122	Escherichia coli	b4401	[pn:aerobic respiration control protein arca] [gn:arca]
CONTIG242	10954577_c2_15	305	5967	687	229	834	2.5(10)-83	Escherichia coli	b4403	[pn:hypothetical 25.3 kd protein in arca-thrl intergenic region] [gn:last]
CONTIG242	4098393_c3_16	306	5968	480	160	698	6.4(10)-69	Escherichia coli	b4397	[pn:crea protein] [gn:crea]
CONTIG243	245337_c1_13	307	5969	864	288	249	2.3(10)-21	Bacteriophage 186	U32222	or:bacteriophage 186 le:181 re:711 di:direct nt:orf38; similar to bacteriophage p2 i protein, probable tail fibre protein (gph).
CONTIG243	23714843_c3_20	308	5970	840	280	163	1.7(10)-9	Bacteriophage p2	P26700	
CONTIG244	16658141_f1_2	309	5971	357	119	502	3.7(10)-48	Escherichia coli	b4195	[pn:hypothetical phosphotransferase enzyme ii] [gn:ptxa]
CONTIG244	24617338_f1_3	310	5972	663	221	1063	1.3(10)-107	Escherichia coli	b4196	[pn:hypothetical 23.6 kd protein in aidb-rpsf intergenic region] [gn:yjfv]
CONTIG244	4197943_f1_4	311	5973	870	290	1382	2.1(10)-141	Escherichia coli	b4197	[pn:hypothetical 32.0 kd protein in aidb-rpsf intergenic region]
CONTIG244	22120707_f2_5	312	5974	654	218	897	5.2(10)-90	Escherichia coli	b4193	[pn:hypothetical 52.9 kd protein in aidb-rpsf intergenic region] [gn:yjfs]
CONTIG244	5156693_f2_6	313	5975	318	106	421	1.5(10)-39	Escherichia coli	b4194	[pn:hypothetical 10.9 kd protein in aidb-rpsf intergenic region] [gn:yjft]
CONTIG244	4473818_f2_7	314	5976	186	62	188	7.0(10)-15	Escherichia coli	b4195	[pn:hypothetical phosphotransferase enzyme ii] [gn:ptxa]
CONTIG244	24870965_f2_11	315	5977	570	190	706	9.1(10)-70	Escherichia coli	b4198	[pn:hypothetical protein] [gn:sgae]
CONTIG245	31377187_f2_6	316	5978	708	236	107	0.00058	Escherichia coli	b4366	[pn:hypothetical 25.6 kd protein in dnaat-hold intergenic region] [gn:yjir]
CONTIG245	198377_f3_9	317	5979	762	254	107	0.0023	Escherichia coli	b0457	[pn:hypothetical protein] [gn:ylab]
CONTIG245	25583543_c2_16	318	5980	186	62	220	2.8(10)-18	Escherichia coli	b3661	[pn:lipoprotein-28 precursor] [gn:nlpa]
CONTIG245	1270761_c2_17	319	5981	555	185	750	2.0(10)-74	Escherichia coli	b3661	[pn:lipoprotein-28 precursor] [gn:nlpa]
CONTIG246	35722707_f1_4	320	5982	240	80	168	9.4(10)-13	Zea mays	S58640	hypothetical protein 137 - maize chloroplast
CONTIG246	31910000_c1_15	321	5983	237	79	351	3.7(10)-32	Escherichia coli	b3319	[pn:50s ribosomal subunit protein 14] [gn:rpId]



CONTIG246	22847717_c1_17	322	5984	819	273	1094	7.0(10)-111	Escherichia coli	b3314	[pn:30s ribosomal subunit protein s3] [gn:rpsc]
CONTIG246	3213568_c1_18	323	5985	423	141	675	1.8(10)-66	Escherichia coli	b3313	[pn:50s ribosomal subunit protein 116] [gn:rplp]
CONTIG246	34627178_c2_19	324	5986	843	281	1349	6.7(10)-138	Escherichia coli	b3317	[pn:50s ribosomal subunit protein 12]
CONTIG246	5289213_c2_20	325	5987	342	114	469	1.2(10)-44	Escherichia coli	b3315	[gn:rplb]
CONTIG246	25500038_c2_22	326	5988	270	90	325	2.2(10)-29	Escherichia coli	b3311	[pn:50s ribosomal subunit protein 122] [gn:rplv]
CONTIG246	35282943_c3_23	327	5989	315	105	209	4.2(10)-17	Escherichia coli	b3318	[pn:30s ribosomal subunit protein s17] [gn:rpsq]
CONTIG246	32134658_c3_26	328	5990	282	94	441	1.1(10)-41	Escherichia coli	b3316	[pn:50s ribosomal subunit protein 123] [gn:rplw]
CONTIG246	6064788_c3_27	329	5991	204	68	210	3.2(10)-17	Escherichia coli	b3312	[pn:30s ribosomal subunit protein s19] [gn:rpss]
CONTIG247	3125207_f1_1	330	5992	1308	436	1735	8.3(10)-179	Escherichia coli	b1309	[pn:50s ribosomal subunit protein 129] [gn:rpnc]
CONTIG247	20593891_f1_2	331	5993	375	125	292	3.5(10)-25	Escherichia coli	b1309	[pn:hypothetical protein]
CONTIG247	24227265_f2_9	332	5994	1326	442	1595	5.7(10)-164	Escherichia coli	b1310	[pn:hypothetical protein]
CONTIG247	15630213_f2_10	333	5995	903	301	1055	9.5(10)-107	Escherichia coli	b1311	[pn:hypothetical protein]
CONTIG247	16835032_f3_15	334	5996	645	215	917	4.0(10)-92	Escherichia coli	b1312	[pn:hypothetical protein]
CONTIG248	17003167_f1_1	335	5997	459	153	607	2.7(10)-59	Escherichia coli	b2210	[pn:hypothetical 60.2 kd protein in eco-alkB intergenic region] [gn:yojh]
CONTIG248	125915_f1_2	336	5998	372	124	221	2.2(10)-18	Escherichia coli	b1332	[pn:hypothetical protein]
CONTIG248	35650802_f1_7	337	5999	410	137	516	1.2(10)-49	Escherichia coli	b2186	[pn:hypothetical 37.8 kd protein in rply-prol intergenic region] [gn:yejk]
CONTIG248	16506555_c1_17	338	6000	1875	625	2506	1.7(10)-260	Escherichia coli	b2188	[pn:hypothetical 67.3 kd protein in rply-prol intergenic region]
CONTIG248	4976561_c2_22	339	6001	519	173	632	6.4(10)-62	Escherichia coli	b2209	[pn:tin precursor] [gn:eco]
CONTIG248	22369000_c3_24	340	6002	231	77	319	9.4(10)-29	Escherichia coli	b2187	[pn:hypothetical 8.3 kd protein in rply-prol intergenic region] [gn:yejl]

CONTIG249	25507783_fl_2	341	6003	1458	486	2340	6.5(10)-243	Escherichia coli	S70165	,
CONTIG249	35807766_fl_4	342	6004	732	244	963	5.2(10)-97	Escherichia coli	S70163	,
CONTIG249	546888_fl_5	343	6005	711	237	620	1.2(10)-60	Escherichia coli	b0571	[pn:hypothetical protein] [gn:ylca]
CONTIG249	10563465_fl_9	344	6006	413	138	102	9.3(10)-6	Escherichia coli	b2861	[pn:insertion element is2 hypothetical 13.4 kd protein]
CONTIG250	12277217_cl_14	345	6007	1230	410	937	3.0(10)-94	Escherichia coli	b2349	[pn:putative prophage sf6-like integrase in argw 3'''region] [gn:yfdb]
CONTIG250	35552032_c2_15	346	6008	1227	409	1710	3.7(10)-176	Escherichia coli	b0243	[pn:gamma-glutamyl phosphate reductase] [gn:proa]
CONTIG251	34417067_cl_12	347	6009	399	133	631	8.0(10)-62	Escherichia coli	b3306	[pn:30s ribosomal subunit protein s8] [gn:rpsh]
CONTIG251	34198580_cl_13	348	6010	546	182	850	5.0(10)-85	Escherichia coli	b3305	[pn:50s ribosomal subunit protein 16] [gn:rp1f]
CONTIG251	4020268_cl_14	349	6011	363	121	434	6.0(10)-41	Escherichia coli	b3304	[pn:50s ribosomal subunit protein 118] [gn:rp1r]
CONTIG251	23572211_cl_18	350	6012	927	309	1047	6.7(10)-106	Escherichia coli	b3300	[pn:preprotein translocase secy subunit] [gn:secy]
CONTIG251	5267592_c3_22	351	6013	510	170	829	8.5(10)-83	Escherichia coli	b3303	[pn:30s ribosomal subunit protein s5] [gn:rpse]
CONTIG251	16832200_c3_23	352	6014	186	62	283	6.0(10)-25	Escherichia coli	b3302	[pn:50s ribosomal subunit protein 130] [gn:rpmd]
CONTIG251	259713_c3_24	353	6015	438	146	398	4.0(10)-37	Escherichia coli	b3301	[pn:50s ribosomal subunit protein 115] [gn:rplo]
CONTIG252	4698250_cl_32	354	6016	273	91	215	9.8(10)-18	Escherichia coli	b1346	[pn:hypothetical protein]
CONTIG252	11744091_c3_43	355	6017	1389	463	306	1.1(10)-24	Escherichia coli	b1350	[pn:exodeoxyribonuclease viii] [gn:rece]
CONTIG252	2845092_c3_45	356	6018	1392	464	1150	8.1(10)-117	Escherichia coli	b1345	[pn:hypothetical protein]
CONTIG253	29503136_fl_1	357	6019	522	174	338	9.0(10)-31	Escherichia coli	b3981	[pn:preprotein translocase secy subunit] [gn:sece]
CONTIG253	1035760_fl_3	358	6020	1221	407	1578	3.6(10)-162	Escherichia coli	b3339	[pn:elongation factor tu] [gn:tufa]
CONTIG253	5281893_fl_7	359	6021	555	185	927	3.5(10)-93	Escherichia coli	b3982	[pn:transcription antitermination protein nusg] [gn:nusg]

CONTIG254	3416407_f1_8	360	6022	309	103	365	1.2(10)-33	Escherichia coli	b2609	[pn:30s ribosomal subunit protein s16] [gn:rpsp]
CONTIG254	7144130_f1_9	361	6023	555	185	812	5.4(10)-81	Escherichia coli	b2608	[pn:hypothetical 21.0 kd protein in trmd-rsp intergenic region]
CONTIG254	22789178_f2_10	362	6024	441	147	622	7.2(10)-61	Escherichia coli	b2614	[pn:heat shock protein] [gn:grpe]
CONTIG254	12268766_f2_12	363	6025	1389	463	1971	8.1(10)-204	Escherichia coli	b2610	[pn:signal recognition particle protein] [gn:ffh]
CONTIG254	26366082_f2_14	364	6026	786	262	1233	1.3(10)-125	Escherichia coli	b2607	[pn:guanine-7methyltransferase] [gn:trmd]
CONTIG254	16304761_c1_27	365	6027	888	296	1070	2.5(10)-108	Escherichia coli	b2611	[pn:hypothetical protein]
CONTIG254	1385936_c2_37	366	6028	1302	434	934	6.2(10)-94	Escherichia coli	b2613	[pn:hypothetical protein in grpe 3''' region] [gn:yfd]
CONTIG255	30328941_f2_6	367	6029	1095	365	1120	1.2(10)-113	Escherichia coli	b1620	[pn:repressor protein] [gn:mali]
CONTIG255	19922527_c1_11	368	6030	693	231	794	4.2(10)-79	Escherichia coli	b1621	[pn:pts system, maltose and glucose-specific ii abc component] [gn:malx]
CONTIG255	13674218_c1_12	369	6031	603	201	785	3.8(10)-78	Escherichia coli	b1622	[pn:maly protein] [gn:maly]
CONTIG255	1036625_c2_15	370	6032	993	331	1307	1.8(10)-133	Escherichia coli	b1621	[pn:pts system, maltose and glucose-specific ii abc component] [gn:malx]
CONTIG256	33725701_f1_2	371	6033	1035	345	1335	2.0(10)-136	Escherichia coli	b1235	[pn:hnr protein] [gn:hnr]
CONTIG256	1299183_f2_4	372	6034	954	318	1218	5.0(10)-124	Escherichia coli	b1234	[pn:hypothetical 34.4 kd protein in hnr-puru intergenic region]
CONTIG256	32126890_f2_6	373	6035	840	280	950	1.3(10)-95	Escherichia coli	b1236	[pn:glucose-1-phosphate uridylyltransferase] [gn:ga]
CONTIG256	14572040_c1_15	374	6036	891	297	1318	1.3(10)-134	Escherichia coli	b1232	[pn:formyltetrahydrofolate deformylase] [gn:puru]
CONTIG256	31416_c2_18	375	6037	495	165	542	2.2(10)-52	Escherichia coli	b1233	[pn:hypothetical 17.0 kd protein in hnr-puru intergenic region] [gn:ychi]
CONTIG257	32145043_f2_3	376	6038	222	74	235	7.5(10)-20	synthetic construct	M15619	or:artificial sequence le:29 re:>232 di:direct sr:c.coli (strain sc5000) synthetic dna, clone pkb1 nt:orf16-lacZ fusion protein
CONTIG257	12781963_c3_15	377	6039	189	63	101	1.2(10)-5	Homo sapiens	A44803	pgl protein - human (fragment)
CONTIG258	22438128_f2_4	378	6040	645	215	846	1.3(10)-84	Escherichia coli	b3740	[pn:glucose inhibited division protein] [gn:gldb]

CONTIG258	24277186_f2_5	379	6041	411	137	600	1.6(10)-58	Escherichia coli	b3738	[pn:atp synthase f0 subunit a] [gn:atpb]
CONTIG258	13147637_f3_6	380	6042	1947	649	2757	4.2(10)-287	Escherichia coli	b3741	[pn:glucose inhibited division protein] [gn:gida]
CONTIG258	2071915_f3_7	381	6043	456	152	436	3.7(10)-41	Escherichia coli	b3739	[pn:atp synthase subunit] [gn:api]
CONTIG259	881516_f2_5	382	6044	768	256	441	1.1(10)-41	Escherichia coli	b2110	[pn:hypothetical 26.6 kd fimbrial chaperone in mrp 5'''region] [gn:yehe]
CONTIG259	3007955_f2_6	383	6045	2508	836	1320	7.9(10)-135	Escherichia coli	b2109	[pn:hypothetical outer membrane usher protein in mrp 5'''region] [gn:yehe]
CONTIG259	3948925_f2_7	384	6046	1080	360	147	9.5(10)-8	Escherichia coli	b2108	[pn:hypothetical 36.9 kd protein in mrp 5'''region precursor] [gn:yehe]
CONTIG259	87513_c3_25	385	6047	498	166	176	1.3(10)-13	Escherichia coli	b2217	[pn:regulator of capsule synthesis b component] [gn:resb]
CONTIG260	16848787_f1_1	386	6048	1545	515	2236	6.7(10)-232	Escherichia coli	b2411	[pn:dna ase] [gn:lig]
CONTIG260	4114465_f3_11	387	6049	603	201	865	1.3(10)-86	Escherichia coli	b2411	[pn:dna ase] [gn:lig]
CONTIG260	5257752_f3_14	388	6050	951	317	1146	2.2(10)-116	Escherichia coli	b2409	[pn:hypothetical protein]
CONTIG260	1188966_c1_19	389	6051	1020	340	1320	7.9(10)-135	Escherichia coli	b2410	[pn:hypothetical 36.4 kd protein in xapb-lig intergenic region] [gn:yfeh]
CONTIG261	35816301_f1_1	390	6052	1587	529	1699	5.4(10)-175	Escherichia coli	b2176	[pn:hypothetical protein] [gn:rtin]
CONTIG261	21676375_f1_2	391	6053	1881	627	2470	1.1(10)-256	Escherichia coli	b2177	[pn:hypothetical protein in bcr 5'''region] [gn:yeja]
CONTIG261	5197318_f1_3	392	6054	834	278	1164	2.7(10)-118	Escherichia coli	b2178	[pn:hypothetical 40.4 kd protein in bcr 5''' region] [gn:yejb]
CONTIG261	35781511_f3_6	393	6055	294	98	394	1.1(10)-36	Escherichia coli	b2175	[pn:hypothetical protein] [gn:spr]
CONTIG262	13930381_c2_20	394	6056	1512	504	735	7.7(10)-73	Escherichia coli	b0139	[pn:outer membrane usher protein htrc precursor] [gn:htrc]
CONTIG262	32287800_c2_22	395	6057	600	200	97	0.00459	Escherichia coli	b2369	[pn:putative positive transcription regulator evga] [gn:evga]
CONTIG262	22460963_c3_24	396	6058	1209	403	454	3.1(10)-42	Escherichia coli	b3144	[pn:hypothetical outer membrane usher protein in agal-mtr intergenic region] [gn:yraj]

CONTIG262	5286713_c3_25	397	6059	1161	387	107	0.016	Saccharomyces cerevisiae	YOL155C	[pn:similarity to glucan 1,4-alpha-glucosidase mal5p]
CONTIG263	5119186_fl_3	398	6060	1214	405	98	0.23	Saccharomyces cerevisiae	YCR089W	[pn:required for efficient mating] [gn:fig2]
CONTIG263	21641312_f2_4	399	6061	903	301	1471	7.9(10)-151	Serratia marcescens	U62007	rep protein,, rep hi2b, putative rep protein; similar to inch1 plasmid r27 trae protein.
CONTIG263	3164077_f2_5	400	6062	789	263	121	5.7(10)-6	Salmonella typhimurium	P12057	
CONTIG264	34386086_f1_2	401	6063	642	214	873	1.8(10)-87	Escherichia coli	b2302	[pn:hypothetical protein] [gn:yfcg]
CONTIG264	12239405_f1_3	402	6064	444	148	560	2.7(10)-54	Escherichia coli	b2303	[pn:hypothetical protein] [gn:folx]
CONTIG264	14505385_f1_4	403	6065	513	171	178	8.1(10)-14	Bacillus subtilis	yjcK	[pn:hypothetical protein]
CONTIG264	16286633_f2_10	404	6066	927	309	1223	1.5(10)-124	Escherichia coli	b2304	[pn:hypothetical protein]
CONTIG264	12001891_e1_24	405	6067	504	168	548	5.0(10)-53	Escherichia coli	b2301	[pn:hypothetical protein] [gn:yfcf]
CONTIG264	24634813_c2_25	406	6068	720	240	1065	8.3(10)-108	Escherichia coli	b2306	[pn:histidine transport atp-binding protein hisp] [gn:hisp]
CONTIG265	3145261_f1_1	407	6069	1050	350	1257	3.7(10)-128	Escherichia coli	b2810	[pn:hypothetical protein]
CONTIG265	16258266_f1_2	408	6070	501	167	574	8.9(10)-56	Escherichia coli	b2811	[pn:hypothetical protein] [gn:ygdK]
CONTIG265	32708250_f2_6	409	6071	210	70	301	7.5(10)-27	Escherichia coli	b2810	[pn:hypothetical protein]
CONTIG265	15136718_c2_20	410	6072	1134	378	1789	1.6(10)-184	Escherichia coli	b2813	[pn:membrane-bound lytic murein transglycosylase a precursor] [gn:mlta]
CONTIG265	10968768_c2_21	411	6073	813	271	1292	7.2(10)-132	Escherichia coli	b2812	[pn:hypothetical protein] [gn:ygdI]
CONTIG265	26756450_c3_29	412	6074	207	69	237	4.5(10)-20	Escherichia coli	b2809	[pn:hypothetical protein]
CONTIG266	30078286_f1_6	413	6075	462	154	452	7.5(10)-43	Escherichia coli	b2171	[pn:hypothetical 30.9 kd protein in frub 5'''region] [gn:yeip]
CONTIG266	16453125_f3_24	414	6076	1272	424	1593	9.3(10)-164	Escherichia coli	b2170	[pn:hypothetical 42.7 kd protein in frub 5'''region] [gn:yeio]
CONTIG266	25521057_c2_35	415	6077	996	332	1477	1.8(10)-151	Escherichia coli	b2168	[pn:1-phosphofructokinase] [gn:fruk]

CONTIG266	5195383_c3_42	416	6078	1167	389	1476	2.2(10)-151	Escherichia coli	b2169	[pn:pts system, fructose-specific iia/fpr component]
CONTIG266	14453287_c3_43	417	6079	1755	585	2135	3.3(10)-221	Escherichia coli	b2167	[pn:pts system, fructose-specific iibc component] [gn:frua]
CONTIG267	36570837_f1_2	418	6080	2496	832	3364	0	Escherichia coli	b3863	[pn:dna polymerase i] [gn:polA]
CONTIG267	24644037_f1_4	419	6081	258	86	116	3.0(10)-7	Escherichia coli	b3866	[pn:hypothetical 19.1 kd protein in pola-hemm intergenic region]
CONTIG267	11890640_f3_12	420	6082	471	157	650	1.1(10)-63	Escherichia coli	b3863	[pn:dna polymerase i] [gn:polA]
CONTIG267	5260155_c1_16	421	6083	663	221	878	5.4(10)-88	Escherichia coli	b3865	[pn:hypothetical protein] [gn:yiha]
CONTIG268	5324009_c1_15	422	6084	650	216	795	3.3(10)-79	Escherichia coli	b2572	[pn:sigma-e factor negative regulatory protein] [gn:rsea]
CONTIG268	20109425_c1_18	423	6085	1632	544	2419	2.7(10)-251	Escherichia coli	b2569	[pn:gtp-binding protein lepa] [gn:lepa]
CONTIG268	29979502_c2_22	424	6086	492	164	513	2.6(10)-49	Escherichia coli	b2570	[pn:sigma-e factor regulatory protein rsec] [gn:rsec]
CONTIG268	24317933_c3_23	425	6087	1023	341	1294	4.5(10)-132	Escherichia coli	b2571	[pn:sigma-e factor regulatory protein rseb precursor] [gn:rseb]
CONTIG269	2428216_f2_5	426	6088	243	81	94	0.0015	Archaeoglobus fulgidus	H69378	[pn:purine ntpase homolog]
CONTIG269	10269038_c2_22	427	6089	816	272	915	6.5(10)-92	Escherichia coli	b0570	[pn:hypothetical protein] [gn:ybcz]
CONTIG269	30111081_c2_23	428	6090	444	148	123	5.5(10)-8	Synechocystis sp.	S77018	[PN:hypothetical protein] [OR:Synechocystis sp.] [SR:PCC 6803,, PCC 6803] [SR:PCC 6803,,]
CONTIG269	12620775_c3_27	429	6091	702	234	342	3.3(10)-31	Escherichia coli	S70162	,
CONTIG27	6511700_f2_2	430	6092	396	132	192	2.5(10)-14	Haemophilus influenzae	HI0539	[pn:urease alpha subunit] [gn:urea]
CONTIG270	14313817_f1_4	431	6093	768	256	727	5.5(10)-72	Shigella flexneri	P07044	mercuric resistance operon regulatory protein.
CONTIG270	31284383_f1_5	432	6094	351	117	553	1.5(10)-53	Enterobacter aerogenes	U67194	or:enterobacter aerogenes pn:tnpa gn:tnpa le:10303 re:11307 di:complement
CONTIG270	31523431_f2_8	433	6095	3222	1074	4832	0	Escherichia coli	P03008	transposase for transposon tn3.

CONTIG270	32204791_c1_14	434	6096	399	133	628	1.7(10)-61	Enterobacteria ceae	S07447	[pn:hypothetical protein, 13.1k]
CONTIG270	1057266_c1_18	435	6097	438	146	122	4.7(10)-7	Streptomyces fradiae	P20188	hypothetical 44.4 kd protein in transposon tn4556.
CONTIG270	22550252_c1_19	436	6098	639	213	684	2.0(10)-67	Escherichia coli	P03011	transposon tn3 resolvase.
CONTIG270	25522168_c2_20	437	6099	297	99	165	2.0(10)-12	Shigella flexneri	P04337	hypothetical mercuric resistance protein merc.
CONTIG271	16820950_f1_1	438	6100	702	234	1040	3.7(10)-105	Escherichia coli	b0437	[pn:atp-dependent clp protease proteolytic subunit] [gn:clpp]
CONTIG271	183188_f2_3	439	6101	1332	444	1904	1.0(10)-196	Escherichia coli	b0438	[pn:atp-dependent clp protease atp- binding subunit clpx] [gn:clpx]
CONTIG271	2500431_f3_7	440	6102	1695	565	2578	3.8(10)-268	Escherichia coli	b0439	[pn:lon protease] [gn:lon]
CONTIG271	35736527_c1_10	441	6103	225	75	99	4.0(10)-5	Bacillus subtilis	Z75208	or:bacillus subtilis pn:hypothetical protein gn:ysoc le:80592 re:81206 di:complement nt:unknown function; putative
CONTIG271	11067152_c1_11	442	6104	192	64	112	1.3(10)-6	Bacillus subtilis	Z75208	or:bacillus subtilis pn:hypothetical protein gn:ysoc le:80592 re:81206 di:complement nt:unknown function; putative
CONTIG272	36595388_f1_4	443	6105	483	161	113	6.2(10)-7	Escherichia coli	b1031	[pn:hypothetical protein] [gn:yedv]
CONTIG272	16526588_f2_7	444	6106	1119	373	1440	1.5(10)-147	Escherichia coli	b1020	[pn:phoh protein] [gn:phoh]
CONTIG272	36620786_f2_8	445	6107	186	62	117	2.3(10)-7	Escherichia coli	b1031	[pn:hypothetical protein] [gn:yedv]
CONTIG272	16853317_f3_9	446	6108	270	90	310	8.4(10)-28	Escherichia coli	b1017	[pn:hypothetical protein]
CONTIG272	13851425_f3_10	447	6109	1155	385	1589	2.5(10)-163	Escherichia coli	b1018	[pn:hypothetical protein]
CONTIG272	25978458_f3_11	448	6110	1287	429	1830	7.0(10)-189	Escherichia coli	b1019	[pn:hypothetical protein in phoh 5' region] [gn:yedv]
CONTIG272	3251026_f3_13	449	6111	312	104	106	3.5(10)-6	Escherichia coli	b1031	[pn:hypothetical protein] [gn:yedv]
CONTIG273	5370841_f1_1	450	6112	1809	603	2802	7.0(10)-292	Escherichia coli	b0680	[pn:glutaminyl-trna synthetase] [gn:glns]

CONTIG273	13931505_f1_2	451	6113	615	205	948	2.1(10)-95	Escherichia coli	b0681	[pn:hypothetical protein]
CONTIG273	267062_f3_12	452	6114	2001	667	2384	1.3(10)-247	Escherichia coli	b0679	[pn:pts system, n-acetylglucosamine-specific iabc component] [gn:nage]
CONTIG273	26659787_c3_23	453	6115	861	287	90	0.024	Homo sapiens	153641	mucin - human (fragment)
CONTIG274	35752258_f2_2	454	6116	1101	367	1209	4.5(10)-123	Escherichia coli	b0356	[pn:alcohol--acetaldehyde dehydrogenase] [gn:adhc]
CONTIG274	20490962_f3_5	455	6117	312	104	243	1.1(10)-20	Escherichia coli	b0357	[pn:hypothetical protein]
CONTIG274	32594790_f3_8	456	6118	312	104	331	5.0(10)-30	Haemophilus influenzae	H10185	[pn:putative alcohol dehydrogenase class iii] [gn:adhc]
CONTIG274	24234380_f3_9	457	6119	1551	517	1357	9.5(10)-139	Escherichia coli	b1421	[pn:methyl-accepting chemotaxis protein iii] [gn:trg]
CONTIG274	4837_c2_18	458	6120	501	167	419	2.3(10)-39	Escherichia coli	b1931	[pn:yedg] [gn:yedk]
CONTIG274	31532686_c2_19	459	6121	333	111	486	1.8(10)-46	Escherichia coli	b1931	[pn:yedg] [gn:yedk]
CONTIG275	9892787_f1_1	460	6122	429	143	582	1.3(10)-56	Escherichia coli	b0240	[pn:curlin genes transcriptional activator] [gn:crl]
CONTIG275	31355201_f2_4	461	6123	507	169	731	2.1(10)-72	Escherichia coli	b0239	[pn:hypothetical protein in gpt-crl intergenic region] [gn:yafa]
CONTIG275	866576_f3_10	462	6124	1143	381	1784	5.4(10)-184	Escherichia coli	b0242	[pn:glutamate 5-kinase] [gn:prob]
CONTIG275	3906666_c1_13	463	6125	1080	360	1533	2.1(10)-157	Escherichia coli	b0241	[pn:outer membrane pore protein e precursor] [gn:phoe]
CONTIG276	20604691_f1_2	464	6126	732	244	856	1.2(10)-85	Escherichia coli	b0192	[pn:copper homeostasis protein precursor] [gn:cuf]
CONTIG276	9961630_f2_8	465	6127	453	151	492	4.4(10)-47	Escherichia coli	b0191	[pn:hypothetical protein in acca-cuf intergenic region] [gn:yaef]
CONTIG276	34648525_f3_14	466	6128	564	188	842	3.5(10)-84	Escherichia coli	b0190	[pn:hypothetical protein in acca-cuf intergenic region] [gn:yaef]
CONTIG276	24725817_c1_20	467	6129	183	61	290	1.1(10)-25	Escherichia coli	b0196	[pn:rcsf protein] [gn:rcsf]
CONTIG276	16100083_c1_21	468	6130	294	98	93	8.5(10)-5	Escherichia coli	C47040	orf3 3' to rcsf - escherichia coli
CONTIG276	907_c1_22	469	6131	1785	595	2738	4.2(10)-285	Escherichia coli	b0194	[pn:prolyl-tma synthetase] [gn:pros]
CONTIG276	3236528_c2_30	470	6132	225	75	202	2.2(10)-16	Escherichia coli	P52099	hypothetical 7.2 kd protein in mesj-cuf intergenic region.



CONTIG276	26066886_c3_31	471	6133	720	240	1037	7.7(10)-105	Escherichia coli	b0195	[pn:hypothetical 26.4 kd protein in pros-rsf intergenic region]
CONTIG277	29886588_c1_14	472	6134	441	147	730	2.6(10)-72	Serratia marcescens	U59129	or:serratia marcescens pn:htdf gn:htdf le:29 re:454 di:direct
CONTIG277	26235000_c2_16	473	6135	3678	1226	92	0.68	Anas platyrhynchos	U27213	or:anas platyrhynchos pn:igm heavy chain, constant region domains one, gn:mu le:<1 re:1350 di:direct sr:domestic duck nt:presumed antibody; immunoglobulin heavy chain
CONTIG277	3937825_c3_17	474	6136	480	160	755	5.9(10)-75	Plasmid R478	A55209	h transfer determinant a - plasmid r478
CONTIG277	32687777_c3_18	475	6137	597	199	904	9.5(10)-91	Serratia marcescens	U59129	or:serratia marcescens pn:htdk gn:htdk le:463 re:996 di:direct
CONTIG278	1362586_f1_4	476	6138	519	173	249	2.3(10)-21	Vibrio cholerae	S81006	or:vibrio cholerae pn:hcp gn:hcp le:690 re:1208 di:direct sr:vibrio cholerae o17 nt:28 kda secreted hydrophilic protein; this sequence
CONTIG278	3333152_c2_17	477	6139	1284	428	1901	2.1(10)-196	Escherichia coli	b1184	[pn:umuc protein] [gn:umuc]
CONTIG278	21759555_c3_18	478	6140	474	158	568	3.7(10)-55	Escherichia coli	b1183	[pn:umud protein] [gn:umud]
CONTIG279	995266_c1_16	479	6141	948	316	122	2.7(10)-6	Methanococcus jannaschii	MJ1494	[pn:aaa superfamily atpase, similar to ftsh]
CONTIG279	2143887_c2_23	480	6142	522	174	604	5.9(10)-59	Escherichia coli	AJ224995	[de:escherichia coli ecorii restriction endonuclease gene.] [pn:restriction endonuclease] [gn:ecorii]
CONTIG28	476381_f1_1	481	6143	243	81	173	2.6(10)-12	Escherichia coli	b4065	[pn:hypothetical 60.5 kd protein in soxr-acs intergenic region] [gn:yice]
CONTIG28	29728382_f2_2	482	6144	315	105	133	5.2(10)-8	Escherichia coli	b4065	[pn:hypothetical 60.5 kd protein in soxr-acs intergenic region] [gn:yice]
CONTIG280	23609515_f3_13	483	6145	294	98	419	2.3(10)-39	Enterobacter agglomerans	A38965	hypothetical protein a (insertion sequence is1222) -enterobacter agglomerans

CONTIG280	36211006_c1_14	484	6146	2109	703	107	0.028	Legionella pneumophila	AF026534	[de:legionella pneumophila dot region ii, cita, dote, dotn, dote, dotf, dotg, doth, doti, dotj, dotk, dotl, and dotm genes, complete cds.] [pn:dotl] [gn:dotl] [nt:similar to trbc from the inci plasmid r64.]
CONTIG280	14538317_c2_15	485	6147	2613	871	104	0.03799	Listeria monocytogenes	M80351	or: listeria monocytogenes pn:p60-related protein le:1 re:1437 di:direct sr: listeria monocytogenes (serovar 1/2a mackness) dna
CONTIG281	21567780_f2_8	486	6148	768	256	755	5.9(10)-75	Escherichia coli	b4190	[pn:hypothetical 27.6 kd protein in aidb-rpsf intergenic region] [gn:yjfp]
CONTIG281	26681526_c1_13	487	6149	759	253	1187	9.8(10)-121	Escherichia coli	b4191	[pn:hypothetical transcriptional regulator in aidb-rpsf intergenic region] [gn:yjfq]
CONTIG281	4392281_c1_15	488	6150	231	77	95	0.00025	Nannocystis exedens	U66220	or: nannocystis exedens pn:unknown le:<1 re:872 di:complement nt:orf1
CONTIG281	23609702_c2_16	489	6151	447	149	408	3.5(10)-38	Escherichia coli	b4189	[pn:hypothetical 16.0 kd protein in aidb-rpsf intergenic region] [gn:yjfo]
CONTIG281	32501253_c3_23	490	6152	333	111	309	1.1(10)-27	Escherichia coli	b4188	[pn:hypothetical 11.0 kd protein in aidb-rpsf intergenic region] [gn:yjfn]
CONTIG282	17080463_f2_12	491	6153	1182	394	1562	1.8(10)-160	Escherichia coli	b4322	[pn:mannonate hydratase] [gn:uxua]
CONTIG282	20752305_c1_18	492	6154	1482	494	2008	9.8(10)-208	Escherichia coli	b2172	[pn:hypothetical 54.0 kd protein in fruk 5'''region] [gn:yeiq]
CONTIG282	13719178_c1_19	493	6155	1026	342	1336	1.6(10)-136	Escherichia coli	b2173	[pn:hypothetical 36.1 kd protein in frub-rtn intergenic region] [gn:yeir]
CONTIG282	895125_c1_20	494	6156	735	245	744	8.5(10)-74	Escherichia coli	b2174	[pn:hypothetical protein]
CONTIG283	32551313_f1_1	495	6157	369	123	362	2.6(10)-33	Escherichia coli	b3228	[pn:stringent starvation protein b] [gn:sspb]
CONTIG283	23850927_f1_3	496	6158	2436	812	1534	1.7(10)-157	Escherichia coli	b3216	[pn:hypothetical outer membrane usher protein in gltf-nant intergenic region] [gn:yhcd]
CONTIG283	4875010_f2_6	497	6159	816	272	565	8.0(10)-55	Escherichia coli	b3215	[pn:hypothetical 25.3 kd fimbrial chaperone in gltf-nant intergenic region] [gn:yhca]
CONTIG283	24415941_f3_8	498	6160	681	227	242	1.3(10)-20	Escherichia coli	b3219	[pn:hypothetical protein] [gn:yhcf]

CONTIG283	24415936_f3_10	499	6161	282	94	160	6.5(10)-12	Escherichia coli	b3219	[pn:hypothetical protein] [gn:yhcf]
CONTIG284	14081275_f1_1	500	6162	228	76	115	2.2(10)-6	Escherichia coli	b1514	[pn:hypothetical protein]
CONTIG284	23464702_f1_2	501	6163	978	326	771	1.2(10)-76	Escherichia coli	b1515	[pn:hypothetical protein]
CONTIG284	22945418_f1_6	502	6164	978	326	1134	4.0(10)-115	Escherichia coli	b1517	[pn:hypothetical protein] [gn:yneb]
CONTIG284	16130028_f2_8	503	6165	1056	352	1315	2.7(10)-134	Escherichia coli	b1516	[pn:hypothetical protein]
CONTIG284	31845937_f3_13	504	6166	543	181	353	2.2(10)-32	Escherichia coli	b1518	[pn:hypothetical protein]
CONTIG285	23922002_f1_2	505	6167	1338	446	1837	1.3(10)-189	Escherichia coli	b1423	[pn:hypothetical protein]
CONTIG285	31753501_c3_22	506	6168	960	320	1315	2.7(10)-134	Escherichia coli	b1422	[pn:hypothetical protein]
CONTIG286	7207808_f2_9	507	6169	1038	346	1460	1.2(10)-149	Escherichia coli	b1097	[pn:hypothetical 38.2 kd protein in pabc-holb intergenic region] [gn:yceg]
CONTIG286	26303760_f2_11	508	6170	759	253	822	4.7(10)-82	Escherichia coli	b1099	[pn:dna polymerase iii, delta"" subunit] [gn:holb]
CONTIG286	84378_f3_12	509	6171	1299	433	1922	1.3(10)-198	Escherichia coli	b1095	[pn:3-oxoacyl-acyl-carrier-protein synthase ii] [gn:fabf]
CONTIG286	4019818_f3_13	510	6172	819	273	864	1.7(10)-86	Escherichia coli	b1096	[pn:4-amino-4-deoxychorismate lyase] [gn:pabc]
CONTIG286	33988967_f3_15	511	6173	660	220	904	9.5(10)-91	Escherichia coli	b1098	[pn:hypothetical protein in pabc-holb intergenic region] [gn:tmk]
CONTIG287	20525451_f1_1	512	6174	1008	336	1549	4.2(10)-159	Escherichia coli	b1779	[pn:glyceraldehyde 3-phosphate dehydrogenase a] [gn:gapa]
CONTIG287	29511057_f3_8	513	6175	900	300	1148	1.3(10)-116	Escherichia coli	b1780	[pn:hypothetical protein]
CONTIG287	36135312_f3_11	514	6176	1574	525	2554	1.3(10)-265	Escherichia coli	b1783	[pn:hypothetical protein]
CONTIG287	20125050_c2_19	515	6177	774	258	934	6.2(10)-94	Escherichia coli	b1782	[pn:hypothetical protein]
CONTIG288	206649_c1_21	516	6178	184	61	183	2.3(10)-14	Escherichia coli	b3179	[pn:cell division protein ftsj] [gn:ftsj]
CONTIG288	2187917_c1_23	517	6179	933	311	1346	1.3(10)-137	Escherichia coli	b3177	[pn:dihydropteroate synthase] [gn:folp]

CONTIG288	34038453_c2_25	518	6180	288	96	374	1.3(10)-34	Escherichia coli	b3179	[pn:cell division protein ftsj] [gn:ftsj]
CONTIG288	23495385_c2_26	519	6181	1959	653	2979	0	Escherichia coli	b3178	[pn:cell division protein ftsj] [gn:ftsj]
CONTIG288	6491462_c2_28	520	6182	1125	375	1407	4.7(10)-144	Escherichia coli	b3176	[pn:mrsa protein] [gn:mrsa]
CONTIG289	14338511_f3_21	521	6183	2178	726	522	3.1(10)-49	Achlya klebsiana	P41755	nad-specific glutamate dehydrogenase (ec 1.4.1.2) (nad-gdh)
CONTIG289	20183505_c1_25	522	6184	687	229	433	7.7(10)-41	Escherichia coli	b0019	[pn:na] [gn:nhaa]
CONTIG289	4877327_c2_27	523	6185	1914	638	2830	7.7(10)-295	Escherichia coli	b0014	[pn:dnak protein] [gn:dnak]
CONTIG289	13672255_c2_28	524	6186	1011	337	922	1.2(10)-92	Escherichia coli	b0019	[pn:na] [gn:nhaa]
CONTIG289	26658142_c2_30	525	6187	987	329	1437	3.2(10)-147	Escherichia coli	b0020	[pn:transcriptional activator protein nhar] [gn:nhar]
CONTIG289	24471938_c3_35	526	6188	1116	372	1243	3.0(10)-146	Escherichia coli	b0015	[pn:dnaj protein] [gn:dnaj]
CONTIG290	9896067_f1_4	527	6189	318	106	273	7.0(10)-24	Enterobacter agglomerans	B38965	hypothetical protein b (insertion sequence is1222) -enterobacter agglomerans
CONTIG290	23609515_f2_8	528	6190	294	98	419	2.3(10)-39	Enterobacter agglomerans	A38965	hypothetical protein a (insertion sequence is1222) -enterobacter agglomerans
CONTIG290	22905462_f3_15	529	6191	561	187	866	1.0(10)-86	Enterobacter agglomerans	B38965	hypothetical protein b (insertion sequence is1222) -enterobacter agglomerans
CONTIG290	24500893_c2_21	530	6192	2418	806	1110	0.012	Escherichia coli	AF005044	[PN:TraN] [GN:traN] [DE:Escherichia coli plasmid R100-1 TraV (traV), TraR (traR), OrfG1(orfG1), OrfH (orfH), OrfI (orfI), TraC (traC), TrbI (trbI), TraW(traW), TraU (traU), TrbC (trbC), TraN (traN), TrbE (trbE) and TraF(traF) genes, c
CONTIG290	32225050_c2_24	531	6193	1680	560	286	5.5(10)-22	Bacillus subtilis	yjcD	[pn:hypothetical protein]
CONTIG291	20370300_f2_10	532	6194	321	107	113	2.8(10)-6	Rhizobium sp.	P55426	hypothetical 34.2 kd protein y4ec,

CONTIG291	1050702_f3_13	533	6195	900	300	95	0.04	Bacillus subtilis	ripX	[pn:integrase/recombinase] [gn:ykqm]
CONTIG291	33879001_f3_17	534	6196	729	243	324	8.4(10)-28	Escherichia coli	P27190	dna primase trac (ec 2.7.7.-) (replication primase).
CONTIG291	16020063_c3_33	535	6197	192	64	259	2.1(10)-22	Enterobacter aerogenes	U67194	or:enterobacter aerogenes pn:tnpa gn:tnpa le:13384 re:14388 di:direct
CONTIG292	4114091_f2_10	536	6198	480	160	439	1.8(10)-41	Escherichia coli	M16489	or:escherichia coli le:627 re:1199 di:complement sr:escherichia coli dna nt:orf 4; putative
CONTIG292	24611566_c1_26	537	6199	450	150	644	3.3(10)-63	Escherichia coli	b0736	[pn:hypothetical 15.6 kd protein in cydb-tolq intergenic region] [gn:ybgc]
CONTIG292	802151_c1_28	538	6200	1128	376	211	2.8(10)-21	Escherichia coli	b0739	[pn:protein] [gn:tola]
CONTIG292	1190692_c2_29	539	6201	1515	505	2417	4.5(10)-251	Escherichia coli	b0733	[pn:cytochrome d ubiquinol oxidase subunit i] [gn:cyda]
CONTIG292	21761538_c2_30	540	6202	1155	385	1632	6.7(10)-168	Escherichia coli	b0734	[pn:cytochrome d ubiquinol oxidase subunit ii] [gn:cydb]
CONTIG292	2867192_c3_38	541	6203	303	101	343	2.7(10)-31	Escherichia coli	b0735	[pn:10.9 kd protein in cydb-tolq intergenic region] [gn:ybge]
CONTIG292	25509808_c3_39	542	6204	696	232	1068	4.0(10)-108	Escherichia coli	b0737	[pn:tolq protein] [gn:tolq]
CONTIG292	33728407_c3_40	543	6205	432	144	529	5.2(10)-51	Escherichia coli	b0738	[pn:tolr protein] [gn:tolr]
CONTIG293	20875925_f2_6	544	6206	903	301	1407	4.7(10)-144	Escherichia coli	b3073	[pn:probable ornithine aminotransferase] [gn:yggg]
CONTIG293	24728780_f3_14	545	6207	576	192	844	2.2(10)-84	Escherichia coli	b3073	[pn:probable ornithine aminotransferase] [gn:yggg]
CONTIG293	25509681_c1_20	546	6208	606	202	126	1.7(10)-7	Escherichia coli	b1434	[pn:hypothetical protein]
CONTIG294	10353302_f1_2	547	6209	414	138	421	1.5(10)-39	Escherichia coli	b0530	[pn:hypothetical protein] [gn:sfma]
CONTIG294	13947040_f1_3	548	6210	612	204	558	4.4(10)-54	Salmonella typhi	Q08456	fimbriin-like protein fimi.
CONTIG294	5963891_f1_5	549	6211	2586	862	2988	0	Escherichia coli	b0532	[pn:hypothetical protein] [gn:sfmd]
CONTIG294	25487750_f2_10	550	6212	738	246	660	6.9(10)-65	Escherichia coli	b0531	[pn:hypothetical protein] [gn:sfmc]

CONTIG294	26845900_f2_14	551	6213	1104	368	1037	7.7(10)-105	Escherichia coli	b0533	[pn:hypothetical protein] [gn:sfmh]
CONTIG294	391957_f3_17	552	6214	240	80	104	8.5(10)-6	Escherichia coli	b0530	[pn:hypothetical protein] [gn:sfma]
CONTIG294	30338206_f3_23	553	6215	498	166	333	3.1(10)-30	Escherichia coli	b0534	[pn:hypothetical fimbrial-like protein in fimz 5''' region] [gn:ybog]
CONTIG294	32426319_c1_25	554	6216	503	167	156	1.8(10)-11	Salmonella typhimurium	P26318	fimbriae y protein.
CONTIG294	10417711_c1_37	555	6217	276	92	249	2.3(10)-21	Escherichia coli	b0529	[pn:methylenetetrahydrofolate dehydrogenase] [gn:fold]
CONTIG294	12697187_c3_44	556	6218	636	212	687	9.4(10)-68	Escherichia coli	b0535	[pn:fimbriae z protein] [gn:fimz]
CONTIG295	24647536_f1_6	557	6219	753	251	428	2.6(10)-40	Bacteriophage P21	P36693	terminase large subunit (gp2) (fragments).
CONTIG295	3960443_f1_7	558	6220	333	111	139	1.1(10)-9	Bacteriophage lambda	P03712	head decoration protein (gpd) (major capsid protein d).
CONTIG295	11207801_f2_16	559	6221	540	180	114	6.7(10)-7	Bacteriophage lambda	J02459	or:bacteriophage lambda le:5132 re:5737 di:direct sr:lambda wild-type and lambda strain ci857s7 nt:nu3 (capsid assembly;201)
CONTIG295	9970952_f2_17	560	6222	1035	345	972	5.9(10)-98	Bacteriophage lambda	P03713	major head protein (gpe) (major coat protein).
CONTIG295	36135213_f2_18	561	6223	444	148	100	1.5(10)-5	Bacteriophage lambda	P03709	dna packaging protein fi.
CONTIG296	6281375_f2_2	562	6224	990	330	701	3.1(10)-69	Haemophilus influenzae	H11144	[pn:udp-3-0-acyl n-acetylglucosamine deacetylase] [gn:lpxc]
CONTIG296	4942010_f2_3	563	6225	453	151	266	3.8(10)-23	Haemophilus influenzae	H10293	[pn:mercury resistance homolog]
CONTIG296	978425_f2_4	564	6226	933	311	711	6.9(10)-70	Escherichia coli	b0462	[pn:acriflavin resistance protein b] [gn:acrb]
CONTIG296	4145218_f3_5	565	6227	3639	1213	282	1.6(10)-34	Bacillus subtilis	yoaE	[pn:hypothetical protein]
CONTIG296	13797188_f3_6	566	6228	1299	433	505	1.8(10)-48	Escherichia coli	b0463	[pn:acriflavin resistance protein a precursor] [gn:acra]
CONTIG297	30347307_f2_10	567	6229	546	182	180	5.9(10)-14	Escherichia coli	b3558	[pn:insertion element is150 hypothetical 33.3 kd protein] [gn:yif5b]
CONTIG297	12948567_f3_13	568	6230	291	97	391	2.2(10)-36	Escherichia coli	I41306	hypothetical protein (argf-lacZ region) - escherichia coli

CONTIG297	1448266_c2_17	569	6231	2382	794	91	0.34	Candida albicans	P43060	phosphoribosylamidoimidazole-succinocarboxamide synthase (ec 6.3.2.6) (saicar synthetase).
CONTIG298	5971942_f1_1	570	6232	279	93	269	1.8(10)-23	Escherichia coli	b1648	[pn:hypothetical protein]
CONTIG298	2348136_f1_2	571	6233	921	307	1273	7.5(10)-130	Escherichia coli	b1647	[pn:hypothetical protein]
CONTIG298	2603262_f1_3	572	6234	573	191	789	1.5(10)-78	Escherichia coli	b1646	[pn:copper-zinc superoxide dismutase] [gn:sodc]
CONTIG298	884512_c1_23	573	6235	282	94	295	3.2(10)-26	Escherichia coli	b1643	[pn:hypothetical protein]
CONTIG298	25907180_c2_31	574	6236	2100	700	2323	4.0(10)-241	Escherichia coli	b1645	[pn:hypothetical protein]
CONTIG298	12634512_c3_35	575	6237	945	315	1096	4.2(10)-111	Escherichia coli	b1644	[pn:hypothetical protein]
CONTIG299	4151412_f1_1	576	6238	1200	400	152	2.8(10)-8	Bacteriophage P1	P06956	recombinase cre.
CONTIG299	10006261_c3_21	577	6239	1032	344	98	0.056	Saccharomyces cerevisiae	YGR155W	[pn:cystathionine beta-synthase] [gn:cys4]
CONTIG3	4863125_c3_6	578	6240	606	202	473	4.5(10)-45	Escherichia coli	b0698	[pn:potassium-transporting atpase, a chain] [gn:kdpA]
CONTIG30	29728277_f1_1	579	6241	303	101	229	9.4(10)-19	Escherichia coli	b1053	[pn:hypothetical 43.9 kd protein in msyb-htrb intergenic region] [gn:ycee]
CONTIG300	5163340_f1_2	580	6242	438	146	493	3.3(10)-47	Erwinia carotovora subsp. carotovora	JC4727	mob protein a - erwinia carotovora subsp. carotovora a cis-acting locus, orit and trans-acting locus, mob are involved in mobilization of pec3, a non-self-transmissible multiple-copy plasmid. the mob consists of five proteins.
CONTIG300	14866452_f3_5	581	6243	432	144	546	8.3(10)-53	Erwinia carotovora subsp. carotovora	JC4729	mob protein c - erwinia carotovora subsp. carotovora a cis-acting locus, orit and trans-acting locus, mob are involved in mobilization of pec3, a non-self-transmissible multiple-copy plasmid. the mob consists of five proteins.

CONTIG301	31755158_fl_1	582	6244	993	331	1496	1.8(10)-153	Escherichia coli	b3974	[pn:pantothenate kinase] [gn:coaa]
CONTIG301	33475302_c1_29	583	6245	1185	395	1621	1.0(10)-166	Escherichia coli	b3543	[pn:dipeptide transport system permease protein dppb] [gn:dppb]
CONTIG301	4332343_c1_30	584	6246	912	304	1144	3.5(10)-116	Escherichia coli	b3542	[pn:dipeptide transport system permease protein dppc] [gn:dppc]
CONTIG301	17011415_c1_32	585	6247	1032	344	1665	2.2(10)-171	Escherichia coli	b3540	[pn:dipeptide transport atp-binding protein dppf] [gn:dppf]
CONTIG301	20167543_c2_35	586	6248	228	76	392	1.7(10)-36	Escherichia coli	b3544	[pn:periplasmic dipeptide transport protein precursor] [gn:dppa]
CONTIG301	34564667_c2_37	587	6249	1002	334	1402	1.6(10)-143	Escherichia coli	b3541	[pn:dipeptide transport atp-binding protein dppd] [gn:dppd]
CONTIG302	13753811_fl_3	588	6250	1221	407	1664	2.7(10)-171	Escherichia coli	b2844	[pn:hypothetical protein]
CONTIG302	6445812_fl_4	589	6251	717	239	136	1.5(10)-7	Escherichia coli	b4311	[pn:hypothetical 28.3 kd protein in feci-fimb intergenic region]
CONTIG302	4461517_f2_6	590	6252	327	109	137	6.4(10)-9	Escherichia coli	b0208	[pn:hypothetical transcriptional regulator in rnh-dhnr intergenic region] [gn:yafc]
CONTIG302	3176081_f2_7	591	6253	555	185	91	0.041	Haemophilus influenzae	H10681	[pn:transcriptional activator protein] [gn:ilvy]
CONTIG302	2078535_c3_37	592	6254	276	92	338	9.0(10)-31	Escherichia coli	b2845	[pn:hypothetical protein]
CONTIG303	894181_fl_1	593	6255	1428	476	834	2.5(10)-83	Escherichia coli	b1525	[pn:hypothetical protein]
CONTIG303	32689752_c1_13	594	6256	1863	621	2460	1.2(10)-255	Salmonella enterica	X99719	or:salmonella enterica le:<142 re:1998 direct nt:similarity with helicase; orf1
CONTIG303	25986000_c3_29	595	6257	567	189	356	1.1(10)-32	Escherichia coli	b2250	[pn:hypothetical protein]
CONTIG304	22656568_f3_8	596	6258	222	74	359	5.4(10)-33	Escherichia coli	b1823	[pn:cold shock-like protein cspc] [gn:cspc]
CONTIG304	1308402_f3_9	597	6259	1779	593	1863	2.2(10)-192	Escherichia coli	b0084	[pn:penicillin-binding protein 3 precursor] [gn:fts]
CONTIG304	9879466_f3_10	598	6260	855	285	1136	2.5(10)-115	Escherichia coli	b1822	[pn:hypothetical 30.4 kd protein in manz-cspc intergenic region] [gn:yebh]
CONTIG304	12531889_c1_13	599	6261	425	141	561	2.1(10)-54	Escherichia coli	b1819	[pn:pts system, mannose-specific iid component] [gn:manz]



CONTIG304	14110327_c1_14	600	6262	459	153	602	9.5(10)-59	Escherichia coli	b1820	[pn:hypothetical protein]
CONTIG304	31770643_c3_26	601	6263	774	258	832	4.0(10)-83	Escherichia coli	b1821	[pn:hypothetical protein]
CONTIG305	1424025_f1_1	602	6264	204	68	101	0.00029	Porphyromonas gingivalis	P46071	protease prth (ec 3.4.22.-).
CONTIG305	34085002_f3_7	603	6265	3627	1209	560	5.2(10)-51	Acinetobacter calcoaceticus	AF011339	[PN:unknown] [DE:Acinetobacter calcoaceticus unknown protein gene, partial cds.] [LE:<1] [RE:2753] [DI:direct]
CONTIG305	6101512_c2_16	604	6266	570	190	172	3.5(10)-13	Escherichia coli	b2846	[pn:hypothetical protein]
CONTIG305	24254687_c2_17	605	6267	1806	602	310	8.5(10)-25	Escherichia coli	X86087	or:escherichia coli gn:hec-hlyb le:3628 re:5748 di:direct
CONTIG306	15750937_f1_1	606	6268	990	330	1061	2.2(10)-107	Escherichia coli	b2100	[pn:hypothetical protein]
CONTIG306	26351577_f2_7	607	6269	1005	335	1345	1.8(10)-137	Escherichia coli	b2099	[pn:hypothetical protein]
CONTIG306	29718840_f3_15	608	6270	1236	412	1733	1.3(10)-178	Escherichia coli	b2098	[pn:hypothetical protein]
CONTIG306	11885284_c1_20	609	6271	896	298	1152	5.0(10)-117	Escherichia coli	b2103	[pn:hypothetical protein] [gn:thid]
CONTIG306	29850408_c1_21	610	6272	789	263	1085	6.2(10)-110	Escherichia coli	b2101	[pn:hypothetical protein]
CONTIG307	4332807_f1_2	611	6273	1629	543	2360	4.9(10)-245	Escherichia coli	b2574	[pn:l-aspartate oxidase] [gn:nadb]
CONTIG307	5978412_f3_12	612	6274	1332	444	1704	1.6(10)-175	Escherichia coli	b2576	[pn:atp-dependent rna helicase srmb] [gn:srmb]
CONTIG307	22449093_c1_14	613	6275	396	132	599	2.0(10)-58	Escherichia coli	b2579	[pn:hypothetical 14.3 kd protein in srmb-ung intergenic region]
CONTIG307	32531250_c2_28	614	6276	1089	363	969	1.2(10)-97	Escherichia coli	b2575	[pn:hypothetical protein in nadb-srmb intergenic region] [gn:yfic]
CONTIG307	9817892_c3_36	615	6277	414	138	602	9.5(10)-59	Escherichia coli	b2573	[pn:rna polymerase sigma-e factor] [gn:rpoe]
CONTIG308	4306635_f2_9	616	6278	915	305	1404	9.9(10)-144	Escherichia coli	b2843	[pn:5-keto-4-deoxyuronate isomerase] [gn:kdui]
CONTIG308	4491568_f2_10	617	6279	771	257	1202	2.5(10)-122	Escherichia coli	b2842	[pn:2-deoxy-d-gluconate 3-dehydrogenase] [gn:kdud]

CONTIG308	3203441_f2_11	618	6280	1557	519	1989	1.0(10)-205	Escherichia coli	b2841	[pn:l-arabinose isomerase] [gn:arae]
CONTIG308	20038125_c3_30	619	6281	741	247	958	1.8(10)-96	Erwinia chrysanthemi	P21258	oligogalacturonate lyase (ec 4.2.2.6)
CONTIG309	25802285_f2_15	620	6282	861	287	696	1.1(10)-68	Mycobacterium tuberculosis	Z95121	unknown,,mtcy20b11.07c,mtcy20b11.07c. len
CONTIG309	15834716_f2_24	621	6283	930	310	978	1.3(10)-98	Escherichia coli	b1473	[pn:yddg]
CONTIG309	23443818_f2_25	622	6284	531	177	606	3.6(10)-59	Escherichia coli	b0553	[pn:outer membrane porin protein nmpe precursor] [gn:nmpc]
CONTIG309	4817250_c1_37	623	6285	282	94	284	9.0(10)-24	Escherichia coli	b1474	[pn:alpha subunit of formate dehydrogenase n] [gn:fdng]
CONTIG309	17083292_c1_38	624	6286	333	111	465	3.6(10)-43	Escherichia coli	b1474	[pn:alpha subunit of formate dehydrogenase n] [gn:fdng]
CONTIG309	14555411_c1_39	625	6287	2460	820	3784	0	Escherichia coli	b1474	[pn:alpha subunit of formate dehydrogenase n] [gn:fdng]
CONTIG309	13784625_c1_42	626	6288	717	239	803	4.7(10)-80	Escherichia coli	b1476	[pn:gamma subunit of formate dehydrogenase n]
CONTIG309	7054750_c2_52	627	6289	270	90	253	1.3(10)-21	Escherichia coli	b0396	[pn:araj protein precursor] [gn:araj]
CONTIG309	26604202_c3_57	628	6290	885	295	1459	1.5(10)-149	Escherichia coli	b1475	[pn:beta subunit of formate dehydrogenase n] [gn:fdnh]
CONTIG31	29877090_c1_7	629	6291	297	99	122	7.0(10)-8	unidentified	X06035	or:unidentified le:107 re:>319 di:direct nt:laci gene product (71 aa)
CONTIG31	32605456_c2_8	630	6292	453	151	226	6.7(10)-19	Haemophilus influenzae	HI0522	[pn:sp]
CONTIG311	21776426_f3_19	631	6293	483	161	233	1.2(10)-19	Haemophilus influenzae	HI1623	[pn:mercury resistance regulatory protein]
CONTIG311	4494003_c1_26	632	6294	468	156	276	3.3(10)-24	Escherichia coli	b0487	[pn:hypothetical protein] [gn:ybbi]
CONTIG311	3959387_c1_27	633	6295	243	81	151	5.9(10)-11	Escherichia coli	b1531	[pn:multiple antibiotic resistance protein] [gn:mara]
CONTIG311	3986268_c2_28	634	6296	1641	547	778	2.1(10)-77	Bacillus subtilis	yvgX	[pn:hypothetical protein]
CONTIG311	26069675_c2_30	635	6297	2556	852	1087	3.8(10)-110	Escherichia coli	b3469	[pn:hypothetical protein] [gn:yhho]
CONTIG311	32714701_c3_31	636	6298	948	316	608	2.2(10)-59	Bacillus subtilis	yvgX	[pn:hypothetical protein]

CONTIG312	34551934_c2_38	637	6299	866	288	1314	3.3(10)-134	Escherichia coli	b1851	[pn:phosphogluconate dehydratase] [gn:edd]
CONTIG312	5082913_c2_39	638	6300	681	227	1022	3.0(10)-103	Escherichia coli	b1850	[pn:2-keto-3-deoxygluconate 6-p aldolase]
CONTIG312	10975301_c2_40	639	6301	2520	840	261	6.5(10)-19	Escherichia coli	AF044503	[de:escherichia coli strain ec11 unknown (498), hcp gene, complete cds;and rhsg accessory genetic element vrg protein, core component anddsorf-g1 genes, complete cds.] [pn:vrg protein]
CONTIG313	2082812_fl_1	640	6302	1311	437	1695	1.3(10)-174	Escherichia coli	b1650	[pn:hypothetical protein] [gn:nema]
CONTIG313	2828405_fl_2	641	6303	435	145	661	5.4(10)-65	Escherichia coli	b1651	[pn:hypothetical protein] [gn:gloa]
CONTIG313	5097838_fl_3	642	6304	669	223	887	6.0(10)-89	Escherichia coli	b1652	[pn:ribonuclease t] [gn:rt]
CONTIG313	5093876_fl_5	643	6305	849	283	749	2.5(10)-74	Escherichia coli	b1655	[pn:hypothetical protein] [gn:ydh]
CONTIG313	4589017_fl_10	644	6306	609	203	840	5.7(10)-84	Escherichia coli	b1649	[pn:hypothetical protein]
CONTIG313	13870968_fl_19	645	6307	1767	589	1323	3.7(10)-135	Escherichia coli	b4114	[pn:hypothetical 61.7 kd protein in bass-ady intergenic region]
CONTIG313	6735627_fl_21	646	6308	822	274	862	2.7(10)-86	Escherichia coli	b1656	[pn:superoxide dismutase] [gn:sodb]
CONTIG313	20114567_c1_22	647	6309	414	138	374	1.3(10)-34	Escherichia coli	b1657	[pn:hypothetical protein]
CONTIG313	26354717_c3_44	648	6310	360	120	593	8.5(10)-58	Escherichia coli	b1654	[pn:hypothetical protein in hr 5'''region] [gn:ydh]
CONTIG313	35411430_c3_46	649	6311	633	211	90	0.066	Rhodobacter sphaeroides	A57140	motb protein homolog - rhodobacter sphaeroides
CONTIG314	4332768_c2_19	650	6312	231	77	224	1.1(10)-18	Escherichia coli	b3267	[pn:hypothetical 7.5 kd protein in acrf-rnd intergenic region]
CONTIG314	12269541_c3_20	651	6313	2973	991	3597	0	Escherichia coli	b3266	[pn:acrf] [gn:acrf]
CONTIG315	9786501_fl_11	652	6314	495	165	451	9.5(10)-43	Escherichia coli	b1668	[pn:hypothetical protein]
CONTIG315	26306432_c2_43	653	6315	318	106	185	1.5(10)-14	Escherichia coli	b3049	[pn:glycogen synthesis protein glgs] [gn:glgs]

CONTIG315	1959386_c3_45	654	6316	522	174	250	1.8(10)-21	Escherichia coli	AF044503	[de:escherichia coli strain ec11 unknown (498), hcp gene, complete cds;and rhsg accessory genetic element vrg protein, core component anddsorf-g1 genes, complete cds.] [pn:unknown] [gn:498]
CONTIG315	2470468_c3_51	655	6317	438	146	174	2.2(10)-13	Escherichia coli	b1419	[pn:hypothetical protein] [gn:ydca]
CONTIG315	23729187_c3_52	656	6318	543	181	212	2.0(10)-17	Escherichia coli	b3686	[pn:hsls] [gn:ibpb]
CONTIG316	36572802_fl_1	657	6319	669	223	998	1.0(10)-100	Escherichia coli	b1040	[pn:putative regulatory protein] [gn:csgd]
CONTIG316	14568878_fl_2	658	6320	450	150	569	3.0(10)-55	Escherichia coli	b1038	[pn:assembly/transport component in curli production] [gn:csgf]
CONTIG316	4881326_f2_11	659	6321	450	150	555	9.1(10)-54	Escherichia coli	b1039	[pn:assembly/transport component in curli production] [gn:csgl]
CONTIG316	24022000_f3_19	660	6322	867	289	1323	3.7(10)-135	Escherichia coli	b1037	[pn:assembly/transport component in curli production] [gn:csgg]
CONTIG316	24407841_c1_34	661	6323	567	189	808	1.3(10)-80	Escherichia coli	b1035	[pn:hypothetical protein] [gn:yedz]
CONTIG316	22351626_c1_35	662	6324	579	193	720	3.0(10)-71	Escherichia coli	b1036	[pn:hypothetical protein] [gn:yedz]
CONTIG316	22781591_c2_37	663	6325	801	267	277	2.6(10)-24	Haemophilus influenzae	HI1364	[pn:hypothetical protein]
CONTIG316	35984376_c2_38	664	6326	435	145	249	2.3(10)-21	Escherichia coli	b1031	[pn:hypothetical protein] [gn:yedv]
CONTIG316	14882337_c3_47	665	6327	951	317	1285	4.0(10)-131	Escherichia coli	b1033	[pn:hypothetical protein] [gn:yedw]
CONTIG316	475201_c3_48	666	6328	774	258	1079	2.7(10)-109	Escherichia coli	b1034	[pn:hypothetical protein] [gn:yedx]
CONTIG317	23937778_f2_8	667	6329	1527	509	591	1.3(10)-57	Escherichia coli	b2155	[pn:colicin i receptor precursor] [gn:cira]
CONTIG317	781555_f2_15	668	6330	1104	368	1339	7.7(10)-137	Escherichia coli	b2158	[pn:hypothetical 36.9 kd protein in lysp-nfo intergenic region] [gn:yeih]
CONTIG317	7281308_f3_23	669	6331	873	291	1163	3.3(10)-118	Escherichia coli	b2159	[pn:endonuclease iv] [gn:nfo]

CONTIG317	22691558_c1_26	670	6332	873	291	1169	7.9(10)-119	Escherichia coli	b2157	[pn:hypothetical transcriptional regulator in lysp- nfo intergenic region] [gn:yeic]
CONTIG317	3922542_c3_41	671	6333	1557	519	2396	7.5(10)-249	Escherichia coli	b2156	[pn:lysine-specific permease] [gn:lysp]
CONTIG318	23722842_f1_4	672	6334	609	203	856	1.2(10)-85	Serratia marcescens	U60283	or:serratia marcescens pn:restriction methylase gn:trag1 le:84 re:1775 di:direct nt:putative restriction methylase
CONTIG318	4351033_f2_9	673	6335	1173	391	1857	9.8(10)-192	Serratia marcescens	U60283	or:serratia marcescens pn:restriction methylase gn:trag1 le:84 re:1775 di:direct nt:putative restriction methylase
CONTIG318	30273558_f3_14	674	6336	1188	396	101	0.012	Pseudomonas putida	P31857	hypothetical 32.4 kd protein in gidb- unci intergenic region.
CONTIG318	10172502_c2_28	675	6337	858	286	424	7.0(10)-40	Haemophilus influenzae	H10209	[pn:dna adenine methylase] [gn:dam]
CONTIG319	14486288_c1_28	676	6338	597	199	116	1.8(10)-5	Escherichia coli	b0535	[pn:fimbriae z protein] [gn:fimz]
CONTIG319	24648952_c1_29	677	6339	7962	2654	1058	3.7(10)-128	Escherichia coli	b1509	[pn:hypothetical protein]
CONTIG320	25492191_f1_3	678	6340	1248	416	1489	9.6(10)-153	Escherichia coli	b2907	[pn:ubih protein] [gn:ubih]
CONTIG320	14850443_f1_4	679	6341	1215	405	1730	2.7(10)-178	Escherichia coli	b2906	[pn:visc protein] [gn:visc]
CONTIG320	2596033_f1_6	680	6342	426	142	630	1.0(10)-61	Escherichia coli	b2904	[pn:glycine cleavage system h protein] [gn:gcvh]
CONTIG320	1291416_f2_7	681	6343	1206	402	1832	4.4(10)-189	Escherichia coli	b2908	[pn:proline aminopeptidase ii] [gn:pepp]
CONTIG320	1289712_f2_9	682	6344	1170	390	1730	2.7(10)-178	Escherichia coli	b2905	[pn:aminomethyltransferase] [gn:gcvt]
CONTIG321	24666092_f2_17	683	6345	1035	345	1509	7.4(10)-155	Escherichia coli	b2151	[pn:mgl repressor and galactose ultrainduction factor] [gn:gals]
CONTIG321	9770002_f3_20	684	6346	1179	393	262	6.7(10)-22	Escherichia coli	b4332	[pn:hypothetical 41.4 kd protein in iada-merd intergenic region] [gn:yij]
CONTIG321	19743927_f3_21	685	6347	759	253	883	1.6(10)-88	Escherichia coli	b2153	[pn:gtp cyclohydrolase i] [gn:fole]
CONTIG321	17067305_f3_22	686	6348	1176	392	1277	2.7(10)-130	Escherichia coli	b2152	[pn:hypothetical 43.4 kd protein in gals-fole intergenic region] [gn:yeib]

CONTIG321	15712632_f3_23	687	6349	966	322	1479	1.1(10)-151	Escherichia coli	b2150	[pn:d-galactose-binding periplasmic protein precursor] [gn:mglb]
CONTIG321	4143937_c1_29	688	6350	879	293	1253	9.9(10)-128	Escherichia coli	b2154	[pn:hypothetical 31.3 kd protein in fole-cira intergenic region] [gn:yeig]
CONTIG321	7078550_c2_37	689	6351	663	221	95	0.01099	Synechococcus PCC7942	U59236	or:synechococcus pcc7942 pn:unknown le:3661 re:4344 di:complement nt:orf227
CONTIG321	5963902_c2_41	690	6352	666	222	230	2.7(10)-18	Escherichia coli	b2155	[pn:colicin i receptor precursor] [gn:cira]
CONTIG322	26855467_f2_9	691	6353	612	204	715	1.0(10)-70	Escherichia coli	b1857	[pn:31.1 kd protein in msbb-ruvb intergenic region] [gn:yebi]
CONTIG322	25664816_f2_10	692	6354	1335	445	1985	2.7(10)-205	Escherichia coli	b1856	[pn:hypothetical 46.7 kd protein in msbb-ruvb intergenic region] [gn:yeba]
CONTIG322	6535437_f2_11	693	6355	216	72	161	1.6(10)-11	Escherichia coli	b1855	[pn:msbb protein] [gn:msbb]
CONTIG322	13797676_f2_12	694	6356	858	286	1320	7.9(10)-135	Escherichia coli	b1855	[pn:msbb protein] [gn:msbb]
CONTIG322	26734465_f2_14	695	6357	771	257	892	1.8(10)-89	Escherichia coli	b1852	[pn:glucose 6-phosphate 1 dehydrogenase] [gn:zwf]
CONTIG322	80333_f3_27	696	6358	759	253	1123	5.9(10)-114	Escherichia coli	b1852	[pn:glucose 6-phosphate 1 dehydrogenase] [gn:zwf]
CONTIG322	24245760_c2_39	697	6359	942	314	1296	2.7(10)-132	Escherichia coli	b1853	[pn:hypothetical 32.0 kd protein in pyka-zwf intergenic region] [gn:yebk]
CONTIG322	13945816_c2_40	698	6360	1554	518	2232	1.8(10)-231	Escherichia coli	b1854	[pn:pyruvate kinase a] [gn:pyka]
CONTIG323	16823587_f3_8	699	6361	240	80	196	1.0(10)-15	Escherichia coli	b2809	[pn:hypothetical protein]
CONTIG323	10651717_c1_11	700	6362	498	166	760	1.7(10)-75	Plasmid pSW200	L42525	or:plasmid psw200 gn:mobb le:2345 re:2830 di:direct sr:plasmid psw200 dna
CONTIG323	26681536_c1_12	701	6363	213	71	270	1.5(10)-23	Plasmid pSW200	L42525	or:plasmid psw200 gn:mobb le:2834 re:3049 di:direct sr:plasmid psw200 dna
CONTIG323	5130252_c3_15	702	6364	336	112	455	3.6(10)-43	Plasmid pSW200	L42525	or:plasmid psw200 gn:moba le:1657 re:3156 di:direct sr:plasmid psw200 dna

CONTIG323	24783153_c3_16	703	6365	708	236	967	2.0(10)-97	Plasmid pSW200	L42525	or:plasmid psw200 gn:moba le:1657 re:3156 di:direct sr:plasmid psw200 dna
CONTIG324	24088337_f2_4	704	6366	3273	1091	276	1.5(10)-23	Escherichia coli	b1350	[pn:exodeoxyribonuclease viii] [gn:rece]
CONTIG324	3914843_f2_5	705	6367	1095	365	259	4.7(10)-44	Escherichia coli	JN0845	enterohemolysin 1 - escherichia coli
CONTIG324	22744032_f2_6	706	6368	216	72	244	8.3(10)-21	Escherichia coli	b1346	[pn:hypothetical protein]
CONTIG324	16143775_f3_8	707	6369	210	70	196	1.0(10)-15	Salmonella typhimurium	AF001386	[de:salmonella typhimurium prophage-like element gifsy-1, partialsequence.] [nt:orf-3; similar to orf sequence of e.coli ""rac""]
CONTIG324	23697218_f3_19	708	6370	1251	417	1525	1.5(10)-156	Escherichia coli	b1345	[pn:hypothetical protein]
CONTIG325	16298292_f1_4	709	6371	252	84	223	1.3(10)-18	Escherichia coli	b3184	[pn:hypothetical 35.0 kd protein in dacb-rpma intergenic region] [gn:yhbc]
CONTIG325	34588378_f1_5	710	6372	1191	397	1569	3.2(10)-161	Escherichia coli	b3183	[pn:hypothetical 43.3 kd gtp-binding protein in dacb-rpma intergenic region] [gn:yhbz]
CONTIG325	16437552_f2_13	711	6373	366	122	522	2.8(10)-50	Escherichia coli	b3186	[pn:50s ribosomal subunit protein 121] [gn:rplu]
CONTIG325	9777178_f2_14	712	6374	843	281	1077	4.4(10)-109	Escherichia coli	b3184	[pn:hypothetical 35.0 kd protein in dacb-rpma intergenic region] [gn:yhbc]
CONTIG325	16587811_f2_22	713	6375	480	160	670	6.0(10)-66	Escherichia coli	b3181	[pn:grea]
CONTIG325	23616702_f3_23	714	6376	258	86	418	3.0(10)-39	Escherichia coli	b3185	[pn:50s ribosomal subunit protein 127] [gn:rpma]
CONTIG325	21666317_c1_35	715	6377	1095	365	736	6.0(10)-73	Escherichia coli	b4112	[pn:sensor protein bass/pmrb] [gn:bass]
CONTIG325	31437785_c2_39	716	6378	375	125	461	8.4(10)-44	Escherichia coli	b3180	[pn:hypothetical 10.8 kd protein in ftsj-grea intergenic region] [gn:yhbv]
CONTIG325	24651051_c2_42	717	6379	669	223	530	4.0(10)-51	Escherichia coli	b4113	[pn:transcriptional regulatory protein basr/pmra] [gn:basr]
CONTIG325	3948541_c3_51	718	6380	1443	481	2243	1.2(10)-232	Escherichia coli	b3182	[pn:d-alanyl-d-alanine carboxypeptidase, fraction b] [gn:"]

CONTIG326	10994040_c1_37	719	6381	696	232	128	2.7(10)-17	Mycoplasma sp.	P43641	modification methylase muni (ec 2.1.1.72) (adenine-specific methyltransferase muni) (m.muni). [pn:hypothetical protein]
CONTIG326	32664142_c1_38	720	6382	768	256	503	3.0(10)-48	Escherichia coli	b1560	[pn:hypothetical protein]
CONTIG326	23708141_c2_45	721	6383	531	177	130	1.0(10)-8	Escherichia coli	b1147	[pn:hypothetical protein]
CONTIG326	2151943_c2_47	722	6384	570	190	184	1.8(10)-14	Escherichia coli	b2357	[pn:hypothetical protein] [gn:yfdn]
CONTIG326	24422952_c2_49	723	6385	417	139	210	3.2(10)-17	Escherichia coli	b0550	[pn:hypothetical protein] [gn:rus]
CONTIG326	32070417_c2_50	724	6386	456	152	279	1.6(10)-24	Escherichia coli	b0551	[pn:hypothetical protein] [gn:ybcq]
CONTIG326	4416592_c3_57	725	6387	936	312	136	5.2(10)-7	Bacteriophage SPP1	S43811	gene 38 protein - phage spp1
CONTIG326	35657508_c3_58	726	6388	375	125	178	8.1(10)-14	Escherichia coli	b4043	[pn:lexa] [gn:lexa]
CONTIG326	12298415_c3_60	727	6389	216	72	329	8.1(10)-30	Escherichia coli	b1560	[pn:hypothetical protein]
CONTIG327	22369627_f1_1	728	6390	1446	482	765	5.0(10)-76	Ralstonia solanacearum	A36929	virulence regulatory protein vsrb - pseudomonas solanacearum
CONTIG327	30746031_f1_3	729	6391	1386	462	1149	1.0(10)-116	Escherichia coli	b4052	[pn:replicative dna helicase] [gn:dnab]
CONTIG327	24072317_f3_15	730	6392	888	296	136	5.2(10)-7	Bacillus subtilis	soj	[pn:hypothetical protein]
CONTIG327	12511376_f3_16	731	6393	1725	575	95	0.11	Agrobacterium rhizogenes	P05683	possible replication protein b.
CONTIG327	33798336_f3_19	732	6394	804	268	91	0.02999	Hyphomicrobi um sp.	Y08074	[de:hyphomicrobium sp. mxaf gene, partial, strain b 69.] [pn:methanol dehydrogenase] [gn:mxaf] [nt:alpha- subunit]
CONTIG328	32505382_f1_5	733	6395	855	285	884	1.3(10)-88	Escherichia coli	b1226	[pn:respiratory nitrate reductase I delta chain] [gn:narj]
CONTIG328	36049181_f2_9	734	6396	1557	519	2592	1.3(10)-269	Escherichia coli	b1225	[pn:respiratory nitrate reductase I beta chain] [gn:narh]
CONTIG328	4771955_f3_12	735	6397	3840	1280	6414	0	Escherichia coli	b1224	[pn:respiratory nitrate reductase I alpha chain] [gn:narg]
CONTIG328	16527216_f3_17	736	6398	273	91	360	4.2(10)-33	Escherichia coli	b1227	[pn:respiratory nitrate reductase I gamma chain] [gn:nari]



CONTIG329	2863535_f1_3	737	6399	897	299	252	1.2(10)-21	Haemophilus influenzae	HI0359	[pn:gb]
CONTIG329	2214012_f1_4	738	6400	633	211	753	9.5(10)-75	Escherichia coli	b2700	[pn:hypothetical 17.6 kd protein in mlb-reca intergenic region] [gn:ygad]
CONTIG329	33595376_f1_7	739	6401	588	196	931	1.3(10)-93	Escherichia coli	b2697	[pn:alanyl-trna synthetase] [gn:alas]
CONTIG329	6034638_f2_9	740	6402	759	253	327	1.3(10)-29	Bacillus subtilis	yigB	[pn:hypothetical protein]
CONTIG329	1442277_f2_11	741	6403	516	172	746	5.2(10)-74	Escherichia coli	b2698	[pn:regulatory protein recx] [gn:oraa]
CONTIG329	16307256_f3_15	742	6404	912	304	409	2.7(10)-38	Haemophilus influenzae	HI0362	[pn:adhesin b precursor] [gn:fima]
CONTIG329	15679167_f3_16	743	6405	1095	365	1601	1.3(10)-164	Escherichia coli	b2699	[pn:reca protein] [gn:reca]
CONTIG33	36343781_f1_2	744	6406	240	80	213	2.3(10)-17	Escherichia coli	b1486	[pn:hypothetical protein]
CONTIG330	23439000_c2_40	745	6407	330	110	119	1.5(10)-7	Haemophilus influenzae	HI1250	[pn:hypothetical protein]
CONTIG330	5105068_c2_43	746	6408	1260	420	325	2.2(10)-29	Proteus vulgaris	S04739	site-specific dna-methyltransferase (cytosine-specific) (ec2.1.1.73) pvuii - proteus vulgaris
CONTIG331	4027290_c1_32	747	6409	816	272	102	0.0085	Treponema denticola	U84257	or:treponema denticola pn:methyl-accepting chemotaxis protein b gn:dmcb le:1 re:>1107 di:direct
CONTIG331	14652281_c2_34	748	6410	585	195	152	4.7(10)-11	Haemophilus influenzae	HI1418	[pn:hypothetical protein]
CONTIG331	392502_c2_38	749	6411	1881	627	1007	1.2(10)-101	Bacteriophage P4	P10277	putative p4-specific dna primase (ec 2.7.7.-)
CONTIG331	20491562_c3_40	750	6412	393	131	104	5.7(10)-6	Escherichia coli	b2624	[pn:prophage cp4-57 regulatory protein alpa] [gn:alpa]
CONTIG332	23993956_f1_1	751	6413	975	325	1387	6.2(10)-142	Escherichia coli	b0025	[pn:hypothetical 34.6 kd protein in rpst-iles intergenic region] [gn:yaac]
CONTIG332	23464537_f1_2	752	6414	2862	954	4572	0	Escherichia coli	b0026	[pn:isoleucyl-trna synthetase] [gn:iles]
CONTIG332	5209791_f2_10	753	6415	549	183	638	1.5(10)-62	Escherichia coli	b0028	[pn:probable flbb-type 16 kd peptidyl-prolyl cis- trans isomerase]
CONTIG332	5098753_f3_15	754	6416	510	170	739	2.8(10)-73	Escherichia coli	b0027	[pn:lipoprotein signal peptidase] [gn:lsps]

CONTIG332	34614462_f3_16	755	6417	987	329	1496	1.8(10)-153	Escherichia coli	b0029	[pn:lytb protein] [gn:lytb]
CONTIG333	23556562_f1_4	756	6418	384	128	397	5.0(10)-37	Escherichia coli	b4168	[pn:hypothetical 16.9 kd protein in psd-amib intergenic region] [gn:yjee]
CONTIG333	32625456_f1_5	757	6419	1368	456	1651	6.5(10)-170	Escherichia coli	b4169	[pn:n-acetylmuramoyl-l-alanine amidase precursor] [gn:amib]
CONTIG333	4307943_f1_6	758	6420	1818	606	2103	8.4(10)-218	Escherichia coli	b4170	[pn:dna mismatch repair protein mutI] [gn:mutI]
CONTIG333	33414068_f1_8	759	6421	324	108	340	5.5(10)-31	Escherichia coli	b4172	[pn:host factor-ij] [gn:hfq]
CONTIG333	21907291_f3_23	760	6422	1692	564	2008	9.8(10)-208	Escherichia coli	b4167	[pn:hypothetical 54.7 kd protein in psd-amib intergenic region] [gn:yjef]
CONTIG333	11022125_f3_24	761	6423	225	75	295	3.2(10)-26	Escherichia coli	b4168	[pn:hypothetical 16.9 kd protein in psd-amib intergenic region] [gn:yjee]
CONTIG333	25509662_f3_26	762	6424	954	318	1439	1.8(10)-147	Escherichia coli	b4171	[pn:trna delta-2-isopentenylpyrophosphate] [gn:miaa]
CONTIG333	32033567_f3_27	763	6425	372	124	586	4.7(10)-57	Escherichia coli	b4173	[pn:gtp-binding protein hflx] [gn:hflx]
CONTIG333	12928592_c3_67	764	6426	1251	417	1857	9.8(10)-192	Escherichia coli	b4166	[pn:hypothetical 43.1 kd protein in psd-amib intergenic region] [gn:yjes]
CONTIG334	35362692_f2_8	765	6427	273	91	449	1.6(10)-42	Escherichia coli	b0954	[pn:d-3-hydroxydecanoyl-acyl carrier-protein] [gn:faba]
CONTIG334	6283552_f2_16	766	6428	450	150	117	6.4(10)-7	Pseudomonas aeruginosa	JQ0133	hypothetical 26.4k protein - pseudomonas aeruginosa
CONTIG334	660278_f3_29	767	6429	315	105	113	1.6(10)-6	Haemophilus influenzae	H11343	[pn:nitrogen fixation protein] [gn:nifs]
CONTIG334	25634630_c1_39	768	6430	1257	419	1795	3.7(10)-185	Escherichia coli	b0950	[pn:pqia] [gn:pqia]
CONTIG334	2463131_c2_43	769	6431	231	77	111	1.2(10)-5	Escherichia coli	b0951	[pn:pqib] [gn:pqib]
CONTIG334	14661713_c2_44	770	6432	579	193	688	7.4(10)-68	Escherichia coli	b0952	[pn:hypothetical protein]
CONTIG334	21541091_c3_45	771	6433	2025	675	3149	0	Escherichia coli	b0948	[pn:hypothetical protein] [gn:yby]
CONTIG334	4688202_c3_46	772	6434	1908	636	2661	6.2(10)-277	Escherichia coli	b0949	[pn:hypothetical protein] [gn:yebh]
CONTIG334	34039693_c3_50	773	6435	1656	552	2259	2.5(10)-234	Escherichia coli	b0951	[pn:pqib] [gn:pqib]

CONTIG335	156328_c1_33	774	6436	399	133	91	0.0008	Escherichia coli	b2354	[pn:hypothetical protein]
CONTIG335	4100463_c1_34	775	6437	1074	358	160	3.3(10)-9	unclassified	JC4865	contractile tail sheath protein - pseudomonas aeruginosaphage psl7
CONTIG335	31427007_c2_38	776	6438	582	194	199	1.7(10)-15	unclassified	JC4865	contractile tail sheath protein - pseudomonas aeruginosaphage psl7
CONTIG335	23471062_c2_39	777	6439	525	175	146	2.0(10)-10	no gb taxonomy match	JC5192	
CONTIG335	6020968_c2_41	778	6440	1539	513	98	0.22	Archaeoglobus fulgidus	E69444	[pn:chromosome segregation protein (smcI) homolog]
CONTIG335	4040875_c3_49	779	6441	219	73	137	1.8(10)-9	Bacteriophage P2	P51772	tail protein x (gpx).
CONTIG335	34431916_c3_52	780	6442	309	103	147	4.4(10)-9	Escherichia coli	b1372	[pn:hypothetical protein]
CONTIG336	3959462_f1_2	781	6443	906	302	1083	1.0(10)-109	Escherichia coli	b4211	[pn:hypothetical 29.7 kd protein in rpli-cpnb intergenic region] [gn:ytfj]
CONTIG336	34647781_f1_3	782	6444	843	281	272	9.0(10)-24	Bacillus subtilis	ybfI	[pn:hypothetical protein]
CONTIG336	26604652_f1_4	783	6445	984	328	1215	1.1(10)-123	Escherichia coli	b4210	[pn:hypothetical 35.5 kd protein in rpli-cpnb intergenic region] [gn:ytfj]
CONTIG336	19616637_f2_14	784	6446	699	233	1036	9.8(10)-105	Escherichia coli	b4209	[pn:hypothetical protein] [gn:ytfj]
CONTIG336	34037501_c1_33	785	6447	417	139	550	3.1(10)-53	Escherichia coli	b4212	[pn:hypothetical protein] [gn:ytfj]
CONTIG336	32214687_c2_35	786	6448	1662	554	1202	2.5(10)-122	Escherichia coli	b4355	[pn:methyl-accepting chemotaxis protein i] [gn:tsr]
CONTIG337	22864163_f1_1	787	6449	720	240	1125	3.6(10)-114	Escherichia coli	b2565	[pn:dna repair protein reco] [gn:reco]
CONTIG337	12539057_f1_2	788	6450	753	251	1131	8.4(10)-115	Escherichia coli	b2564	[pn:pyridoxal phosphate biosynthetic protein pdxj] [gn:pdxj]
CONTIG337	24627067_f2_11	789	6451	435	145	699	5.0(10)-69	Escherichia coli	b2566	[pn:gtp-binding protein] [gn:era]
CONTIG337	19687915_f2_17	790	6452	936	312	815	2.6(10)-81	Escherichia coli	b2428	[pn:hypothetical protein] [gn:yfeu]
CONTIG337	3916713_f2_18	791	6453	1374	458	702	2.3(10)-69	Bacillus subtilis	ybbF	[pn:hypothetical protein]
CONTIG337	33728143_f2_19	792	6454	639	213	774	5.7(10)-77	Escherichia coli	b2560	[pn:hypothetical 21.9 kd protein in purl-dpj intergenic region] [gn:yfbb]

CONTIG337	20720005_f2_20	793	6455	564	188	687	9.4(10)-68	Escherichia coli	b2559	[pn:hypothetical 20.0 kd protein in purl-dpj intergenic region] [gn:yfhc]
CONTIG337	32120791_f3_25	794	6456	543	181	611	1.1(10)-59	Escherichia coli	b2563	[pn:dpj protein] [gn:acps]
CONTIG337	7133415_c1_36	795	6457	288	96	95	0.00058	Mycobacterium leprae	Z70722	[de:mycobacterium leprae cosmid b1770.] [pn:probable phosphoprotein phosphatase] [gn:ppp] [nt:mlb1770.13c, ppp; putative phosphoprotein]
CONTIG337	15631650_c1_39	796	6458	891	297	1094	7.0(10)-111	Escherichia coli	b2561	[pn:hypothetical protein in purl-dpj intergenic region] [gn:yfhf]
CONTIG337	34425883_c2_44	797	6459	819	273	977	1.8(10)-98	Escherichia coli	b2558	[pn:hypothetical 53.2 kd protein in purl-dpj intergenic region] [gn:yfhf]
CONTIG337	15813833_c3_61	798	6460	306	102	468	1.5(10)-44	Escherichia coli	b2562	[pn:hypothetical protein] [gn:yfhf]
CONTIG338	26301586_f2_11	799	6461	1548	516	2212	2.3(10)-229	Escherichia coli	b2011	[pn:exodeoxyribonuclease i] [gn:sbcb]
CONTIG338	30663417_c1_29	800	6462	1167	389	1330	6.9(10)-136	Escherichia coli	b2008	[pn:hypothetical 40.0 kd protein in cobu-sbcb intergenic region] [gn:yeaa]
CONTIG338	627287_c1_30	801	6463	384	128	589	2.2(10)-57	Escherichia coli	b2007	[pn:hypothetical protein]
CONTIG338	3145787_c2_31	802	6464	930	310	1209	4.5(10)-123	Escherichia coli	b2015	[pn:hypothetical protein]
CONTIG338	4120381_c2_34	803	6465	1182	394	1718	5.2(10)-177	Escherichia coli	b2010	[pn:yeec] [gn:yeec]
CONTIG338	14492842_c3_39	804	6466	1371	457	2127	2.3(10)-220	Escherichia coli	b2014	[pn:hypothetical 49.8 kd transport protein in sbcb 3'''region] [gn:yeef]
CONTIG338	29695755_c3_45	805	6467	525	175	569	3.0(10)-55	Escherichia coli	b2009	[pn:hypothetical 18.1 kd protein in phse 5'''region] [gn:sbmc]
CONTIG339	22073430_f2_23	806	6468	195	65	218	4.7(10)-18	Escherichia coli	D90862	or:escherichia coli gn:cvpa le:13907 re:14203 di:direct sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [pir accession number a04446]
CONTIG339	1445258_c1_28	807	6469	819	273	778	2.1(10)-77	Escherichia coli	b2313	[pn:colicin v production protein] [gn:cvpa]
CONTIG339	30353282_c1_29	808	6470	1551	517	2388	5.2(10)-248	Escherichia coli	b2312	[pn:amidophosphoribosyltransferase] [gn:purf]

CONTIG339	12244766_c1_32	809	6471	759	253	951	1.0(10)-95	Escherichia coli	b2308	[pn:histidine transport system permease protein hisq] [gn:hisq]
CONTIG339	36042669_c2_33	810	6472	464	154	404	9.1(10)-38	Escherichia coli	b2314	[pn:dedd protein] [gn:dedd]
CONTIG339	9766650_c2_41	811	6473	987	329	1107	2.8(10)-112	Escherichia coli	b2310	[pn:lysine-arginine-ornithine-binding periplasmic protein precursor] [gn:argt]
CONTIG339	24705382_c3_47	812	6474	216	72	141	6.7(10)-10	Escherichia coli	D90862	or:escherichia coli gn:cvpa le:13892 re:14173 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [swissprot accession number p03820];
CONTIG339	22391387_c3_49	813	6475	612	204	864	1.7(10)-86	Escherichia coli	b2311	[pn:3-octaprenyl-4-hydroxybenzoate carboxy-lyase] [gn:ubix]
CONTIG339	1174038_c3_51	814	6476	804	268	1160	7.0(10)-118	Escherichia coli	b2309	[pn:histidine-binding periplasmic protein hisj] [gn:hisj]
CONTIG339	25431631_c3_52	815	6477	717	239	1070	2.5(10)-108	Escherichia coli	b2307	[pn:histidine transport system permease protein hism] [gn:hism]
CONTIG340	34004425_f2_9	816	6478	927	309	1467	2.1(10)-150	Escherichia coli	b1344	[pn:hypothetical protein] [gn:ydao]
CONTIG340	24320165_c1_32	817	6479	1164	388	572	1.5(10)-55	Escherichia coli	b1885	[pn:methyl-accepting chemotaxis protein iv] [gn:tap]
CONTIG340	36221037_c1_33	818	6480	999	333	1494	2.8(10)-153	Escherichia coli	b1342	[pn:hypothetical protein]
CONTIG340	867076_c2_38	819	6481	1401	467	2143	4.9(10)-222	Escherichia coli	b1343	[pn:atp-dependent ma helicase dbpa] [gn:dbpa]
CONTIG340	1959386_c3_40	820	6482	522	174	250	1.8(10)-21	Escherichia coli	AF044503	[de:escherichia coli strain ec11 unknown (498), hcp gene, complete cds,and rhsg accessory genetic element vrg protein, core component anddsorf-g1 genes, complete cds.] [pn:unknown] [gn:498]
CONTIG341	5992932_f1_3	821	6483	657	219	967	2.0(10)-97	Escherichia coli	b1662	[pn:riboflavin synthase alpha chain] [gn:ribe]
CONTIG341	33854677_f1_7	822	6484	1332	444	1429	2.2(10)-146	Escherichia coli	b1659	[pn:hypothetical transcriptional regulator in cfa- purr intergenic region] [gn:ydhb]

CONTIG341	35673775_f2_17	823	6485	837	279	1176	1.3(10)-119	Escherichia coli	b1657	[pn:hypothetical protein]
CONTIG341	33630193_c1_33	824	6486	327	109	93	0.00031	Corynebacterium glutamicum	U85507	or:corynebacterium glutamicum pn:unknown gn:orf6 lc:3532 re:4290 di:complement
CONTIG341	23992051_c1_45	825	6487	1395	465	1790	1.2(10)-184	Escherichia coli	b1663	[pn:hypothetical protein in ribc 5'''region] [gn:ydhe]
CONTIG341	16834386_c2_48	826	6488	1032	344	1704	1.6(10)-175	Escherichia coli	b1658	[pn:purine nucleotide synthesis repressor] [gn:purf]
CONTIG341	12988952_c2_51	827	6489	1290	430	1494	2.8(10)-153	Escherichia coli	b1660	[pn:hypothetical 43.4 kd protein in pur-cfa intergenic region] [gn:ydhe]
CONTIG341	36120180_c3_63	828	6490	1173	391	1810	9.4(10)-187	Escherichia coli	b1661	[pn:cyclopropane-fatty-acyl- phospholipid synthase] [gn:cfa]
CONTIG342	17036458_f1_6	829	6491	1091	364	343	1.8(10)-30	Escherichia coli	P22520	colicin v secretion atp-binding protein cvab.
CONTIG342	2926577_f2_7	830	6492	624	208	241	1.7(10)-20	Escherichia coli	b0293	[pn:hypothetical 20.1 kd protein in intf-eaeh intergenic region precursor] [gn:yagz]
CONTIG342	4564068_f2_8	831	6493	687	229	439	1.8(10)-41	Escherichia coli	b0292	[pn:hypothetical 24.5 kd protein in intf-eaeh intergenic region precursor] [gn:yagy]
CONTIG342	7067568_f2_9	832	6494	2601	867	1162	4.4(10)-118	Escherichia coli	b0291	[pn:hypothetical 91.2 kd protein in intf-eaeh intergenic region precursor] [gn:yagx]
CONTIG342	35806508_f3_11	833	6495	828	276	127	3.7(10)-6	Escherichia coli	b1608	[pn:rstal] [gn:rstal]
CONTIG342	32135053_f3_15	834	6496	1740	580	138	5.0(10)-6	Escherichia coli	b0290	[pn:hypothetical 60.0 kd protein in intf-eaeh intergenic region] [gn:yagw]
CONTIG342	14879055_f3_16	835	6497	660	220	319	9.4(10)-29	Escherichia coli	b0289	[pn:hypothetical 28.2 kd protein in intf-eaeh intergenic region] [gn:ada]
CONTIG343	5182963_f1_3	836	6498	1071	357	1273	7.5(10)-130	Escherichia coli	b2213	[pn:ada regulatory protein] [gn:ada]
CONTIG343	10940625_f1_5	837	6499	1668	556	2187	1.1(10)-226	Escherichia coli	b2211	[pn:hypothetical abc transporter in eco-alkb intergenic region] [gn:yaji]
CONTIG343	7130050_f2_10	838	6500	1035	345	1454	5.0(10)-149	Escherichia coli	b2215	[pn:outer membrane protein c precursor] [gn:ompc]
CONTIG343	4119056_f2_11	839	6501	393	131	288	1.8(10)-25	Escherichia coli	b2214	[pn:hypothetical 38.5 kd protein in ada-ompc intergenic region]

CONTIG343	36572037_f2_12	840	6502	618	206	897	5.2(10)-90	Escherichia coli	b2214	[pn:hypothetical 38.5 kd protein in ada-ompC intergenic region]
CONTIG343	12679793_f3_24	841	6503	672	224	843	2.7(10)-84	Escherichia coli	b2212	[pn:alkb protein] [gn:alkb]
CONTIG343	7119075_f3_28	842	6504	1440	480	322	4.5(10)-29	Bacillus subtilis	ykoK	[pn:hypothetical protein]
CONTIG343	16491667_f3_29	843	6505	204	68	93	0.0018	Caenorhabditis elegans	Z81518	[de:caenorhabditis elegans cosmid f28d9, complete sequence.]
										[pn:f28d9.a] [nt:protein predicted using genefinder; preliminary]
CONTIG343	10400826_f3_30	844	6506	977	326	1363	2.2(10)-139	Escherichia coli	b2210	[pn:hypothetical 60.2 kd protein in eco-alkb intergenic region] [gn:yohh]
CONTIG343	23446927_c1_31	845	6507	1401	467	705	1.2(10)-69	Bacillus subtilis	ybaR	[pn:hypothetical protein]
CONTIG344	4197127_f1_12	846	6508	606	202	578	3.3(10)-56	Escherichia coli	b1304	[pn:phage shock protein a] [gn:pspa]
CONTIG344	4551958_f1_13	847	6509	255	85	342	3.3(10)-31	Escherichia coli	b1305	[pn:phage shock protein b] [gn:pspb]
CONTIG344	23463300_f3_26	848	6510	615	205	137	1.8(10)-9	Escherichia coli	b0464	[pn:potential acrab operon repressor] [gn:acrr]
CONTIG344	21601437_f3_37	849	6511	294	98	344	2.1(10)-31	Escherichia coli	b1304	[pn:phage shock protein a] [gn:pspa]
CONTIG344	14713512_f3_38	850	6512	429	143	401	1.8(10)-37	Escherichia coli	b1306	[pn:phage shock protein c] [gn:pspc]
CONTIG344	19649088_f3_39	851	6513	237	79	287	2.2(10)-25	Escherichia coli	b1307	[pn:phage shock protein d] [gn:pspd]
CONTIG344	23494193_c2_53	852	6514	996	332	1387	6.2(10)-142	Escherichia coli	b1303	[pn:hypothetical protein in pspa 5'" region] [gn:pspf]
CONTIG344	34102188_c2_55	853	6515	990	330	1302	6.4(10)-133	Escherichia coli	b1293	[pn:peptide transport system permease protein sapb] [gn:sapb]
CONTIG344	32704131_c2_58	854	6516	1041	347	1550	3.2(10)-159	Escherichia coli	b1291	[pn:peptide transport system atp-binding protein sapd] [gn:sapd]
CONTIG344	10392093_c3_68	855	6517	1761	587	2401	2.2(10)-249	Escherichia coli	b1294	[pn:peptide transport periplasmic protein sapa precursor] [gn:sapa]
CONTIG344	24881542_c3_69	856	6518	906	302	1169	7.9(10)-119	Escherichia coli	b1292	[pn:peptide transport system permease protein sapc] [gn:sapc]
CONTIG344	26448807_c3_70	857	6519	825	275	1289	1.5(10)-131	Escherichia coli	b1290	[pn:peptide transport system atp-binding protein sapf] [gn:sapf]

CONTIG345	33617808_f2_11	858	6520	435	145	329	8.1(10)-30	Escherichia coli	b1856	[pn:hypothetical 46.7 kd protein in msbb-ruvb intergenic region] [gn:yeba]
CONTIG345	16303831_c2_48	859	6521	1341	447	1384	1.3(10)-141	Escherichia coli	b0574	[pn:hypothetical protein] [gn:ylcd]
CONTIG345	24629011_c2_53	860	6522	2496	832	1315	2.7(10)-134	Bacillus subtilis	yvgX	[pn:hypothetical protein]
CONTIG345	1219468_c3_55	861	6523	573	191	584	7.7(10)-57	Escherichia coli	b0572	[pn:hypothetical protein] [gn:ylcb]
CONTIG345	4688752_c3_56	862	6524	381	127	253	9.1(10)-22	Escherichia coli	b0573	[pn:hypothetical protein] [gn:ylcc]
CONTIG345	21520677_c3_58	863	6525	3177	1059	4318	0	Escherichia coli	b0575	[pn:hypothetical protein in phev 5''' region] [gn:ybde]
CONTIG346	878876_f1_1	864	6526	402	134	518	7.7(10)-50	Escherichia coli	b2582	[pn:hypothetical protein in the ung 3''' region] [gn:yfig]
CONTIG346	32617793_f1_3	865	6527	2715	905	3772	0	Escherichia coli	b2584	[pn:hypothetical protein] [gn:yfiq]
CONTIG346	29782832_f2_8	866	6528	783	261	999	8.1(10)-101	Escherichia coli	b2583	[pn:hypothetical protein] [gn:yfip]
CONTIG346	24807932_f2_14	867	6529	1377	459	1923	1.0(10)-198	Escherichia coli	b2585	[pn:cdp-diacylglycerol-serine o-phosphatidyltransferase] [gn:pssa]
CONTIG346	19972900_f3_26	868	6530	423	141	417	3.8(10)-39	Escherichia coli	b2586	[pn:hypothetical 9.9 kd protein in pss-kgtp intergenic region] [gn:yfim]
CONTIG346	50066_c3_52	869	6531	1377	459	1610	1.5(10)-165	Escherichia coli	b2587	[pn:alpha-ketoglutarate permease] [gn:kgtp]
CONTIG347	6658593_f1_1	870	6532	441	147	479	1.0(10)-45	Escherichia coli	AJ224995	[de:escherichia coli ecorii restriction endonuclease gene.] [pn:restriction endonuclease] [gn:ecorii]
CONTIG347	10605292_f1_5	871	6533	1557	519	124	3.2(10)-8	Bacillus subtilis	yefF	[pn:hypothetical protein]
CONTIG347	20722933_f1_6	872	6534	1644	548	93	1.0(10)0	Caenorhabditis elegans	U33058	or:caenorhabditis elegans pn:unc-89 gn:unc-89 le:join(4920 re:4969,5656 di:direct nt:giant ig superfamily member located in the middle
CONTIG347	2133550_f3_21	873	6535	1404	468	1121	9.6(10)-114	Pseudomonas alcaligenes	U77945	[de:pseudomonas alcaligenes maturase-related protein gene, completecds.] [pn:maturase-related protein]



CONTIG347	24729507_c2_50	874	6536	204	68	116	1.5(10)-6	Anabaena PCC7120	AF047044	[de:anabaena pcc7120 insertion sequence is1594 putative transposase, complete cds.] [pn:putative transposase] [nt:tnp1594]
CONTIG347	12319757_c2_51	875	6537	1596	532	1528	7.2(10)-157	Escherichia coli	b1961	[pn:dna-cytosine methyltransferase] [gn:dcn]
CONTIG347	35597962_c3_58	876	6538	306	102	176	3.2(10)-13	Pseudomonas atlantica	A32816	hypothetical protein, 33k (insertion sequence is492) -pseudomonas atlantica
CONTIG348	25422808_fl_1	877	6539	1383	461	2037	8.3(10)-211	Escherichia coli	b0211	[pn:regulatory protein dnir and hypothetical yafg] [gn:mld]
CONTIG348	3393955_fl_6	878	6540	306	102	108	6.7(10)-6	Corynebacterium glutamicum	U85507	or:corynebacterium glutamicum pn:unknown gn:orf6 le:3532 re:4290 di:complement
CONTIG348	26687512_fl_7	879	6541	927	309	1370	4.0(10)-140	Escherichia coli	b0208	[pn:hypothetical transcriptional regulator in rrh- dnir intergenic region] [gn:yafc]
CONTIG348	16047027_f2_12	880	6542	192	64	212	2.0(10)-17	Escherichia coli	b0212	[pn:hypothetical protein] [gn:glob]
CONTIG348	24884441_c1_24	881	6543	819	273	1204	1.5(10)-122	Escherichia coli	b0207	[pn:hypothetical 29.4 kd protein in aspu-dnir intergenic region]
CONTIG348	12148381_c1_29	882	6544	873	291	1288	1.8(10)-131	Escherichia coli	b0209	[pn:hypothetical protein in aspu-dnir intergenic region] [gn:yafd]
CONTIG348	6926692_c2_34	883	6545	1191	397	1054	1.2(10)-106	Bacillus subtilis	ytdD	[pn:hypothetical protein]
CONTIG348	24098587_c3_41	884	6546	786	262	771	1.2(10)-76	Escherichia coli	b0210	[pn:hypothetical 23.0 kd protein in aspu-dnir intergenic region]
CONTIG349	31489806_fl_10	885	6547	1011	337	782	8.0(10)-78	Escherichia coli	b1899	[pn:hypothetical protein]
CONTIG349	2907256_fl_11	886	6548	914	305	702	2.3(10)-69	Escherichia coli	b1897	[pn:trehalose phosphatase] [gn:otsb]
CONTIG349	35397937_f2_12	887	6549	420	140	307	1.7(10)-27	Escherichia coli	P52092	hypothetical 9.1 kd protein in araf-fn intergenic region.
CONTIG349	4886041_f2_15	888	6550	609	203	194	1.6(10)-15	Escherichia coli	b0424	[pn:thij protein] [gn:thij]
CONTIG349	32301093_f2_17	889	6551	1554	518	2320	8.5(10)-241	Escherichia coli	b1900	[pn:l-arabinose transport atp-binding protein arag] [gn:arag]

CONTIG349	22125378_f3_22	890	6552	1041	347	1573	1.2(10)-161	Escherichia coli	b1901	[pn:l-arabinose-binding periplasmic protein precursor] [gn:araf]
CONTIG349	34069716_c1_39	891	6553	540	180	539	4.5(10)-52	Escherichia coli	b1902	[pn:ferritin-like protein] [gn:yecj]
CONTIG349	4574191_c1_40	892	6554	1356	452	1123	5.9(10)-114	Escherichia coli	b4123	[pn:anaerobic c4-dicarboxylate transporter dcub] [gn:dcub]
CONTIG349	24880458_c2_56	893	6555	1266	422	713	1.7(10)-70	Escherichia coli	b3754	[pn:hypothetical 51.5 kd protein in rbsr-rssc intergenic region] [gn:yieo]
CONTIG349	32441526_c3_70	894	6556	240	80	171	4.5(10)-13	Escherichia coli	b1903	[pn:hypothetical protein]
CONTIG350	31913255_f2_14	895	6557	636	212	946	3.3(10)-95	Escherichia coli	b2186	[pn:hypothetical 37.8 kd protein in rply-prol intergenic region] [gn:yejk]
CONTIG350	4298562_f2_20	896	6558	717	239	1101	1.3(10)-111	Escherichia coli	b2183	[pn:hypothetical 25.9 kd protein in bcr-rply intergenic region] [gn:yejd]
CONTIG350	24823252_f2_21	897	6559	1215	405	1519	6.5(10)-156	Escherichia coli	b2182	[pn:bicyclomycin resistance protein] [gn:bcr]
CONTIG350	15632767_f2_22	898	6560	372	124	460	1.1(10)-43	Escherichia coli	b2181	[pn:hypothetical 12.5 kd protein in bcr 5''' region] [gn:yejl]
CONTIG350	26602187_c1_45	899	6561	399	133	94	0.00339	Listeria innocua	Q01836	protein p60 precursor (invasion-associated protein).
CONTIG350	26439076_c1_47	900	6562	1776	592	2642	6.4(10)-275	Escherichia coli	b2184	[pn:hypothetical 66.4 kd protein in rsua-rply intergenic region] [gn:yejh]
CONTIG350	5103968_c2_51	901	6563	1035	345	1250	2.1(10)-127	Escherichia coli	b2179	[pn:hypothetical 38.1 kd protein in bcr 5''' region] [gn:yeje]
CONTIG350	682093_c3_56	902	6564	906	302	1310	9.0(10)-134	Escherichia coli	b2178	[pn:hypothetical 40.4 kd protein in bcr 5''' region] [gn:yejb]
CONTIG350	16803125_c3_58	903	6565	1644	548	1921	1.6(10)-198	Escherichia coli	b2180	[pn:hypothetical abc transporter in bcr 5''' region] [gn:yejf]
CONTIG350	25885842_c3_64	904	6566	480	160	117	2.3(10)-7	Escherichia coli	S24805	hypothetical protein fwdl 566 - escherichia coli
CONTIG350	14504807_c3_65	905	6567	363	121	172	3.5(10)-13	Escherichia coli	P28247	very hypothetical 19.2 kd protein in bcr 3' region.
CONTIG350	20500018_c3_68	906	6568	309	103	343	2.7(10)-31	Escherichia coli	b2185	[pn:50s ribosomal protein l25] [gn:rply]
CONTIG351	4015938_f1_1	907	6569	1488	496	1348	8.5(10)-138	Bacillus subtilis	ybaR	[pn:hypothetical protein]

CONTIG351	36219711_f1_2	908	6570	846	282	1339	7.7(10)-137	Yersinia enterocolitica	Y13308	[PN:hypothetical protein] [DE:Yersinia enterocolitica plasmid DNA fragment, strain 15673.] [NT:ORF2] [LE:3722] [RE:4588] [DI:complement]
CONTIG351	30275251_f2_16	909	6571	726	242	1028	6.9(10)-104	Yersinia enterocolitica	U58366	or:yersinia enterocolitica pn:arsh gn:arsh le:3823 re:4521 di:direct nt:required for arsenic resistance
CONTIG351	9947125_f3_29	910	6572	1536	512	536	9.5(10)-52	Escherichia coli	b3558	[pn:insertion element is150 hypothetical 33.3 kd protein] [gn:y15b]
CONTIG351	10979628_c1_32	911	6573	1443	481	942	9.0(10)-95	Rhizobium sp.	P55373	putative transposase y4bf,
CONTIG351	1282876_c2_43	912	6574	465	155	105	4.5(10)-6	Escherichia coli	P19770	insertion element is150 hypothetical 14 kd protein (orf).
CONTIG351	22350087_c2_48	913	6575	375	125	385	9.5(10)-36	Escherichia coli	b3501	[pn:arsenical resistance operon arsefg repressor] [gn:arsr]
CONTIG351	1964202_c2_49	914	6576	1302	434	1649	1.1(10)-169	Escherichia coli	b3502	[pn:arsenical pump membrane protein] [gn:arsb]
CONTIG351	4110263_c2_50	915	6577	435	145	659	8.8(10)-65	Escherichia coli	b3503	[pn:arsenate reductase] [gn:arsc]
CONTIG352	30475831_f1_4	916	6578	624	208	90	0.51	Rattus norvegicus	S54307	myosin heavy chain - rat
CONTIG352	5156338_f1_6	917	6579	486	162	101	9.4(10)-5	Mycobacterium tuberculosis	Z95586	unknown,,mtey336.26,mtey336.26. len
CONTIG352	35329530_f2_11	918	6580	1326	442	532	2.5(10)-51	Bacteriophage HK97	P49859	portal protein (gp3).
CONTIG352	33694458_f3_14	919	6581	408	136	153	3.6(10)-11	Bacteriophage phi-105	L35561	or:bacteriophage phi-105 pn:holin le:796 re:1170 di:direct sr:bacteriophage phi-105 dna nt:orf2; potential dual start motif; putative
CONTIG352	31922905_f3_15	920	6582	1770	590	611	1.1(10)-59	Escherichia coli	b1149	[pn:hypothetical protein]
CONTIG352	3230287_f3_17	921	6583	876	292	108	0.00035	Haemophilus influenzae	HI0714	[pn:atp-dependent clp protease proteolytic component] [gn:clpp]
CONTIG353	5352308_f1_3	922	6584	1251	417	1489	9.6(10)-153	Haemophilus influenzae	HI0166	[pn:nitrogen fixation protein] [gn:mfe]
CONTIG353	4900311_f1_5	923	6585	1227	409	1512	3.6(10)-155	Haemophilus influenzae	HI0171	[pn:phenolhydroxylase component]

CONTIG353	23600408_f1_11	924	6586	489	163	608	2.2(10)-59	Haemophilus influenzae	HI0674	[pn:xanthine guanine phosphoribosyl transferase gpt] [gn:hi0692]
CONTIG353	31257276_f2_12	925	6587	1449	483	1348	8.5(10)-138	Haemophilus influenzae	HI0164	[pn:nadh]
CONTIG353	886265_f2_14	926	6588	879	293	617	2.5(10)-60	Haemophilus influenzae	HI0167	[pn:hypothetical protein]
CONTIG353	4015843_f2_19	927	6589	1068	356	1529	5.5(10)-157	Escherichia coli	b0231	[pn:hypothetical protein dinp] [gn:dinp]
CONTIG353	25525302_f3_25	928	6590	642	214	586	4.7(10)-57	Haemophilus influenzae	HI0168	[pn:nadh]
CONTIG353	16835331_f3_26	929	6591	603	201	672	3.7(10)-66	Haemophilus influenzae	HIN_166	[pn:nadh] [gn:hi0170]
CONTIG353	3402000_f3_27	930	6592	216	72	102	9.3(10)-6	Haemophilus influenzae	HI0173	[pn:hypothetical protein]
CONTIG353	897881_c2_41	931	6593	558	186	857	9.0(10)-86	Escherichia coli	b0237	[pn:aminoacyl-histidine dipeptidase precursor] [gn:pepd]
CONTIG353	5283561_c2_42	932	6594	942	314	1379	4.4(10)-141	Escherichia coli	b0237	[pn:aminoacyl-histidine dipeptidase precursor] [gn:pepd]
CONTIG354	32426885_f1_1	933	6595	1701	567	2515	1.8(10)-261	Escherichia coli	b0779	[pn:excision nuclease abc subunit b] [gn:uvrb]
CONTIG354	5113562_f1_5	934	6596	252	84	310	8.4(10)-28	Escherichia coli	b0784	[pn:molybdopterin converting factor, subunit 1] [gn:moad]
CONTIG354	26442752_f1_11	935	6597	444	148	374	1.3(10)-34	Escherichia coli	b0791	[pn:hypothetical protein]
CONTIG354	29941531_f2_18	936	6598	513	171	694	1.7(10)-68	Escherichia coli	b0785	[pn:molybdopterin converting factor, subunit 2] [gn:moae]
CONTIG354	5192562_f3_31	937	6599	1212	404	1473	4.7(10)-151	Escherichia coli	b0781	[pn:molybdenum cofactor biosynthesis protein a] [gn:moaa]
CONTIG354	33208555_f3_32	938	6600	537	179	816	2.0(10)-81	Escherichia coli	b0782	[pn:molybdenum cofactor biosynthesis protein b] [gn:moab]
CONTIG354	22353383_f3_33	939	6601	489	163	768	2.5(10)-76	Escherichia coli	b0783	[pn:molybdenum cofactor biosynthesis protein c] [gn:moac]
CONTIG354	31853402_f3_35	940	6602	711	237	998	1.0(10)-100	Escherichia coli	b0786	[pn:hypothetical protein]
CONTIG354	36582349_c1_45	941	6603	1297	432	1608	2.3(10)-165	Escherichia coli	b0792	[pn:hypothetical protein]
CONTIG354	15907653_c1_48	942	6604	1155	385	1314	3.3(10)-134	Escherichia coli	b0788	[pn:hypothetical protein]

CONTIG354	4194068_c1_54	943	6605	918	306	974	3.7(10)-98	Escherichia coli	b0780	[pn:hypothetical protein]
CONTIG354	24878930_c2_61	944	6606	306	102	454	4.5(10)-43	Escherichia coli	b0790	[pn:hypothetical protein]
CONTIG354	32048331_c2_62	945	6607	1344	448	1686	1.3(10)-173	Escherichia coli	b0789	[pn:hypothetical protein]
CONTIG354	34018807_c3_74	946	6608	717	239	721	2.3(10)-71	Escherichia coli	b0790	[pn:hypothetical protein]
CONTIG355	6350712_fl_4	947	6609	717	239	1055	9.5(10)-107	Escherichia coli	b3651	[pn:spou protein] [gn:spou]
CONTIG355	36134652_fl_5	948	6610	2085	695	3169	0	Escherichia coli	b3652	[pn:dna recombinase] [gn:recg]
CONTIG355	644762_fl_8	949	6611	1425	475	1955	4.0(10)-202	Escherichia coli	b3654	[pn:hypothetical 48.9 kd protein in glts 3''' region] [gn:yice]
CONTIG355	21488907_f2_14	950	6612	330	110	347	1.0(10)-31	Escherichia coli	b3649	[pn:dna-directed ma polymerase omega chain] [gn:rp0z]
CONTIG355	6016411_f2_24	951	6613	1728	576	2085	6.7(10)-216	Escherichia coli	b3655	[pn:hypothetical 62.3 kd protein in glts-selc intergenic region] [gn:yich]
CONTIG355	11223782_f3_26	952	6614	2130	710	3369	0	Escherichia coli	b3650	[pn:diphosphate 3'''-pyrophosphohydrolase] [gn:spot]
CONTIG355	104836_c1_42	953	6615	1215	405	1441	1.2(10)-147	Escherichia coli	b3653	[pn:sodium/glutamate symport carrier protein] [gn:glts]
CONTIG355	24035758_c2_62	954	6616	585	195	632	6.4(10)-62	Escherichia coli	A30374	hypothetical 77k protein (spot 3' region) - escherichia coli
CONTIG356	23962828_fl_8	955	6617	255	85	398	4.0(10)-37	Escherichia coli	b3610	[pn:glutaredoxin 3] [gn:grxc]
CONTIG356	21532290_fl_9	956	6618	1086	362	1613	7.0(10)-166	Escherichia coli	b3608	[pn:l-glycerol 3-phosphate dehydrogenase] [gn:gpsa]
CONTIG356	13066577_fl_10	957	6619	876	292	1374	1.5(10)-140	Escherichia coli	b3607	[pn:serine acetyltransferase] [gn:cysc]
CONTIG356	31515805_f2_15	958	6620	459	153	626	2.7(10)-61	Escherichia coli	b3611	[pn:hypothetical 15.6 kd protein in secb-tdh intergenic region] [gn:yibn]
CONTIG356	602133_f2_16	959	6621	504	168	736	6.0(10)-73	Escherichia coli	b3609	[pn:protein-export protein secb] [gn:secb]
CONTIG356	31773430_c1_32	960	6622	519	173	345	1.6(10)-31	Escherichia coli	P20343	very hypothetical cysx protein.
CONTIG356	24348137_c2_46	961	6623	1587	529	2409	3.2(10)-250	Escherichia coli	b3612	[pn:putative 2,3-bisphosphoglycerate independent phosphoglycerate] [gn:yibo]

CONTIG356	4393955_c2_47	962	6624	1293	431	1601	1.3(10)-164	Escherichia coli	b3613	[pn:hypothetical 47.5 kd protein in secb-tdh intergenic region] [gn:yibp]
CONTIG356	21509378_c2_48	963	6625	966	322	1034	1.6(10)-104	Escherichia coli	b3614	[pn:hypothetical 30.7 kd protein in secb-tdh intergenic region] [gn:yibq]
CONTIG356	25562927_c3_51	964	6626	417	139	523	2.2(10)-50	Escherichia coli	P15041	very hypothetical 17.7 kd protein in secb region.
CONTIG357	33777166_fl_3	965	6627	606	202	717	6.2(10)-71	Escherichia coli	b2136	[pn:hypothetical 21.4 kd protein in pbpg-cdd intergenic region] [gn:yohd]
CONTIG357	25595075_fl_10	966	6628	432	144	404	9.1(10)-38	Escherichia coli	b2141	[pn:hypothetical 14.6 kd protein in pbpg-cdd intergenic region]
CONTIG357	22048902_f2_25	967	6629	360	120	126	6.0(10)-8	Escherichia coli	U00007	or:escherichia coli pn:yohk le:43789 re:44535 di:direct sr:escherichia coli k12 bbb2600
CONTIG357	4348418_f2_26	968	6630	894	298	1234	1.0(10)-125	Escherichia coli	b2143	[pn:cytidine deaminase] [gn:cdd]
CONTIG357	36580143_f3_41	969	6631	720	240	784	5.0(10)-78	Escherichia coli	b2142	[pn:hypothetical 24.5 kd protein in pbpg-cdd intergenic region] [gn:yohk]
CONTIG357	20989806_f3_42	970	6632	765	255	1098	2.6(10)-111	Escherichia coli	b2144	[pn:sana protein] [gn:sana]
CONTIG357	13804192_c1_50	971	6633	978	326	1378	5.5(10)-141	Escherichia coli	b2140	[pn:hypothetical 35.2 kd protein in pbpg-cdd intergenic region]
CONTIG357	33625277_c1_56	972	6634	537	179	399	3.1(10)-37	Escherichia coli	b2134	[pn:penicillin-binding protein 7 precursor] [gn:pbpg]
CONTIG357	9960917_c2_65	973	6635	930	310	1038	6.0(10)-105	Escherichia coli	b2137	[pn:hypothetical oxidoreductase in pbpg-cdd intergenic region]
CONTIG357	29820255_c2_67	974	6636	645	215	899	3.2(10)-90	Escherichia coli	b2135	[pn:hypothetical 22.4 kd protein in pbpg-cdd intergenic region] [gn:yohc]
CONTIG357	34547640_c2_68	975	6637	267	89	349	6.2(10)-32	Escherichia coli	b2134	[pn:penicillin-binding protein 7 precursor] [gn:pbpg]
CONTIG357	6532628_c3_76	976	6638	1431	477	1491	6.0(10)-153	Escherichia coli	b2138	[pn:hypothetical 43.3 kd protein in pbpg-cdd intergenic region] [gn:yohg]
CONTIG358	35257828_fl_2	977	6639	2559	853	3835	0	Escherichia coli	b1049	[pn:periplasmic glucans biosynthesis protein mdoh] [gn:mdoh]
CONTIG358	4506967_fl_3	978	6640	237	79	333	3.1(10)-30	Escherichia coli	b1050	[pn:hypothetical protein in mdoh-msyb intergenic region] [gn:ycek]

CONTIG358	30476516_f2_17	979	6641	1137	379	1610	1.5(10)-165	Escherichia coli	b1055	[pn:hypothetical 40.0 kd protein in hrb 5'''region] [gn:ycea]
CONTIG358	23472503_f3_22	980	6642	1614	538	2500	7.2(10)-260	Escherichia coli	b1048	[pn:periplasmic glucans biosynthesis protein mdog precursor] [gn:mdog]
CONTIG358	632292_c1_35	981	6643	621	207	788	1.8(10)-78	Escherichia coli	b1056	[pn:hypothetical 18.7 kd protein in hrb 5'''region] [gn:ycei]
CONTIG358	15112506_c2_56	982	6644	957	319	1353	2.5(10)-138	Escherichia coli	b1054	[pn:membrane protein affecting cell division, growth and high temperature survival] [gn:htrb]
CONTIG358	19688586_c2_57	983	6645	1290	430	1516	1.3(10)-155	Escherichia coli	b1053	[pn:hypothetical 43.9 kd protein in msyb-htrb intergenic region] [gn:ycee]
CONTIG358	11132918_c2_65	984	6646	471	157	714	1.3(10)-70	Escherichia coli	b1047	[pn:hypothetical protein]
CONTIG358	32632692_c3_66	985	6647	309	103	392	1.7(10)-36	Escherichia coli	b1059	[pn:hypothetical protein] [gn:sola]
CONTIG358	788387_c3_67	986	6648	651	217	655	2.2(10)-64	Escherichia coli	b1057	[pn:hypothetical protein]
CONTIG358	1985430_c3_73	987	6649	423	141	608	2.2(10)-59	Escherichia coli	b1051	[pn:acidic protein msyb, multicopy suppressor of secy] [gn:msyb]
CONTIG358	11892042_c3_74	988	6650	273	91	90	0.00063	Mycobacterium tuberculosis	Q10700	hypothetical 26.0 kd protein cy49.31c.
CONTIG359	23704667_f1_3	989	6651	1299	433	1955	4.0(10)-202	Escherichia coli	b3780	[pn:rhb] [gn:rhb]
CONTIG359	3157813_f1_4	990	6652	1491	497	1952	8.4(10)-202	Escherichia coli	b3779	[pn:guanosine pentaphosphatase] [gn:gppa]
CONTIG359	36110330_f1_8	991	6653	277	93	344	2.1(10)-31	Escherichia coli	b3775	[pn:peptidyl-prolyl cis-trans isomerase c] [gn:ppic]
CONTIG359	15812950_c1_32	992	6654	405	135	595	5.2(10)-58	Escherichia coli	b3781	[pn:thioredoxin] [gn:trxa]
CONTIG359	32453180_c2_37	993	6655	2031	677	3208	0	Escherichia coli	b3778	[pn:atp-dependent dna helicase rep] [gn:rep]
CONTIG359	4382713_c2_41	994	6656	498	166	643	4.2(10)-63	Escherichia coli	b3784	[pn:putative undecaprenyl-phosphate alpha-n-acetylglucosaminyltransferase] [gn:rfc]
CONTIG359	26599182_c3_44	995	6657	1338	446	2038	6.5(10)-211	Escherichia coli	b3783	[pn:transcription termination factor] [gn:rho]

CONTIG36	7089053_c3_5	996	6658	525	175	891	2.2(10)-89	Escherichia coli	b2580	[pn:uracil-dna glycosylase] [gn:ung]
CONTIG360	6522187_f1_2	997	6659	885	295	949	1.6(10)-95	Klebsiella pneumoniae	AF040380	[dc:klebsiella pneumoniae ribosomal protein l11 methyltransferase(prma) gene, partial cds; carbonic anhydrase (cat) and yhdg homologues, complete cds; and small dna binding protein f <sup>11</sup> ] [pn:carbonic anhydrase] [gn:cat]
CONTIG360	4741568_f1_9	998	6660	1149	383	1396	7.0(10)-143	Escherichia coli	b3265	[pn:acrc] [gn:acrc]
CONTIG360	6292163_f2_10	999	6661	1275	425	2007	1.3(10)-207	Escherichia coli	b3256	[pn:biotin carboxylase] [gn:accc]
CONTIG360	30163283_f2_11	1000	6662	291	97	200	3.7(10)-16	Escherichia coli	b3257	[pn:hypothetical 9.1 kd protein in acce-panf intergenic region] [gn:yhdt]
CONTIG360	34569061_f2_14	1001	6663	900	300	1313	4.4(10)-134	Escherichia coli	b3259	[pn:ribosomal protein l11 methyltransferase] [gn:prma]
CONTIG360	33797942_f2_17	1002	6664	312	104	401	1.8(10)-37	Haemophilus influenzae	HI0980	[pn:factor-for-inversion stimulation protein] [gn:fis]
CONTIG360	2908507_f2_19	1003	6665	2253	751	809	1.3(10)-94	Bacillus subtilis	ykoW	[pn:hypothetical protein]
CONTIG360	3991262_f3_22	1004	6666	1488	496	1914	9.0(10)-198	Escherichia coli	b3258	[pn:sodium/pantothenate symporter] [gn:panf]
CONTIG360	4556526_f3_25	1005	6667	969	323	1585	6.5(10)-163	Escherichia coli	b3260	[pn:hypothetical 35.9 kd protein in pmra-fis intergenic region] [gn:yhdg]
CONTIG360	26036410_f3_31	1006	6668	233	78	145	6.5(10)-9	Escherichia coli	b3266	[pn:acrf] [gn:acrf]
CONTIG360	16834836_c3_58	1007	6669	654	218	555	9.1(10)-54	Escherichia coli	b3264	[pn:potential acrf/envcd operon repressor] [gn:envr]
CONTIG361	16853457_f1_1	1008	6670	918	306	1425	5.9(10)-146	Escherichia coli	b0750	[pn:quinolate synthetase a protein] [gn:nada]
CONTIG361	14457650_f2_13	1009	6671	789	263	881	2.6(10)-88	Escherichia coli	b0751	[pn:pnuc protein] [gn:pnuc]
CONTIG361	33876317_f2_17	1010	6672	1077	359	1703	2.1(10)-175	Escherichia coli	b0754	[pn:phospho-2-dehydro-3-deoxyheptanate aldolase]
CONTIG361	34241057_c1_38	1011	6673	1113	371	1434	6.5(10)-147	Escherichia coli	b0756	[pn:aldose 1-epimerase] [gn:galm]



CONTIG361	11995900_c1_42	1012	6674	447	149	149	9.6(10)-11	Escherichia coli	b0753	[pn:hypothetical protein] [gn:ybgs]
CONTIG361	25980277_c1_43	1013	6675	954	318	930	1.7(10)-93	Escherichia coli	b0752	[pn:hypothetical protein] [gn:ybgr]
CONTIG361	877066_c2_47	1014	6676	1182	394	1686	1.3(10)-173	Escherichia coli	b0757	[pn:galactokinase] [gn:galk]
CONTIG361	23572188_c2_50	1015	6677	756	252	1120	1.2(10)-113	Escherichia coli	b0755	[pn:phosphoglycerate mutase 1] [gn:gpmA]
CONTIG361	36541291_c3_57	1016	6678	1218	406	1734	1.1(10)-178	Escherichia coli	b0759	[pn:udp-glucose 4-epimerase] [gn:gale]
CONTIG361	4343818_c3_58	1017	6679	1056	352	1772	1.0(10)-182	Escherichia coli	b0758	[pn:galactose-1-phosphate uridylyltransferase] [gn:galt]
CONTIG362	20177211_fl_1	1018	6680	696	232	798	1.6(10)-79	Escherichia coli	b0489	[pn:hypothetical protein] [gn:ybbk]
CONTIG362	1054782_fl_5	1019	6681	858	286	824	2.8(10)-82	Escherichia coli	b0482	[pn:hypothetical protein] [gn:ybap]
CONTIG362	12913181_fl_6	1020	6682	579	193	666	1.6(10)-65	Escherichia coli	b0481	[pn:hypothetical protein] [gn:ybak]
CONTIG362	2283_fl_13	1021	6683	2640	880	3509	0	Escherichia coli	b0484	[pn:hypothetical protein] [gn:ybar]
CONTIG362	25431562_f3_20	1022	6684	456	152	449	1.6(10)-42	Escherichia coli	b0488	[pn:hypothetical protein] [gn:ybbj]
CONTIG362	6456561_f3_27	1023	6685	1026	342	1248	3.3(10)-127	Escherichia coli	b0479	[pn:fosmidomycin resistance protein] [gn:fsr]
CONTIG362	4735278_c1_29	1024	6686	1698	566	2631	9.4(10)-274	Escherichia coli	b0480	[pn:udp-sugar hydrolase precursor] [gn:ushA]
CONTIG362	21659407_c3_62	1025	6687	444	148	603	7.5(10)-59	Escherichia coli	b0487	[pn:hypothetical protein] [gn:ybbi]
CONTIG362	672192_c3_66	1026	6688	207	69	113	6.7(10)-6	Escherichia coli	b0544	[pn:hypothetical protein] [gn:ybck]
CONTIG363	21666540_fl_5	1027	6689	735	245	1128	1.8(10)-114	Transposon Tn1525	M12900	or:transposon tn1525 gn:p12 le:996 re:>1721 di:direct sr:transposon tn1525 dna nt:putative
CONTIG363	4085387_f3_19	1028	6690	228	76	96	0.00018	Thiobacillus ferrooxidans	AF032884	[de:thiobacillus ferrooxidans n-acetylglucosamine-1-phosphateuridylyltransferase (glmU) gene, partial cds; glucosamine synthase(glmS)] [pn:transposition complex] [gn:tnsa]

CONTIG363	2928432_f3_25	1029	6691	861	287	1457	2.3(10)-149	Cloning vector pKF296	D63840	or:cloning vector pkf296 pn:aminoglucoside phosphotransferase in supe host gn:apg3 le:322 re:1137 di:complement sr:cloning vector pkf296 dna nt:cag for gln at the
CONTIG363	21666540_f3_26	1030	6692	786	262	1128	1.8(10)-114	Transposon Tn1525	M12900	or:transposon tn1525 gn:p12 le:996 re:>1721 di:direct sr:transposon tn1525 dna nt:putative
CONTIG363	6439528_f3_31	1031	6693	255	85	345	1.6(10)-31	Escherichia coli	I77547	hypothetical protein 2 (insertion sequence is903) - escherichia coli
CONTIG363	4772550_c1_32	1032	6694	1665	555	1066	6.5(10)-108	Escherichia coli	b4114	[pn:hypothetical 61.7 kd protein in bass-adyi intergenic region]
CONTIG363	11891882_c1_35	1033	6695	738	246	1268	2.6(10)-129	Salmonella ordonez	S34451	hypothetical protein (insertion sequence is261) - salmonellaordonez plasmid pip173
CONTIG363	25558159_c2_40	1034	6696	821	273	1387	6.2(10)-142	Escherichia coli	X02527	or:escherichia coli le:199 re:1122 di:direct nt:orf1 (aa 1-307)
CONTIG363	11891882_c2_45	1035	6697	738	246	1268	2.6(10)-129	Salmonella ordonez	S34451	hypothetical protein (insertion sequence is261) - salmonellaordonez plasmid pip173
CONTIG363	2397500_c2_47	1036	6698	999	333	1492	4.7(10)-153	Yersinia pestis	AF053946	[de:yersinia pestis plasmid pcd1, complete plasmid sequence.] [pn:transposase (tn1000) homolog] [gn:tnpa]
CONTIG363	12553761_c3_49	1037	6699	2148	716	2685	1.8(10)-279	Escherichia coli	I56963	transposase (transposons) - escherichia coli
CONTIG364	4979758_f1_7	1038	6700	591	197	571	1.8(10)-55	Escherichia coli	b0622	[pn:hypothetical protein in cspe 5'''region] [gn:ybeg]
CONTIG364	25882937_f1_12	1039	6701	2715	905	1008	9.0(10)-102	Methanobacter ium thermoautotro phicum	MTH1516	[pn:cation-transporting p-atpase pac]
CONTIG364	2350761_f2_26	1040	6702	540	180	370	3.7(10)-34	Escherichia coli	b0607	[pn:hypothetical protein] [gn:ybdq]
CONTIG364	14260803_f3_46	1041	6703	713	238	928	2.7(10)-93	Escherichia coli	b1990	[pn:31.6 kd protein in cobt 3'''region precursor] [gn:erfk]
CONTIG364	19689037_c1_55	1042	6704	1110	370	186	7.7(10)-12	Escherichia coli	b2074	[pn:hypothetical protein]

CONTIG364	16103578_c2_70	1043	6705	477	159	169	7.4(10)-13	Saccharomyces cerevisiae	X85757	or:saccharomyces cerevisiae pn:unknown gn:internal orf g 1669 le:6964 re:7365 di:direct sr:baker's yeast
CONTIG364	33632800_c2_77	1044	6706	1344	448	1931	1.3(10)-199	Escherichia coli	b1981	[pn:hypothetical protein] [gn:shia]
CONTIG364	4027332_c3_85	1045	6707	3075	1025	633	2.5(10)-59	Haemophilus influenzae	H10895	[pn:acriflavine resistance protein] [gn:acrb]
CONTIG364	32523467_c3_88	1046	6708	1551	517	2201	3.5(10)-228	Escherichia coli	b1982	[pn:amp nucleosidase] [gn:amn]
CONTIG365	24237891_fl_12	1047	6709	538	180	509	6.9(10)-49	Escherichia coli	b3939	[pn:cystathionine gamma-synthase] [gn:metb]
CONTIG365	7110281_f2_15	1048	6710	312	104	106	3.5(10)-6	Helicobacter pylori	AC000108	or:helicobacter pylori pn:orf2 le:155 re:595 di:complement nt:orf2 - probable transmembrane regions @ aa 18-36;
CONTIG365	4391268_f2_24	1049	6711	222	74	345	1.6(10)-31	Escherichia coli	b3936	[pn:50s ribosomal protein 131] [gn:rpme]
CONTIG365	15738762_f2_25	1050	6712	765	255	345	1.6(10)-31	Salmonella typhimurium	P43022	hypothetical 15.6 kd protein in rhat 5'region.
CONTIG365	24266301_c1_46	1051	6713	1080	360	1477	1.8(10)-151	Escherichia coli	b3934	[pn:transcriptional repressor cytr] [gn:cytr]
CONTIG365	31885957_c1_48	1052	6714	459	153	124	1.6(10)-7	Escherichia coli	L06547	or:escherichia coli pn:suppressor protein gn:msga le:92 re:961 di:direct sr:escherichia coli (strain
CONTIG365	2246056_c1_49	1053	6715	546	182	757	3.6(10)-75	Escherichia coli	b3932	[pn:heat shock protein hslv] [gn:hslv]
CONTIG365	4400693_c1_50	1054	6716	1074	358	1692	3.0(10)-174	Escherichia coli	b3931	[pn:heat shock protein hslu] [gn:hslu]
CONTIG365	32656378_c2_53	1055	6717	2325	775	3256	0	Escherichia coli	b3935	[pn:primosomal protein replication factor] [gn:pria]
CONTIG365	4313842_c3_57	1056	6718	327	109	518	7.7(10)-50	Escherichia coli	b3938	[pn:metf aporepressor] [gn:metf]
CONTIG365	2365316_c3_63	1057	6719	996	332	421	2.0(10)-81	Escherichia coli	b3933	[pn:cell division protein flsn] [gn:flsn]
CONTIG366	25676576_fl_4	1058	6720	2142	714	2255	6.5(10)-234	Escherichia coli	b0661	[pn:hypothetical protein] [gn:ylea]
CONTIG366	34414182_fl_5	1059	6721	474	158	593	8.5(10)-58	Escherichia coli	b0659	[pn:hypothetical protein] [gn:ybey]

CONTIG366	25503555_f1_6	1060	6722	963	321	1269	2.0(10)-129	Escherichia coli	b0658	[pn:hypothetical protein]
CONTIG366	33703178_f2_14	1061	6723	1227	409	1648	1.3(10)-169	Escherichia coli	b0660	[pn:hypothetical protein]
CONTIG366	4532311_f2_17	1062	6724	768	256	1032	2.6(10)-104	Escherichia coli	b0655	[pn:hypothetical protein in glj 5''region] [gn:ybej]
CONTIG366	15632327_f3_25	1063	6725	1539	513	2176	1.5(10)-225	Escherichia coli	b0657	[pn:apolipoprotein n-acyltransferase] [gn:lnl]
CONTIG366	14897193_c1_35	1064	6726	1188	396	1499	8.5(10)-154	Escherichia coli	b0662	[pn:hypothetical protein]
CONTIG366	4586018_c1_36	1065	6727	297	99	232	1.6(10)-19	Escherichia coli	b0663	[pn:hypothetical protein]
CONTIG366	22678556_c1_37	1066	6728	246	82	139	1.1(10)-9	Escherichia coli	b0667	[pn:hypothetical protein]
CONTIG366	20510955_c2_48	1067	6729	246	82	169	7.4(10)-13	Escherichia coli	b0669	[pn:hypothetical protein]
CONTIG366	10400328_c3_58	1068	6730	378	126	149	9.6(10)-11	Escherichia coli	b0669	[pn:hypothetical protein]
CONTIG367	33828125_f1_9	1069	6731	588	196	736	6.0(10)-73	Escherichia coli	b1158	[pn:dna-invertase pin] [gn:pin]
CONTIG367	13800432_f2_20	1070	6732	501	167	567	4.9(10)-55	Escherichia coli	b1183	[pn:umud protein] [gn:umud]
CONTIG367	21878768_f3_24	1071	6733	531	177	829	8.5(10)-83	Enterobacter agglomerans	B38965	hypothetical protein b (insertion sequence is1222) -enterobacter agglomerans
CONTIG367	17000680_f3_25	1072	6734	645	215	264	6.2(10)-23	Haemophilus influenzae	HI1415	[pn:hypothetical protein]
CONTIG367	11468_f3_26	1073	6735	1023	341	115	0.00016	Bacillus subtilis	xtmA	[pn:pbsx defective prophage terminase] [gn:ykxf]
CONTIG367	859550_f3_30	1074	6736	1299	433	1893	1.5(10)-195	Escherichia coli	b1184	[pn:umuc protein] [gn:umuc]
CONTIG367	16813157_c2_45	1075	6737	360	120	277	2.6(10)-24	Escherichia coli	b1931	[pn:yedg] [gn:yedk]
CONTIG367	284787_c2_50	1076	6738	636	212	98	1.8(10)-5	coliphage T2	P07067	tail fiber protein gp37.
CONTIG367	9823576_c3_61	1077	6739	318	106	496	1.6(10)-47	Escherichia coli	b1931	[pn:yedg] [gn:yedk]
CONTIG368	4964080_c1_40	1078	6740	1146	382	1728	4.5(10)-178	Escherichia coli	b3786	[gn:rffe]

CONTIG368	12969003_c1_44	1079	6741	1266	422	1583	1.1(10)-162	Escherichia coli	b3792	[pn:hypothetical 45.0 kd protein in rffe-rft intergenic region] [gn:yifj]
CONTIG368	1223875_c2_53	1080	6742	981	327	1423	9.5(10)-146	Escherichia coli	b3785	[pn:hypothetical protein] [gn:yifc]
CONTIG368	32695160_c2_56	1081	6743	696	232	620	1.2(10)-60	Escherichia coli	b3790	[pn:hypothetical 19.6 kd protein in rffe-rft intergenic region] [gn:yifh]
CONTIG368	23886_c2_59	1082	6744	1425	475	1579	2.7(10)-162	Escherichia coli	b3793	[pn:4-alpha-l-fucosyltransferase] [gn:rfti]
CONTIG368	3949178_c2_60	1083	6745	753	251	1103	7.7(10)-112	Escherichia coli	b3794	[pn:probable udp-n-acetyl-d-mannosaminuronic acid transferase] [gn:rftm]
CONTIG368	31285313_c2_61	1084	6746	984	328	1153	3.8(10)-117	Escherichia coli	b3795	[pn:probable transport protein yifk] [gn:yifk]
CONTIG368	36413932_c3_64	1085	6747	1281	427	1871	3.2(10)-193	Escherichia coli	b3787	[pn:udp-mannac dehydrogenase] [gn:rftd]
CONTIG368	16927331_c3_66	1086	6748	1305	435	1799	1.3(10)-185	Escherichia coli	b3791	[pn:hypothetical 41.9 kd protein in rffe-rft intergenic region] [gn:yifj]
CONTIG368	14880207_c3_70	1087	6749	1086	362	863	2.1(10)-86	Escherichia coli	b4405	[pn:hypothetical protein]
CONTIG369	12144586_fl_4	1088	6750	873	291	116	0.00024	Chlorella virus SC-1A	U65736	or:chlorella virus sc-1a pn:dna adenine methyltransferase gn:m.cvisile:51 re:1169 di:direct
CONTIG369	14875251_c3_55	1089	6751	1095	365	1548	5.5(10)-159	Serratia marcescens	U62006	rep protein,,rep hi2a,putative rep protein; similar to f plasmid rep
CONTIG370	6413317_c1_37	1090	6752	819	273	166	1.0(10)-10	Methanococcus jannaschii	MJ1187	[pn:dinitrogenase reductase activating glycohydrolase] [gn:drag]
CONTIG370	26056543_c1_42	1091	6753	4371	1457	354	4.4(10)-64	Escherichia coli	b3593	[pn:rhsa protein precursor] [gn:rhsa]
CONTIG370	6369787_c2_45	1092	6754	2283	761	656	1.8(10)-64	Escherichia coli	AF044503	[de:escherichia coli strain ec11 unknown (498), hcp gene, complete cds;and rhsg accessory genetic element vrg protein, core component anddsorf-g1 genes, complete cds.] [pn:vrg protein]
CONTIG371	4538312_fl_10	1093	6755	1071	357	606	3.6(10)-59	Escherichia coli	b0846	[pn:hypothetical protein]
CONTIG371	32531952_f3_34	1094	6756	738	246	1154	3.1(10)-117	Escherichia coli	b0839	[pn:penicillin-binding protein 6 precursor] [gn:dacc]

CONTIG371	4532942_f3_36	1095	6757	1242	414	1804	4.0(10)-186	Escherichia coli	b0842	[pn:hypothetical protein] [gn:cmr]
CONTIG371	4454693_c1_46	1096	6758	489	163	170	5.7(10)-13	Vibrio cholerae	S81006	or:vibrio cholerae pn:hcp gn:hcp le:690 re:1208 di:direct sr:vibrio cholerae o17 nt:28 kda secreted hydrophilic protein; this sequence
CONTIG371	1962837_c1_48	1097	6759	1293	431	1611	1.2(10)-165	Escherichia coli	b3502	[pn:arsenical pump membrane protein] [gn:arsb]
CONTIG371	4119678_c1_49	1098	6760	441	147	605	4.5(10)-59	Escherichia coli	b3503	[pn:arsenate reductase] [gn:arsc]
CONTIG371	10400802_c2_66	1099	6761	1377	459	1517	1.1(10)-155	Escherichia coli	b0845	[pn:hypothetical protein]
CONTIG371	36407965_c2_71	1100	6762	741	247	700	4.0(10)-69	Escherichia coli	b0841	[pn:hypothetical protein]
CONTIG371	21894191_c3_75	1101	6763	429	143	364	1.6(10)-33	Escherichia coli	b3501	[pn:arsenical resistance operon arsefg repressor] [gn:arsr]
CONTIG371	12896930_c3_78	1102	6764	885	295	987	1.5(10)-99	Escherichia coli	b0844	[pn:hypothetical protein]
CONTIG371	2472658_c3_82	1103	6765	801	267	1094	7.0(10)-111	Escherichia coli	b0840	[pn:deoxyribose operon repressor] [gn:deor]
CONTIG372	25500018_f1_2	1104	6766	408	136	636	2.3(10)-62	Escherichia coli	b3294	[pn:50s ribosomal subunit protein 117] [gn:rp1q]
CONTIG372	15808568_f1_11	1105	6767	486	162	756	4.5(10)-75	Escherichia coli	b3284	[pn:smg protein] [gn:smg]
CONTIG372	15626037_f1_13	1106	6768	627	209	817	1.6(10)-81	Escherichia coli	b3282	[pn:hypothetical protein in aroe-smg intergenic region] [gn:yrdc]
CONTIG372	6676963_f2_14	1107	6769	366	122	570	2.3(10)-55	Escherichia coli	b3298	[pn:30s ribosomal subunit protein s13] [gn:rpsm]
CONTIG372	34119062_f2_15	1108	6770	624	208	1008	9.0(10)-102	Escherichia coli	b3296	[pn:30s ribosomal subunit protein s4] [gn:rpsd]
CONTIG372	13859837_f2_16	1109	6771	459	153	632	6.4(10)-62	Escherichia coli	b3292	[pn:hypothetical transcriptional regulator in msc- rplq intergenic region] [gn:yhdm]
CONTIG372	36020676_f2_17	1110	6772	234	78	319	9.4(10)-29	Escherichia coli	P36675	hypothetical 8.1 kd protein in msc- rplq intergenic region.
CONTIG372	6447956_f2_24	1111	6773	306	102	354	1.8(10)-32	Escherichia coli	b3281	[pn:shikimate dehydrogenase] [gn:aroel]
CONTIG372	24333427_f3_25	1112	6774	393	131	522	2.8(10)-50	Escherichia coli	b3297	[pn:30s ribosomal subunit protein s11] [gn:rpsk]

CONTIG372	26445160_f3_27	1113	6775	1008	336	1641	7.5(10)-169	Escherichia coli	b3295	[pn:rpoa] [gn:rpoa]
CONTIG372	26440705_f3_28	1114	6776	510	170	508	8.8(10)-49	Escherichia coli	b3293	[pn:hypothetical 13.9 kd protein in mscL-rplq intergenic region] [gn:yhdn]
CONTIG372	31735932_f3_38	1115	6777	1200	400	683	2.5(10)-67	Escherichia coli	b3286	[pn:hypothetical protein]
CONTIG372	15751342_f3_39	1116	6778	597	199	744	8.5(10)-74	Escherichia coli	b3283	[pn:hypothetical 18.6 kd protein in aroE-smg intergenic region]
CONTIG372	1228918_c1_45	1117	6779	1323	441	1963	5.7(10)-203	Escherichia coli	b3289	[pn:fmj] [gn:sun]
CONTIG372	25667180_c2_56	1118	6780	1410	470	2235	8.5(10)-232	Escherichia coli	b3290	[pn:trka protein of the constitutive k+ transport system trk] [gn:trka]
CONTIG372	29711432_c3_69	1119	6781	528	176	795	3.3(10)-79	Escherichia coli	b3287	[pn:n-formylmethionylaminoacyl-tRNA ornithylase] [gn:dfi]
CONTIG372	26660166_c3_70	1120	6782	972	324	1435	5.0(10)-147	Escherichia coli	b3288	[pn:methionyl-tRNA formyltransferase] [gn:fmt]
CONTIG372	23484430_c3_74	1121	6783	459	153	329	8.1(10)-30	Escherichia coli	b3291	[pn:large conductance mechanosensitive channel] [gn:msc]
CONTIG373	3961086_f1_10	1122	6784	408	136	515	1.6(10)-49	Escherichia coli	b3148	[pn:hypothetical 14.8 kd protein in agai-mtr intergenic region] [gn:yrar]
CONTIG373	6361092_f1_11	1123	6785	600	200	974	3.7(10)-98	Escherichia coli	b3149	[pn:hypothetical 21.1 kd protein in agai-mtr intergenic region] [gn:yrar]
CONTIG373	4312943_f1_12	1124	6786	585	195	693	2.2(10)-68	Escherichia coli	b3150	[pn:hypothetical 20.0 kd protein in agai-mtr intergenic region] [gn:yrar]
CONTIG373	5371093_f2_15	1125	6787	1173	391	949	1.6(10)-95	Escherichia coli	b1621	[pn:pts system, maltose and glucose-specific ii abc component] [gn:malx]
CONTIG373	6464836_f2_16	1126	6788	1185	395	658	1.1(10)-64	Bacillus subtilis	patB	[pn:aminotransferase]
CONTIG373	23696933_f2_20	1127	6789	2169	723	1318	1.1(10)-251	Escherichia coli	b3147	[pn:hypothetical 72.8 kd protein in agai-mtr intergenic region]
CONTIG373	13864777_c1_39	1128	6790	885	295	1225	9.1(10)-125	Escherichia coli	b3146	[pn:hypothetical 31.3 kd protein in agai-mtr intergenic region] [gn:yrar]
CONTIG373	2376312_c2_42	1129	6791	339	113	395	8.3(10)-37	Escherichia coli	b3151	[pn:hypothetical 37.3 kd protein in agai-mtr intergenic region] [gn:yrar]
CONTIG374	5995468_f1_1	1130	6792	291	97	426	4.2(10)-40	Escherichia coli	b2606	[pn:50s ribosomal subunit protein 119] [gn:rpl]
CONTIG374	22854707_f1_16	1131	6793	996	332	1524	1.8(10)-156	Escherichia coli	b2594	[pn:fish suppressor protein sfhb] [gn:sfbb]

CONTIG374	6095760_f2_24	1132	6794	1134	378	1646	2.2(10)-169	Escherichia coli	b2601	[pn:phospho-2-dehydro-3-deoxyheptonate aldolase, tyrosensitive] [gn:aroF]
CONTIG374	16125281_f2_33	1133	6795	387	129	103	4.5(10)-5	Escherichia coli	D90887	or:escherichia coli gn:yfii le:13236 re:14294 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [swissprot accession number p33643];
CONTIG374	33464032_f2_35	1134	6796	1554	518	2377	7.7(10)-247	Escherichia coli	b2592	[pn:clpb protein] [gn:clpb]
CONTIG374	3384667_f3_36	1135	6797	1599	533	214	1.3(10)-30	Pseudomonas aeruginosa	D28119	or:pseudomonas aeruginosa le:3035 re:4450 di:direct sr:pseudomonas aeruginosa, (strain pao1), dna, (clone pin100) nt:putative
CONTIG374	29957332_f3_45	1136	6798	1161	387	1630	1.1(10)-167	Escherichia coli	b2600	[pn:chorismate mutase/prephenate dehydrogenase] [gn:tyra]
CONTIG374	6064015_f3_46	1137	6799	933	311	354	1.8(10)-32	Bacillus subtilis	yvrE	[pn:hypothetical protein]
CONTIG374	35413193_f3_50	1138	6800	750	250	1091	1.5(10)-110	Escherichia coli	b2593	[pn:hypothetical 26.3 kd protein in clpb 5'''region] [gn:yfih]
CONTIG374	36359756_f3_51	1139	6801	1077	359	1542	2.3(10)-158	Escherichia coli	b2592	[pn:clpb protein] [gn:clpb]
CONTIG374	29323783_c1_64	1140	6802	699	233	1092	1.1(10)-110	Escherichia coli	b2595	[pn:hypothetical protein]
CONTIG374	35605011_c1_65	1141	6803	396	132	542	2.2(10)-52	Escherichia coli	b2597	[pn:12.7 kd protein in sfhb-phel intergenic region] [gn:yfia]
CONTIG374	32145012_c2_93	1142	6804	429	143	328	1.0(10)-29	Escherichia coli	b2602	[pn:hypothetical protein in arof-rpls intergenic region] [gn:yfil]
CONTIG374	837807_c2_94	1143	6805	600	200	484	3.1(10)-46	Escherichia coli	b2603	[pn:hypothetical protein]
CONTIG374	26292262_c3_104	1144	6806	1179	393	1642	6.0(10)-169	Escherichia coli	b2599	[pn:chorismate mutase-p/prephenate dehydratase] [gn:phea]
CONTIG374	35728790_c3_112	1145	6807	1242	414	1260	1.8(10)-128	Escherichia coli	b2604	[pn:hypothetical protein in rpls 5'''region] [gn:yfin]
CONTIG374	25992337_c3_113	1146	6808	498	166	651	6.2(10)-64	Escherichia coli	b2605	[pn:hypothetical 17.2 kd protein in rpls 5'''region] [gn:yfib]



CONTIG375	3251076_f1_6	1147	6809	816	272	1019	6.2(10)-103	Escherichia coli	b0933	[pn:hypothetical abc transporter atp-binding protein in pepn-pyrd intergenic region] [gn:ycbe]
CONTIG375	13839807_f1_8	1148	6810	249	83	283	6.0(10)-25	Escherichia coli	b0931	[pn:nicotinate phosphoribosyltransferase] [gn:pncb]
CONTIG375	4413568_f2_12	1149	6811	585	195	812	5.4(10)-81	Escherichia coli	b0937	[pn:hypothetical protein] [gn:ycbp]
CONTIG375	4882842_f2_14	1150	6812	1158	386	1803	5.2(10)-186	Escherichia coli	b0935	[pn:hypothetical protein] [gn:ycbn]
CONTIG375	4410443_f2_15	1151	6813	801	267	1066	6.5(10)-108	Escherichia coli	b0934	[pn:hypothetical protein] [gn:ycbm]
CONTIG375	5362907_f3_27	1152	6814	1014	338	1333	3.2(10)-136	Escherichia coli	b0936	[pn:hypothetical protein]
CONTIG375	1292785_c1_50	1153	6815	306	102	410	2.1(10)-38	Escherichia coli	b0946	[pn:hypothetical protein]
CONTIG375	17070301_c3_60	1154	6816	2718	906	3941	0	Escherichia coli	b0932	[pn:aminopeptidase n] [gn:pepn]
CONTIG375	35641405_c3_69	1155	6817	1050	350	1608	2.3(10)-165	Escherichia coli	b0945	[pn:dihydroorotate dehydrogenase] [gn:pyrd]
CONTIG376	204126_f1_5	1156	6818	1404	468	1811	7.4(10)-187	Escherichia coli	b2509	[pn:exodeoxyribonuclease large subunit] [gn:xsea]
CONTIG376	16273437_f2_13	1157	6819	1107	369	1263	8.6(10)-129	Erwinia carotovora	Q99132	extracellular metalloprotease precursor (ec 3.4.24.-)
CONTIG376	31765664_c1_29	1158	6820	437	145	644	3.3(10)-63	Escherichia coli	b2511	[pn:hypothetical protein]
CONTIG376	6534456_c1_30	1159	6821	1599	533	429	4.2(10)-71	Bacillus subtilis	yclF	[pn:hypothetical protein]
CONTIG376	11207056_c1_36	1160	6822	1491	497	1945	4.7(10)-201	Escherichia coli	b2508	[pn:inosine-5'-monophosphate dehydrogenase] [gn:guab]
CONTIG376	863205_c2_40	1161	6823	348	116	276	3.3(10)-24	Escherichia coli	b2510	[pn:hypothetical protein]
CONTIG376	24319441_c2_46	1162	6824	1686	562	2669	8.8(10)-278	Escherichia coli	b2507	[pn:gmp synthase] [gn:guaa]
CONTIG377	15704432_f1_8	1163	6825	510	170	685	1.5(10)-67	Escherichia coli	b0965	[pn:hypothetical protein]
CONTIG377	160378_f2_11	1164	6826	702	234	541	2.7(10)-52	Escherichia coli	b0959	[pn:hypothetical protein]
CONTIG377	10830131_f3_31	1165	6827	2076	692	2895	1.0(10)-301	Escherichia coli	b0962	[pn:helicase iv] [gn:held]

CONTIG377	35585843_c1_40	1166	6828	759	253	786	3.1(10)-78	Escherichia coli	b0964	[pn:hypothetical protein]
CONTIG377	32616015_c1_42	1167	6829	255	85	99	5.2(10)-5	Paramoecium bursaria	U42580	or: paramoecium bursaria chlorella virus 1 gn:a316r le:158299 re:158976 di:direct nt:papk (17x); similar to pbcv-1 orf a41r, encoded
CONTIG377	34406561_c2_51	1168	6830	1197	399	1740	2.5(10)-179	Escherichia coli	b0967	[pn:hypothetical protein]
CONTIG377	23678515_c2_52	1169	6831	348	116	506	1.3(10)-48	Escherichia coli	b0966	[pn:hypothetical protein]
CONTIG377	32453218_c2_61	1170	6832	453	151	627	2.2(10)-61	Escherichia coli	b0961	[pn:hypothetical protein in held 5''' region] [gn:yccf]
CONTIG377	16150193_c2_62	1171	6833	2178	726	2662	4.9(10)-277	Escherichia coli	b0960	[pn:hypothetical protein]
CONTIG377	2533566_c3_67	1172	6834	483	161	714	1.3(10)-70	Escherichia coli	b0963	[pn:hypothetical 17.3 kd protein in held-sert intergenic region] [gn:ycog]
CONTIG378	5267193_f1_8	1173	6835	672	224	779	1.7(10)-77	Escherichia coli	b3552	[pn:hypothetical 22.2 kd lipoprotein in bisc-cspa intergenic regi]
CONTIG378	32212756_f2_19	1174	6836	585	195	716	8.0(10)-71	Escherichia coli	b3549	[pn:3-methyladenine dna glycosylase i, constitutive] [gn:tag]
CONTIG378	35742087_f2_25	1175	6837	1170	390	573	1.1(10)-55	Bacillus subtilis	kdgR	[pn:transcriptional regulator]
CONTIG378	24009657_f2_27	1176	6838	942	314	762	1.1(10)-75	Bacillus subtilis	kdgK	[pn:2-keto-3-deoxygluconate kinase]
CONTIG378	20754033_f2_28	1177	6839	1338	446	336	1.5(10)-30	Escherichia coli	b3691	[pn:hypothetical 48.8 kd protein in ibpa-gyrb intergenic region]
CONTIG378	24218902_f2_29	1178	6840	1002	334	1353	2.5(10)-138	Escherichia coli	b3553	[pn:putative 2-hydroxyacid dehydrogenase in bisc- cspa intergenic region] [gn:yiae]
CONTIG378	2128201_f3_30	1179	6841	507	169	92	0.028	Escherichia coli	b2321	[pn:div protein] [gn:div]
CONTIG378	16926562_f3_37	1180	6842	552	184	460	1.1(10)-43	Escherichia coli	b3550	[pn:hypothetical 17.1 kd protein in tag-bisc intergenic region] [gn:yiae]
CONTIG378	10838453_c1_64	1181	6843	231	77	249	2.8(10)-20	Escherichia coli	b3551	[pn:biotin sulfoxide reductase] [gn:bisc]
CONTIG378	12975261_c1_66	1182	6844	375	125	211	2.6(10)-17	Escherichia coli	b3548	[pn:hypothetical 26.0 kd protein in prok-tag intergenic region] [gn:yhj]
CONTIG378	2110138_c2_69	1183	6845	324	108	283	6.0(10)-25	Escherichia coli	b3554	[pn:hypothetical 30.2 kd protein in bisc-cspa intergenic region] [gn:yiaf]

CONTIG378	21503156_c2_70	1184	6846	294	98	354	1.8(10)-32	Escherichia coli	b3556	[pn:cold shock protein cspa] [gn:cspa]
CONTIG378	6726562_c2_80	1185	6847	534	178	265	3.7(10)-27	Escherichia coli	b3548	[pn:hypothetical 26.0 kd protein in prok-tag intergenic region] [gn:yhjy]
CONTIG378	36463138_c2_83	1186	6848	1287	429	1411	1.8(10)-144	Escherichia coli	b3546	[pn:64.9 kd protein in prok-tag intergenic region] [gn:yhjw]
CONTIG378	4769752_c3_96	1187	6849	2232	744	2986	0	Escherichia coli	b3551	[pn:biotin sulfoxide reductase] [gn:bisc]
CONTIG378	22659625_c3_102	1188	6850	1224	408	1401	2.1(10)-143	Escherichia coli	b3547	[pn:hypothetical 43.0 kd protein in prok-tag intergenic region] [gn:yhix]
CONTIG379	14580087_f1_8	1189	6851	1194	398	956	3.0(10)-96	Escherichia coli	b1163	[pn:hypothetical protein]
CONTIG379	6855218_c1_40	1190	6852	858	286	266	3.8(10)-23	Escherichia coli	b1559	[pn:hypothetical protein]
CONTIG379	35283591_c2_44	1191	6853	699	233	710	3.5(10)-70	Bacteriophage phi-80	P14814	replication protein 14.
CONTIG379	25628816_c2_45	1192	6854	312	104	129	1.3(10)-8	Bacillus subtilis	ydfG	[pn:hypothetical protein]
CONTIG379	9974138_c2_48	1193	6855	486	162	101	1.2(10)-5	Escherichia coli	b1651	[pn:hypothetical protein] [gn:gloa]
CONTIG379	23542083_c2_51	1194	6856	474	158	510	5.4(10)-49	Escherichia coli	b0550	[pn:hypothetical protein] [gn:rus]
CONTIG379	9896067_c2_53	1195	6857	894	298	1229	3.5(10)-125	Enterobacter agglomerans	B38965	hypothetical protein b (insertion sequence is l222) -enterobacter agglomerans
CONTIG379	35320333_c3_56	1196	6858	459	153	252	1.2(10)-21	Bacteriophage phi-80	P14820	regulatory protein cii.
CONTIG379	13086467_c3_57	1197	6859	990	330	1068	4.0(10)-108	Bacteriophage phi-80	P14815	replication protein 15.
CONTIG379	4145052_c3_64	1198	6860	267	89	157	1.3(10)-11	Escherichia coli	b1061	[pn:dna-damage-inducible protein i] [gn:dini]
CONTIG379	23609515_c3_68	1199	6861	294	98	419	2.3(10)-39	Enterobacter agglomerans	A38965	hypothetical protein a (insertion sequence is l222) -enterobacter agglomerans
CONTIG38	24714410_c3_7	1200	6862	783	261	1109	1.8(10)-112	Escherichia coli	b4371	[pn:hypothetical 37.6 kd protein in dnaat-hold intergenic region] [gn:yjit]
CONTIG380	12695760_f1_15	1201	6863	938	313	1475	3.0(10)-151	Escherichia coli	b2114	[pn:methionyl-tRNA synthetase] [gn:metg]

CONTIG380	3992202_f2_17	1202	6864	1101	367	136	4.7(10)-6	Haemophilus influenzae	U38617	or:haemophilus influenzae pn:hia gn:hia le:251 re:3547 di:direct sr:haemophilus influenzae strain=nontypeable strain 11
CONTIG380	35833441_f2_18	1203	6865	1401	467	1336	1.6(10)-136	Haemophilus influenzae	HI0949	[pn:gamma-aminobutyric acid transaminase]
CONTIG380	792331_f2_19	1204	6866	1485	495	1247	4.2(10)-127	Haemophilus influenzae	HIN_937	[pn:l-2,4-diaminobutyrate decarboxylase] [gn:hi0946]
CONTIG380	22112882_f2_28	1205	6867	399	133	318	1.2(10)-28	Escherichia coli	b2107	[pn:hypothetical protein]
CONTIG380	12932325_f3_42	1206	6868	492	164	154	2.8(10)-11	Escherichia coli	b3550	[pn:hypothetical 17.1 kd protein in tag-bisc intergenic region] [gn:yiacc]
CONTIG380	36042042_c1_54	1207	6869	2250	750	1965	3.5(10)-203	Escherichia coli	b0981	[pn:hypothetical protein in appa 3'''region] [gn:yccc]
CONTIG380	33463915_c1_61	1208	6870	504	168	475	2.7(10)-45	Escherichia coli	b2104	[pn:hypothetical protein] [gn:thim]
CONTIG380	24473131_c2_68	1209	6871	492	164	487	1.5(10)-46	Escherichia coli	b0982	[pn:hypothetical protein] [gn:yecy]
CONTIG380	21515768_c3_78	1210	6872	1140	380	1734	1.1(10)-178	Escherichia coli	b2113	[pn:mrp] [gn:mrp]
CONTIG380	24805387_c3_80	1211	6873	1203	401	1445	4.5(10)-148	Escherichia coli	b0983	[pn:hypothetical protein] [gn:yecz]
CONTIG381	25503383_f1_1	1212	6874	840	280	265	4.9(10)-23	Escherichia coli	b0300	[pn:hypothetical transcriptional regulator in each-beta intergenic region] [gn:ykga]
CONTIG381	4032952_f1_10	1213	6875	762	254	1010	5.5(10)-102	Escherichia coli	b1284	[pn:hypothetical protein]
CONTIG381	1306711_f2_14	1214	6876	243	81	300	9.6(10)-27	Escherichia coli	b4248	[pn:hypothetical 14.6 kd protein in pyrI-argI intergenic region] [gn:yigh]
CONTIG381	14742013_f2_16	1215	6877	1002	334	596	7.9(10)-68	Escherichia coli	b1287	[pn:hypothetical protein]
CONTIG381	4103433_f3_20	1216	6878	813	271	1271	1.2(10)-129	Escherichia coli	b1288	[pn:enoyl-acyl-carrier-protein reductase nadh] [gn:fabi]
CONTIG381	1203412_f3_23	1217	6879	1941	647	3006	0	Escherichia coli	b1286	[pn:exoribonuclease ii] [gn:mb]
CONTIG381	20181625_f3_24	1218	6880	2025	675	2702	2.7(10)-281	Escherichia coli	b1285	[pn:hypothetical protein] [gn:ycir]
CONTIG381	9954777_f3_27	1219	6881	243	81	156	1.8(10)-11	Escherichia coli	b1283	[pn:osmotically inducible lipoprotein b precursor] [gn:osmb]

CONTIG381	25503415_c1_29	1220	6882	351	117	405	7.2(10)-38	Escherichia coli	b1282	[pn:hypothetical 11.4 kd protein in pyrF-osmb intergenic region] [gn:ycih]
CONTIG381	34667968_c2_41	1221	6883	870	290	958	1.8(10)-96	Escherichia coli	b1281	[pn:orotidine-5 <sup>'''</sup> -p decarboxylase] [gn:pyrf]
CONTIG381	34505012_c2_54	1222	6884	1407	469	768	2.5(10)-76	Escherichia coli	b1025	[pn:hypothetical protein] [gn:ycdt]
CONTIG381	15742130_c3_58	1223	6885	915	305	390	2.7(10)-36	Xanthomonas campestris	Y09701	„rpff
CONTIG382	2361512_fl_3	1224	6886	1323	441	2095	5.9(10)-217	Escherichia coli	b0154	[pn:glutamate-1-semialdehyde 2,1-aminomutase] [gn:hemi]
CONTIG382	21650204_c1_36	1225	6887	587	195	810	8.6(10)-81	Escherichia coli	b0149	[pn:peptidoglycan synthetase] [gn:mreb]
CONTIG382	9956407_c1_37	1226	6888	2334	778	1322	1.8(10)-253	Escherichia coli	b0150	[pn:ferrichrome-iron receptor precursor] [gn:fhua]
CONTIG382	15836458_c1_38	1227	6889	2142	714	2080	2.2(10)-215	Escherichia coli	b0153	[pn:ferrichrome transport protein fhub precursor] [gn:fhub]
CONTIG382	35785137_c1_41	1228	6890	825	275	675	1.8(10)-66	Escherichia coli	b0155	[pn:hypothetical protein in heml-pfs intergenic region] [gn:yadq]
CONTIG382	35285941_c2_46	1229	6891	975	325	1086	4.9(10)-110	Escherichia coli	b0152	[pn:ferrichrome-binding periplasmic protein precursor] [gn:fhud]
CONTIG382	21892677_c3_56	1230	6892	933	311	1265	5.2(10)-129	Escherichia coli	b0151	[pn:ferrichrome transport atp-binding protein fhuc] [gn:fhuc]
CONTIG383	35281502_fl_2	1231	6893	2499	833	3387	0	Escherichia coli	b4179	[pn:vacb protein] [gn:vacb]
CONTIG383	14082000_fl_3	1232	6894	795	265	1184	2.0(10)-120	Escherichia coli	b4180	[pn:hypothetical 26.6 kd protein in vacb-aidb intergenic region]
CONTIG383	16804581_f2_12	1233	6895	1392	464	2087	4.2(10)-216	Escherichia coli	b4177	[pn:adenylosuccinate synthetase] [gn:pura]
CONTIG383	29969426_f2_13	1234	6896	501	167	669	7.5(10)-66	Escherichia coli	b4178	[pn:hypothetical 15.6 kd protein in pura-vacb intergenic region] [gn:yjeb]
CONTIG383	21957031_f2_19	1235	6897	1695	565	2361	3.7(10)-245	Escherichia coli	b4187	[pn:aidb protein] [gn:aidb]
CONTIG383	4547906_f3_22	1236	6898	186	62	222	1.8(10)-18	Escherichia coli	b4176	[pn:hypothetical protein] [gn:yjet]
CONTIG383	4067692_c1_36	1237	6899	1383	461	545	1.1(10)-52	Escherichia coli	b1886	[pn:methyl-accepting chemotaxis protein ii] [gn:tar]

CONTIG383	23948293_c1_43	1238	6900	1347	449	154	1.0(10)-7	Shigella flexneri	P30851	vacb protein.
CONTIG383	31875013_c2_52	1239	6901	411	137	101	0.00011	Gluconobacter suboxydans	O05543	[sp:o05543] [de:hypothetical protein in adhs 5'''region (orf3) (fragment)]
CONTIG383	9922177_c3_70	1240	6902	1311	437	570	2.3(10)-55	Escherichia coli	b4111	[pn:proline/betaine transporter] [gn:prop]
CONTIG384	33413132_f2_31	1241	6903	732	244	757	3.6(10)-75	Escherichia coli	b1908	[pn:hypothetical 25.0 kd protein in tyrp-leuz intergenic region]
CONTIG384	19532255_f2_33	1242	6904	264	88	314	3.2(10)-28	Escherichia coli	b1906	[pn:hypothetical 7.3 kd protein in tyrp-rsga intergenic region]
CONTIG384	171887_f3_37	1243	6905	933	311	239	2.7(10)-20	Escherichia coli	b2808	[pn:regulatory protein for glycine cleavage pathway] [gn:gcva]
CONTIG384	32228407_c1_54	1244	6906	342	114	239	2.7(10)-20	Escherichia coli	b1904	[pn:hypothetical protein]
CONTIG384	24881642_c1_64	1245	6907	1293	431	702	2.3(10)-69	Escherichia coli	b3456	[pn:high-affinity branched-chain amino acid transport permease protein livm] [gn:livm]
CONTIG384	15051625_c2_69	1246	6908	1287	429	1480	8.8(10)-152	Escherichia coli	b1907	[pn:tyrosine-specific transport protein] [gn:tyrp]
CONTIG384	4424067_c2_74	1247	6909	996	332	808	1.3(10)-80	Escherichia coli	b3457	[pn:high-affinity branched-chain amino acid transport permease protein livh] [gn:livh]
CONTIG384	2945312_c2_76	1248	6910	765	255	662	4.2(10)-65	Escherichia coli	b3454	[pn:high-affinity branched-chain amino acid transport atp-binding] [gn:livf]
CONTIG384	19725927_c3_85	1249	6911	555	185	787	2.3(10)-78	Escherichia coli	b1905	[pn:ferritin-like protein] [gn:ftn]
CONTIG384	32547515_c3_89	1250	6912	1281	427	765	5.0(10)-76	Escherichia coli	b3460	[pn:leu/ile/val-binding protein precursor] [gn:livj]
CONTIG384	16299092_c3_93	1251	6913	888	296	541	2.2(10)-72	Escherichia coli	b3455	[pn:high-affinity branched-chain amino acid transport atp-binding protein livg] [gn:livg]
CONTIG385	12150208_f1_1	1252	6914	885	295	1078	3.5(10)-109	Escherichia coli	b3424	[pn:glpg protein] [gn:glpg]
CONTIG385	5203167_f2_13	1253	6915	243	81	367	7.7(10)-34	Escherichia coli	b3425	[pn:protein] [gn:glpe]
CONTIG385	786635_f2_14	1254	6916	903	301	1076	5.7(10)-109	Escherichia coli	b3423	[pn:glycerol-3-phosphate regulon repressor] [gn:glpr]

CONTIG385	30507666_f3_29	1255	6917	2433	811	3553	0	Escherichia coli	b3417	[pn:maltodextrin phosphorylase] [gn:malp]
CONTIG385	5208443_f3_30	1256	6918	2091	697	3271	0	Escherichia coli	b3416	[pn:4-alpha-glucanotransferase] [gn:malq]
CONTIG385	22128751_c1_38	1257	6919	267	89	90	0.00044	Rattus norvegicus	P04474	acidic proline-rich protein precursor (clone prp33).
CONTIG385	16289550_c1_39	1258	6920	2760	920	3776	0	Escherichia coli	b3418	[pn:malt] [gn:malt]
CONTIG385	34629027_c3_53	1259	6921	189	63	193	1.1(10)-14	Escherichia coli	b3415	[pn:high-affinity gluconate transporter] [gn:gnt]
CONTIG386	22479712_f2_19	1260	6922	1053	351	931	1.3(10)-93	Escherichia coli	b0419	[pn:hypothetical protein] [gn:yajp]
CONTIG386	3017328_f2_26	1261	6923	1023	341	515	1.6(10)-49	Escherichia coli	b0900	[pn:hypothetical protein] [gn:ycan]
CONTIG386	13166067_f2_29	1262	6924	537	179	135	8.8(10)-9	Escherichia coli	b1967	[pn:hypothetical protein] [gn:yedu]
CONTIG386	4384828_f3_30	1263	6925	186	62	164	2.5(10)-12	Escherichia coli	I41306	hypothetical protein (argf-lacZ region) - escherichia coli
CONTIG386	32667087_f3_37	1264	6926	528	176	169	2.2(10)-12	Haemophilus influenzae	M10821	[pn:gals] [gn:galr]
CONTIG386	15861432_c2_67	1265	6927	1068	356	901	2.0(10)-90	Bacillus subtilis	yrpG	[pn:hypothetical protein]
CONTIG386	34589712_c2_80	1266	6928	453	151	97	0.01799	Methanococcus jannaschii	MJ1643	[pn:chromosome segregation protein]
CONTIG386	3942063_c3_82	1267	6929	909	303	301	7.5(10)-27	Escherichia coli	b0900	[pn:hypothetical protein] [gn:ycan]
CONTIG386	14925383_c3_86	1268	6930	1290	430	700	4.0(10)-69	Helicobacter pylori	HP1193	[pn:aldo-keto reductase, putative]
CONTIG386	11807803_c3_87	1269	6931	975	325	163	9.8(10)-10	Zymomonas mobilis	Q01578	gluconolactonase precursor (ec 3.1.1.17) (d-glucono-delta-lactone lactonohydrolase).
CONTIG387	36415791_f1_1	1270	6932	561	187	748	3.2(10)-74	Escherichia coli	b1181	[pn:hypothetical protein]
CONTIG387	10667163_f3_38	1271	6933	894	298	1177	1.1(10)-119	Escherichia coli	b1187	[pn:fatty acid--fatty acyl responsive dna-binding protein] [gn:fadr]
CONTIG387	29501675_f3_42	1272	6934	1335	445	2065	8.9(10)-214	Escherichia coli	b1189	[pn:d-amino acid dehydrogenase] [gn:dada]
CONTIG387	4816068_f3_43	1273	6935	1080	360	1507	1.2(10)-154	Escherichia coli	b1190	[pn:alanine racemase, catabolic precursor] [gn:dadx]

CONTIG387	34016706_f3_48	1274	6936	633	211	892	1.8(10)-89	Escherichia coli	b1193	[pn:hypothetical protein]
CONTIG387	33805281_c1_49	1275	6937	771	257	845	1.7(10)-84	Escherichia coli	b1194	[pn:hypothetical protein]
CONTIG387	21518877_c1_50	1276	6938	939	313	1176	1.3(10)-119	Escherichia coli	b1192	[pn:hypothetical protein]
CONTIG387	33673808_c1_64	1277	6939	552	184	716	8.0(10)-71	Escherichia coli	b1185	[pn:disulfide bond formation protein b] [gn:dsbb]
CONTIG387	23714768_c2_72	1278	6940	1752	584	2241	2.0(10)-232	Escherichia coli	b1191	[pn:hypothetical protein]
CONTIG387	2089212_c2_78	1279	6941	1581	527	2647	1.8(10)-275	Escherichia coli	b1188	[pn:hypothetical protein in fadr-dada intergenic region] [gn:ycgb]
CONTIG387	437640_c2_81	1280	6942	1620	540	1544	1.3(10)-158	Escherichia coli	b1186	[pn:regulator of intracellular ph] [gn:nhab]
CONTIG387	3986291_c3_95	1281	6943	375	125	226	4.0(10)-18	Escherichia coli	b1186	[pn:regulator of intracellular ph] [gn:nhab]
CONTIG387	13681887_c3_96	1282	6944	186	62	161	5.2(10)-12	Escherichia coli	M83655	or:escherichia coli lc:1764 re:1952 di:direct sr:escherichia coli (sub_strain w1333, strain k-12) dna nt:putative orf
CONTIG388	25900302_f1_4	1283	6945	585	195	844	2.2(10)-84	Escherichia coli	b1627	[pn:hypothetical protein]
CONTIG388	31770037_f1_7	1284	6946	2106	702	2255	6.5(10)-234	Escherichia coli	b1629	[pn:hypothetical protein]
CONTIG388	33594202_f1_8	1285	6947	1053	351	1560	2.8(10)-160	Escherichia coli	b1630	[pn:hypothetical protein] [gn:ydgo]
CONTIG388	4416068_f1_9	1286	6948	696	232	874	1.3(10)-87	Escherichia coli	b1632	[pn:hypothetical protein] [gn:ydgg]
CONTIG388	1269557_f1_12	1287	6949	1548	516	2011	4.7(10)-208	Escherichia coli	b1634	[pn:hypothetical protein]
CONTIG388	24849011_f1_13	1288	6950	639	213	850	5.0(10)-85	Escherichia coli	b1635	[pn:glutathione s-transferase] [gn:gst]
CONTIG388	24346900_f2_23	1289	6951	345	115	252	1.2(10)-21	Escherichia coli	b1625	[pn:hypothetical protein]
CONTIG388	32556875_f2_27	1290	6952	780	260	832	4.0(10)-83	Escherichia coli	b1631	[pn:hypothetical protein]
CONTIG388	22784386_f3_41	1291	6953	468	156	386	7.4(10)-36	Escherichia coli	b1626	[pn:hypothetical protein]



CONTIG388	25672590_f3_42	1292	6954	597	199	757	3.6(10)-75	Escherichia coli	b1628	[pn:hypothetical protein]
CONTIG388	13932707_f3_48	1293	6955	672	224	955	3.7(10)-96	Escherichia coli	b1633	[pn:endonuclease iii] [gn:ntb]
CONTIG388	32522791_c1_57	1294	6956	1371	457	1780	1.3(10)-183	Escherichia coli	b1637	[pn:tyrosyl-trna synthetase] [gn:tyrs]
CONTIG388	30132800_c1_58	1295	6957	927	309	1353	2.5(10)-138	Escherichia coli	b1636	[pn:hypothetical protein]
CONTIG388	4557818_c2_70	1296	6958	1134	378	1559	3.7(10)-160	Escherichia coli	b1640	[pn:hypothetical protein]
CONTIG388	26737658_c2_71	1297	6959	360	120	399	3.1(10)-37	Escherichia coli	b1639	[pn:hypothetical protein in pdxh 5' region] [gn:ydh]
CONTIG388	4785791_c2_72	1298	6960	687	229	1069	3.1(10)-108	Escherichia coli	b1638	[pn:pyridoxamine 5'-phosphate oxidase] [gn:pdxh]
CONTIG388	14511040_c3_109	1299	6961	1158	386	1601	1.3(10)-164	Escherichia coli	b1624	[pn:hypothetical protein]
CONTIG389	4557750_f1_3	1300	6962	774	258	1117	2.6(10)-113	Escherichia coli	b0652	[pn:glutamate/aspartate transport alp-binding protein glt] [gn:glt]
CONTIG389	2379182_f1_4	1301	6963	534	178	783	6.2(10)-78	Escherichia coli	b0651	[pn:hypothetical 33.8 kd protein in leus-glt intergenic region] [gn:ybek]
CONTIG389	13671885_f1_6	1302	6964	507	169	171	4.5(10)-13	Vibrio cholerae	S81006	or:vibrio cholerae pn:hcp gn:hcp le:690 re:1208 di:direct sr:vibrio cholerae o17 nt:28 kda secreted hydrophilic protein; this sequence
CONTIG389	33634701_f1_14	1303	6965	1035	345	1390	3.0(10)-142	Escherichia coli	b0640	[pn:dna polymerase iii, delta subunit] [gn:hol]
CONTIG389	32604625_f1_16	1304	6966	1914	638	3029	0	Escherichia coli	b0635	[pn:mrda] [gn:mrda]
CONTIG389	291465_f1_17	1305	6967	528	176	572	1.5(10)-55	Escherichia coli	b0633	[pn:rare lipoprotein a precursor] [gn:rlpa]
CONTIG389	11769203_f2_18	1306	6968	828	276	1110	1.3(10)-112	Escherichia coli	b0654	[pn:glutamate/aspartate transport system permease protein gltj] [gn:gltj]
CONTIG389	207577_f2_19	1307	6969	675	225	971	7.5(10)-98	Escherichia coli	b0653	[pn:glutamate/aspartate transport system permease protein gltk] [gn:gltk]
CONTIG389	10751002_f2_24	1308	6970	609	203	650	7.9(10)-64	Escherichia coli	b0641	[pn:rare lipoprotein b precursor] [gn:rlpb]

CONTIG389	13714662_f2_27	1309	6971	753	251	887	6.0(10)-89	Escherichia coli	b0639	[pn:hypothetical protein] [gn:yben]
CONTIG389	26801538_f3_34	1310	6972	252	84	336	1.5(10)-30	Escherichia coli	b0655	[pn:hypothetical protein in gltj 5'''region] [gn:ybej]
CONTIG389	9766277_f3_36	1311	6973	582	194	681	4.0(10)-67	Escherichia coli	b0651	[pn:hypothetical 33.8 kd protein in leus-glt intergenic region] [gn:ybek]
CONTIG389	24344502_f3_40	1312	6974	2712	904	4217	0	Escherichia coli	b0642	[pn:leucyl-trna synthetase] [gn:leus]
CONTIG389	31728136_f3_44	1313	6975	405	135	240	2.2(10)-20	Escherichia coli	b0637	[pn:hypothetical 7.7 kd protein in mrdA/pbpa 3'''region] [gn:ybeb]
CONTIG389	1369165_f3_45	1314	6976	471	157	792	7.0(10)-79	Escherichia coli	b0636	[pn:hypothetical 17.3 kd protein in mrdA-phb intergenic region] [gn:ybea]
CONTIG389	33878937_f3_47	1315	6977	1149	383	1390	3.0(10)-142	Escherichia coli	b0634	[pn:rod shape-determining protein rodA] [gn:mrdB]
CONTIG389	787643_c3_97	1316	6978	525	175	688	7.4(10)-68	Escherichia coli	b0643	[pn:hypothetical 18.8 kd protein in leus-glt intergenic region] [gn:ybel]
CONTIG39	22785700_c2_2	1317	6979	444	148	575	7.0(10)-56	Escherichia coli	b1830	[pn:tail-specific protease precursor] [gn:prc]
CONTIG390	32706312_f1_12	1318	6980	354	118	95	0.0015	Kaposi's sarcoma-associated herpesvirus	U93872	[OR:Kaposi's sarcoma-associated herpesvirus] [SR:Kaposi's sarcoma-associated herpesvirus - Human herpesvirus 8] [DE:Kaposi's sarcoma-associated herpesvirus glycoprotein M, DNA replication protein, glycoprotein, DNA replication protein, [pn:hypothetical protein]
CONTIG390	24256457_f2_29	1319	6981	843	281	635	3.1(10)-62	Bacillus subtilis	yusA	[pn:hypothetical protein]
CONTIG390	6926543_f2_30	1320	6982	228	76	274	5.5(10)-24	Escherichia coli	b1675	[pn:hypothetical protein]
CONTIG390	4959691_f3_43	1321	6983	1032	344	816	2.0(10)-81	Bacillus subtilis	yusC	[pn:hypothetical protein]
CONTIG390	9782192_f3_44	1322	6984	663	221	387	5.7(10)-36	Bacillus subtilis	yusB	[pn:hypothetical protein]
CONTIG390	14642966_c1_54	1323	6985	864	288	339	7.0(10)-31	Bacillus subtilis	glnH	[pn:glutamine abc transporter]

CONTIG390	31304138_c1_56	1324	6986	909	303	273	7.0(10)-24	Escherichia coli	b0654	[pn:glutamate/aspartate transport system permease protein gltj] [gn:gltj]
CONTIG390	23722712_c2_72	1325	6987	828	276	645	2.7(10)-63	Bacillus subtilis	glnQ	[pn:glutamine abc transporter]
CONTIG390	30602341_c2_74	1326	6988	735	245	264	6.2(10)-23	Bacillus subtilis	glnP	[pn:glutamine abc transporter]
CONTIG390	35833276_c2_81	1327	6989	876	292	1193	2.2(10)-121	Escherichia coli	b1676	[pn:pyruvate kinase] [gn:pykf]
CONTIG390	4422550_c3_92	1328	6990	1602	534	554	1.7(10)-57	Saccharomyces cerevisiae	YGR155W	[pn:cystathionine beta-synthase] [gn:cys4]
CONTIG390	24401712_c3_93	1329	6991	1158	386	980	8.4(10)-99	Helicobacter pylori	HP0106	[pn:cystathionine gamma-synthase] [gn:metb]
CONTIG391	16620843_f1_1	1330	6992	855	285	211	3.2(10)-35	Escherichia coli	b1559	[pn:hypothetical protein]
CONTIG391	12928328_f1_4	1331	6993	363	121	158	1.1(10)-11	Bacteriophage phi-105	L35561	or:bacteriophage phi-105 pn:holin le:796 re:1170 di:direct sr:bacteriophage phi-105 dna nt:orf2; potential dual start motif; putative [pn:hypothetical protein]
CONTIG391	31922906_f1_6	1332	6994	1770	590	611	1.1(10)-59	Escherichia coli	b1149	
CONTIG391	3230287_f1_9	1333	6995	876	292	106	0.00067	Helicobacter pylori	HP0794	[pn:atp-dependent clp protease proteolytic component] [gn:clpp]
CONTIG391	3913318_f1_10	1334	6996	1221	407	591	1.3(10)-57	Bacteriophage HK97	P49861	major capsid protein precursor (gp5) (head protein).
CONTIG391	26817202_f2_16	1335	6997	666	222	241	1.7(10)-20	Haemophilus influenzae	HI1415	[pn:hypothetical protein]
CONTIG391	21775277_f2_18	1336	6998	948	316	372	2.2(10)-34	Bacteriophage phi-80	S43823	hypothetical protein - phage phi-80
CONTIG391	5156338_f2_21	1337	6999	486	162	101	9.4(10)-5	Mycobacterium tuberculosis	Z95586	unknown,,mtcy336.26,mtcy336.26, len
CONTIG391	35329530_f3_35	1338	7000	1326	442	524	1.8(10)-50	Bacteriophage HK97	P49859	portal protein (gp3).
CONTIG392	5162802_f1_5	1339	7001	471	157	214	1.3(10)-17	Escherichia coli	b1160	[pn:hypothetical protein]
CONTIG392	33447837_f1_6	1340	7002	1335	445	1719	4.0(10)-177	Escherichia coli	b1609	[pn:unknown sensor protein in terminator region] [gn:rsib]
CONTIG392	5315675_f2_15	1341	7003	1419	473	1891	2.5(10)-195	Escherichia coli	b1605	[pn:hypothetical protein]

CONTIG392	26757882_f2_16	1342	7004	735	245	994	2.7(10)-100	Escherichia coli	b1606	[pn:hypothetical protein] [gn:ydgb]
CONTIG392	7161263_f2_17	1343	7005	765	255	924	7.2(10)-93	Escherichia coli	b1608	[pn:rsta] [gn:rsta]
CONTIG392	22054561_f3_28	1344	7006	990	330	1008	9.0(10)-102	Escherichia coli	b1604	[pn:hypothetical protein]
CONTIG392	15761550_f3_30	1345	7007	942	314	1161	5.5(10)-118	Escherichia coli	b1610	[pn:dna replication terminus site-binding protein, dna sequence-specific contrahellicase] [gn:tus]
CONTIG392	1385410_f3_36	1346	7008	1215	405	1645	2.8(10)-169	Escherichia coli	b1613	[pn:mannose-6-phosphate isomerase] [gn:mana]
CONTIG392	282081_f3_37	1347	7009	1608	536	1656	2.0(10)-170	Escherichia coli	b1614	[pn:hypothetical protein]
CONTIG392	4960968_c1_51	1348	7010	345	115	425	5.5(10)-40	Escherichia coli	b1607	[pn:hypothetical protein] [gn:ydgc]
CONTIG392	4806553_c2_63	1349	7011	1398	466	1910	2.3(10)-197	Escherichia coli	b1611	[pn:fumarate hydratase class ii] [gn:fumc]
CONTIG392	23438526_c3_81	1350	7012	1716	572	2604	6.7(10)-271	Escherichia coli	b1612	[pn:fumarate hydratase class i] [gn:fuma]
CONTIG393	5265713_f1_2	1351	7013	939	313	1400	2.6(10)-143	Escherichia coli	b0003	[pn:homoserine kinase] [gn:thrb]
CONTIG393	2931566_f1_3	1352	7014	1290	430	2030	4.5(10)-210	Escherichia coli	b0004	[pn:threonine synthase] [gn:thrc]
CONTIG393	14258462_f1_8	1353	7015	969	323	1541	3.0(10)-158	Escherichia coli	b0008	[pn:transaldolase b] [gn:talb]
CONTIG393	25651656_f2_19	1354	7016	537	179	188	7.0(10)-15	Escherichia coli	b0005	[pn:hypothetical protein]
CONTIG393	22382155_f2_22	1355	7017	1350	450	323	3.5(10)-29	Escherichia coli	b3523	[pn:hypothetical metabolite transport protein in tref-kdkg intergenic region] [gn:yhje]
CONTIG393	6900287_f3_25	1356	7018	2466	822	3632	0	Escherichia coli	b0002	[pn:aspartokinase i/homoserine dehydrogenase i] [gn:thra]
CONTIG393	31411253_f3_34	1357	7019	750	250	900	2.5(10)-90	Escherichia coli	b0009	[pn:molybdopterin biosynthesis mog protein] [gn:mog]
CONTIG393	30210303_c2_62	1358	7020	1500	500	1004	2.3(10)-101	Escherichia coli	b0007	[pn:hypothetical 51.7 kd protein in thr-talb intergenic region] [gn:yaa]
CONTIG393	5162711_c2_63	1359	7021	840	280	1183	2.6(10)-120	Escherichia coli	b0006	[pn:hypothetical 29.6 kd protein in thr-talb intergenic region] [gn:yaaa]

CONTIG393	29850786_c3_74	1360	7022	564	188	759	2.2(10)-75	Escherichia coli	b0010	[pn:hypothetical 20.1 kd protein in mog-higa intergenic region] [gn:yaah]
CONTIG394	25932_f2_20	1361	7023	906	302	1412	1.3(10)-144	Escherichia coli	b0819	[pn:hypothetical protein]
CONTIG394	6447130_f2_33	1362	7024	1008	336	1393	1.5(10)-142	Escherichia coli	b0815	[pn:hypothetical protein] [gn:ybip]
CONTIG394	12145327_c1_51	1363	7025	1110	370	1327	1.3(10)-135	Escherichia coli	b0818	[pn:hypothetical protein] [gn:ybir]
CONTIG394	26852088_c1_57	1364	7026	1551	517	364	1.6(10)-33	Escherichia coli	b4080	[pn:hypothetical 53.4 kd protein in fdhF-phn intergenic region]
CONTIG394	5199068_c1_64	1365	7027	726	242	467	1.8(10)-44	Bacillus subtilis	ydhQ	[pn:hypothetical protein]
CONTIG394	4802193_c2_68	1366	7028	483	161	707	7.2(10)-70	Escherichia coli	b0817	[pn:hypothetical protein]
CONTIG394	21773532_c2_73	1367	7029	1590	530	762	1.1(10)-75	Haemophilus influenzae	HI0897	[pn:multidrug resistance protein] [gn:emrb]
CONTIG394	24253551_c2_77	1368	7030	1398	466	871	3.0(10)-87	Bacillus subtilis	yckE	[pn:hypothetical protein]
CONTIG394	10336090_c2_78	1369	7031	1338	446	528	6.7(10)-51	Bacillus subtilis	ywbA	[pn:hypothetical protein] [gn:ipa-16d]
CONTIG394	26306507_c3_85	1370	7032	1098	366	330	6.5(10)-35	Haemophilus influenzae	HI0898	[pn:multidrug resistance protein] [gn:emra]
CONTIG395	29766093_f1_9	1371	7033	1062	354	1078	3.5(10)-109	Escherichia coli	b1428	[pn:hypothetical protein]
CONTIG395	13865937_f2_24	1372	7034	1254	418	703	1.8(10)-69	Bacillus subtilis	pepT	[pn:peptidase t]
CONTIG395	32509638_f3_45	1373	7035	846	282	328	1.0(10)-29	Escherichia coli	b2808	[pn:regulatory protein for glycine cleavage pathway] [gn:gcva]
CONTIG395	24804715_c1_50	1374	7036	1524	508	481	6.4(10)-46	Bacillus subtilis	pnbA	[pn:intracellular esterase b] [gn:eslb]
CONTIG395	21520675_c1_53	1375	7037	1200	400	932	1.0(10)-93	Escherichia coli	b0600	[pn:hypothetical protein]
CONTIG395	1171956_c1_54	1376	7038	1746	582	2562	1.8(10)-266	Escherichia coli	b1424	[pn:hypothetical protein]
CONTIG395	7271905_c1_55	1377	7039	645	215	627	2.2(10)-61	Escherichia coli	b1427	[pn:ribosomal-protein-serine acetyltransferase] [gn:riml]
CONTIG395	4726568_c1_62	1378	7040	603	201	861	3.3(10)-86	Escherichia coli	b1430	[pn:tellurite resistance protein tehB] [gn:tehb]

CONTIG395	2034780_c2_74	1379	7041	861	287	565	8.0(10)-55	Escherichia coli	b2310	[pn:lysine-arginine-ornithine-binding periplasmic protein precursor] [gn:argt]
CONTIG395	24412506_c2_80	1380	7042	1008	336	1338	9.8(10)-137	Escherichia coli	b1429	[pn:tellurite resistance protein teha] [gn:teha]
CONTIG395	34181577_c3_94	1381	7043	1599	533	1893	1.5(10)-195	Escherichia coli	b3544	[pn:periplasmic dipeptide transport protein precursor] [gn:dppa]
CONTIG396	976575_f2_25	1382	7044	825	275	778	2.1(10)-77	Escherichia coli	b2558	[pn:hypothetical 53.2 kd protein in purt-dpj intergenic region] [gn:yfhd]
CONTIG396	35807756_f3_26	1383	7045	777	259	1109	1.8(10)-112	Escherichia coli	b2552	[pn:flavohemoprotein] [gn:hmpa]
CONTIG396	36369777_f3_29	1384	7046	786	262	160	1.3(10)-10	Aeromonas hydrophila	U56832	[DE:Aeromonas hydrophila FK506 binding protein (fkpA) gene, completecds in 3.9 kb fragment.] [NT:ORF5; no significant similarity with known] [LE:2969] [RE:3721] [DI:complement]
CONTIG396	10366717_c1_47	1385	7047	1485	495	1795	3.7(10)-185	Escherichia coli	b2556	[pn:hypothetical protein] [gn:yfhk]
CONTIG396	4410193_c1_51	1386	7048	348	116	553	1.5(10)-53	Escherichia coli	b2553	[pn:nitrogen regulatory protein p-ii] [gn:glnb]
CONTIG396	14660413_c2_56	1387	7049	1365	455	2071	2.1(10)-214	Escherichia coli	b2554	[pn:hypothetical protein in glnb 5''' region] [gn:yfha]
CONTIG396	32675833_c3_61	1388	7050	3918	1306	6216	0	Escherichia coli	b2557	[pn:phosphoribosylformylglycineamide synthetase] [gn:purI]
CONTIG396	4883533_c3_63	1389	7051	771	257	574	8.9(10)-56	Escherichia coli	b2555	[pn:hypothetical 27.3 kd protein in glnb 5''' region] [gn:yfhg]
CONTIG397	33635886_f1_11	1390	7052	1293	431	2031	3.6(10)-210	Escherichia coli	b1136	[pn:isocitrate dehydrogenase] [gn:icd]
CONTIG397	13105186_f1_12	1391	7053	300	100	340	5.5(10)-31	Escherichia coli	b1295	[pn:hypothetical protein]
CONTIG397	25423317_f2_15	1392	7054	1035	345	1559	3.7(10)-160	Escherichia coli	b1127	[pn:peptidase t] [gn:pept]
CONTIG397	24032750_f3_46	1393	7055	282	94	332	3.8(10)-30	Escherichia coli	b1795	[pn:hypothetical protein]
CONTIG397	32459633_c1_49	1394	7056	669	223	942	9.0(10)-95	Escherichia coli	b1135	[pn:hypothetical protein] [gn:ymfc]
CONTIG397	14630207_c1_51	1395	7057	1287	429	1694	1.8(10)-174	Escherichia coli	b1133	[pn:hypothetical protein in purb 5''' region] [gn:ycfb]

NA aa

CONTIG397	14297282_c1_52	1396	7058	657	219	865	1.3(10)-86	Escherichia coli	b1132	[pn:hypothetical 22.9 kd protein in purb 5'''region] [gn:ycfc]
CONTIG397	2688338_c1_55	1397	7059	1143	381	1745	7.2(10)-180	Escherichia coli	b1128	[pn:hypothetical protein in pept-phoq intergenic region] [gn:ycfd]
CONTIG397	22925191_c2_67	1398	7060	1425	475	2217	7.0(10)-230	Escherichia coli	b1131	[pn:adenylosuccinate lyase] [gn:purb]
CONTIG397	1058467_c3_71	1399	7061	297	99	227	1.2(10)-18	Escherichia coli	b1794	[pn:hypothetical protein]
CONTIG397	34025381_c3_74	1400	7062	1548	516	300	3.7(10)-24	Escherichia coli	b4061	[pn:hypothetical 60.8 kd protein in ssb-soxs intergenic region] [gn:yjcc]
CONTIG397	5101693_c3_76	1401	7063	483	161	684	2.0(10)-67	Escherichia coli	b1134	[pn:hypothetical protein]
CONTIG397	6519052_c3_82	1402	7064	720	240	936	3.8(10)-94	Escherichia coli	b1130	[pn:transcriptional regulatory protein phop] [gn:phop]
CONTIG397	5367202_c3_83	1403	7065	1464	488	1829	9.0(10)-189	Escherichia coli	b1129	[pn:sensor protein phoq] [gn:phoq]
CONTIG398	22117656_f1_4	1404	7066	1431	477	1108	2.2(10)-112	Escherichia coli	b3657	[pn:hypothetical 51.0 kd protein in glts-selc intergenic region] [gn:yicj]
CONTIG398	1301712_f1_5	1405	7067	2052	684	2519	7.0(10)-262	Erwinia herbicola	Q01336	hypothetical protein in crte 3region (orf2) (fragment)
CONTIG398	5324127_f1_9	1406	7068	411	137	429	2.1(10)-40	Escherichia coli	b1792	[pn:hypothetical protein]
CONTIG398	24821017_f2_13	1407	7069	279	93	110	1.3(10)-6	Escherichia coli	b1141	[pn:hypothetical protein]
CONTIG398	15728292_f2_19	1408	7070	513	171	400	2.3(10)-37	Escherichia coli	b1789	[pn:hypothetical protein]
CONTIG398	21581381_f2_26	1409	7071	711	237	734	7.5(10)-75	Escherichia coli	b1794	[pn:hypothetical protein]
CONTIG398	31382327_f3_31	1410	7072	1143	381	930	1.3(10)-95	Escherichia coli	b1140	[pn:hypothetical protein]
CONTIG398	20813561_f3_32	1411	7073	537	179	667	1.2(10)-65	Escherichia coli	b1787	[pn:hypothetical protein]
CONTIG398	4725963_f3_37	1412	7074	1197	399	1339	7.7(10)-137	Escherichia coli	b1791	[pn:hypothetical protein]
CONTIG398	30081918_c1_44	1413	7075	753	251	216	5.5(10)-17	Bacillus subtilis	ctrA	[pn:ctp synthase] [gn:pyrg]
CONTIG398	5112842_c2_61	1414	7076	267	89	211	2.6(10)-17	Escherichia coli	b1793	[pn:hypothetical protein]

CONTIG398	270387_c2_66	1415	7077	825	275	833	3.2(10)-83	Escherichia coli	b1790	[pn:hypothetical protein]
CONTIG399	13167057_fl_1	1416	7078	567	189	888	4.7(10)-89	Escherichia coli	b0860	[pn:arginine-binding periplasmic protein 2 precursor] [gn:artj]
CONTIG399	3963966_fl_20	1417	7079	333	111	357	8.8(10)-33	Haemophilus influenzae	H11532	[pn:grx] [gn:grxa]
CONTIG399	11758450_fl_23	1418	7080	882	294	99	0.03699	Saccharomyces cerevisiae	S57378	hypothetical protein y091w - yeast (saccharomycescerevisiae)
CONTIG399	20990640_fl_24	1419	7081	1803	601	2483	4.5(10)-258	Escherichia coli	b0847	[pn:hypothetical protein]
CONTIG399	36385956_c1_65	1420	7082	480	160	124	4.2(10)-8	Escherichia coli	P12009	multi-copy suppressor of livr.
CONTIG399	33260086_c1_66	1421	7083	528	176	373	1.8(10)-34	Escherichia coli	b0848	[pn:hypothetical protein]
CONTIG399	16525416_c1_68	1422	7084	1086	362	1365	1.3(10)-139	Escherichia coli	b0852	[pn:ribosomal protein s6 modification protein] [gn:rimk]
CONTIG399	32211630_c1_69	1423	7085	513	171	424	7.0(10)-40	Escherichia coli	b0853	[pn:hypothetical protein]
CONTIG399	26303807_c1_74	1424	7086	876	292	1150	8.1(10)-117	Escherichia coli	b0857	[pn:putrescine transport system permease protein poti] [gn:poti]
CONTIG399	20832265_c1_75	1425	7087	519	173	691	3.6(10)-68	Escherichia coli	b0858	[pn:hypothetical protein] [gn:ybj0]
CONTIG399	12932812_c1_80	1426	7088	1542	514	320	1.3(10)-26	Bacillus subtilis	phoR	[pn:two-component sensor histidine kinase]
CONTIG399	16589783_c2_88	1427	7089	1218	406	1754	8.0(10)-181	Escherichia coli	b0855	[pn:putrescine transport atp-binding protein potg] [gn:potg]
CONTIG399	4354193_c2_89	1428	7090	963	321	1321	6.2(10)-135	Escherichia coli	b0856	[pn:putrescine transport system permease protein poti] [gn:poti]
CONTIG399	4078143_c2_91	1429	7091	1137	379	1608	2.3(10)-165	Escherichia coli	b0859	[pn:hypothetical protein] [gn:ybjf]
CONTIG399	16970910_c3_101	1430	7092	813	271	994	2.7(10)-100	Escherichia coli	b0851	[pn:modulator of drug activity a] [gn:mdaa]
CONTIG399	33594691_c3_102	1431	7093	1137	379	1799	1.3(10)-185	Escherichia coli	b0854	[pn:putrescine-binding periplasmic protein precursor] [gn:potf]
CONTIG399	25431561_c3_110	1432	7094	729	243	517	9.8(10)-50	Bacillus subtilis	yclJ	[pn:hypothetical protein]
CONTIG400	24025326_fl_5	1433	7095	2097	699	3000	0	Escherichia coli	b0984	[pn:hypothetical protein] [gn:ymca]



CONTIG400	5292787_f1_7	1434	7096	477	159	368	6.0(10)-34	Escherichia coli	b0982	[pn:hypothetical protein] [gn:yecy]
CONTIG400	3908300_f1_8	1435	7097	2187	729	2495	2.3(10)-259	Escherichia coli	b0981	[pn:hypothetical protein in appa 3'''region] [gn:yccc]
CONTIG400	15626550_f2_17	1436	7098	1818	606	1732	1.7(10)-178	Escherichia coli	b3647	[pn:hypothetical 63.2 kd protein in gmk 5'''region] [gn:yicf]
CONTIG400	31802025_f2_20	1437	7099	825	275	537	7.4(10)-52	Escherichia coli	b0985	[pn:hypothetical protein] [gn:ymcb]
CONTIG400	6445327_f2_26	1438	7100	1134	378	1617	2.7(10)-166	Escherichia coli	b0983	[pn:hypothetical protein] [gn:yeccz]
CONTIG400	4765950_f3_42	1439	7101	678	226	707	7.2(10)-70	Escherichia coli	b0986	[pn:hypothetical protein] [gn:ymcc]
CONTIG400	35413343_f3_45	1440	7102	249	83	130	1.0(10)-8	Escherichia coli	b0982	[pn:hypothetical protein] [gn:yecy]
CONTIG400	1268807_f3_50	1441	7103	840	280	954	4.7(10)-96	Escherichia coli	b3643	[pn:rmase ph] [gn:rph]
CONTIG400	4153_f3_51	1442	7104	690	230	1030	4.2(10)-104	Escherichia coli	b3642	[pn:orotate phosphoribosyltransferase] [gn:pyre]
CONTIG400	26660317_c1_57	1443	7105	219	73	250	5.2(10)-21	Escherichia coli	b3639	[pn:dfp protein] [gn:dfp]
CONTIG400	24017165_c1_59	1444	7106	684	228	922	1.2(10)-92	Escherichia coli	b3641	[pn:ttk protein] [gn:ttk]
CONTIG400	25488563_c1_60	1445	7107	918	306	1362	2.7(10)-139	Escherichia coli	b3644	[pn:33.2 kd protein in dind-rph intergenic region] [gn:yicc]
CONTIG400	24397161_c1_80	1446	7108	579	193	853	2.3(10)-85	Escherichia coli	b3648	[pn:5'''guanylate kinase] [gn:gmk]
CONTIG400	2032005_c2_86	1447	7109	645	215	709	4.4(10)-70	Escherichia coli	b3646	[pn:hypothetical 22.0 kd protein in rph-gmk intergenic region] [gn:yicg]
CONTIG400	16977086_c3_99	1448	7110	570	190	737	4.7(10)-73	Escherichia coli	b3640	[pn:deoxyuridine 5'''-triphosphate nucleotidohydrolase] [gn:dut]
CONTIG401	36460400_f1_11	1449	7111	1578	526	2446	3.7(10)-254	Escherichia coli	b3749	[pn:high affinity ribose transport protein] [gn:rbsa]
CONTIG401	10256930_f1_13	1450	7112	270	90	163	8.4(10)-12	Escherichia coli	b3752	[pn:ribokinase] [gn:rbsk]
CONTIG401	14196916_f1_14	1451	7113	996	332	1421	1.6(10)-145	Escherichia coli	b3753	[pn:rbs repressor] [gn:rbsr]
CONTIG401	14337755_f2_24	1452	7114	1872	624	2382	2.2(10)-247	Escherichia coli	b3747	[pn:kup]

CONTIG401	23714652_f2_27	1453	7115	831	277	945	4.2(10)-95	Escherichia coli	b3750	[pn:high affinity ribose transport protein] [gn:rbse]
CONTIG401	14650302_f3_35	1454	7116	996	332	1573	1.2(10)-161	Escherichia coli	b3744	[pn:aspartate--ammonia ligase] [gn:asna]
CONTIG401	9817191_f3_41	1455	7117	450	150	640	9.0(10)-63	Escherichia coli	b3748	[pn:high affinity ribose transport protein rbsd] [gn:rbsd]
CONTIG401	13917556_f3_42	1456	7118	546	182	90	0.02999	Escherichia coli	b3750	[pn:high affinity ribose transport protein] [gn:rbse]
CONTIG401	2854076_f3_43	1457	7119	918	306	1354	2.0(10)-138	Escherichia coli	b3751	[pn:periplasmic ribose-binding protein precursor] [gn:rbsb]
CONTIG401	24319718_f3_44	1458	7120	897	299	1009	7.0(10)-102	Escherichia coli	b3752	[pn:ribokinase] [gn:rbsk]
CONTIG401	29800187_c1_49	1459	7121	441	147	451	9.5(10)-43	Escherichia coli	b3755	[pn:hypothetical 20.8 kd protein in rbsr-rsc intergenic region] [gn:yiep]
CONTIG401	11067932_c2_80	1460	7122	1464	488	1648	1.3(10)-169	Escherichia coli	b3745	[pn:hypothetical 49.6 kd protein in asna 3''' region]
CONTIG401	24819702_c3_83	1461	7123	1434	478	1844	2.2(10)-190	Escherichia coli	b3754	[pn:hypothetical 51.5 kd protein in rbsr-rsc intergenic region] [gn:yieo]
CONTIG401	9880166_c3_97	1462	7124	1587	529	2038	6.5(10)-211	Escherichia coli	b3746	[pn:hypothetical 57.4 kd protein in asna-kup intergenic region] [gn:yien]
CONTIG401	4007813_c3_100	1463	7125	468	156	605	4.5(10)-59	Escherichia coli	b3743	[pn:regulatory protein] [gn:asnc]
CONTIG401	196080_c3_101	1464	7126	540	180	559	3.5(10)-54	Escherichia coli	b3742	[pn:involved in modulation of initiation at oric] [gn:mioc]
CONTIG402	7082715_f1_3	1465	7127	978	326	1390	3.0(10)-142	Escherichia coli	b2378	[pn:hypothetical protein]
CONTIG402	6837836_f1_8	1466	7128	735	245	1148	1.3(10)-116	Escherichia coli	b2381	[pn:hypothetical protein]
CONTIG402	20703533_f1_11	1467	7129	1260	420	1163	3.3(10)-118	Escherichia coli	b2389	[pn:hypothetical protein]
CONTIG402	19724077_f2_15	1468	7130	957	319	1145	2.7(10)-116	Escherichia coli	b2347	[pn:hypothetical 34.5 kd protein in argw 5'''region] [gn:yfde]
CONTIG402	24782786_f3_31	1469	7131	381	127	94	0.00072	Pseudomonas sp.	L81125	pn:monooxygenase subunit le:502 re:2016 di:direct sr:pseudomonas sp (strain int37) dna
CONTIG402	6343752_f3_38	1470	7132	1725	575	2476	2.5(10)-257	Escherichia coli	b2380	[pn:hypothetical protein]

CONTIG402	24745641_c1_45	1471	7133	1311	437	1930	1.8(10)-199	Enterobacter cloacae	P23234	indole-3-pyruvate decarboxylase (ec 4.1.1.74) (indolepyruvate decarboxylase).
CONTIG402	25473516_c1_51	1472	7134	1254	418	2026	1.2(10)-209	Escherichia coli	b2379	[pn: hypothetical protein]
CONTIG402	33474091_c2_61	1473	7135	981	327	1556	7.7(10)-160	Escherichia coli	b2388	[pn: glucokinase] [gn: glk]
CONTIG402	11770652_c2_72	1474	7136	1416	472	1587	4.0(10)-163	Escherichia coli	b1490	[pn: hypothetical protein]
CONTIG402	7292291_c2_74	1475	7137	888	296	1111	1.1(10)-112	Escherichia coli	b2346	[pn: vacj lipoprotein precursor] [gn: vacj]
CONTIG403	25567885_fl_10	1476	7138	525	175	898	4.0(10)-90	Escherichia coli	b0264	[pn: insertion element is1 protein insb] [gn: insb_2]
CONTIG403	2911290_fl_15	1477	7139	234	78	254	7.2(10)-22	Escherichia coli	S40546	hypothetical protein - escherichia coli
CONTIG403	24489756_f2_28	1478	7140	942	314	101	0.05299	Methanococcus jannaschii	MJ1322	[pn: purine ntpase]
CONTIG403	6735393_f2_29	1479	7141	285	95	446	3.2(10)-42	Escherichia coli	b4294	[pn: insertion element is1f protein insa] [gn: insa_7]
CONTIG403	25567885_f2_36	1480	7142	525	175	829	8.5(10)-83	Escherichia coli	b0264	[pn: insertion element is1 protein insb] [gn: insb_2]
CONTIG403	2911290_f3_50	1481	7143	234	78	296	2.6(10)-26	Escherichia coli	S40546	hypothetical protein - escherichia coli
CONTIG403	6735393_f3_55	1482	7144	285	95	446	3.2(10)-42	Escherichia coli	b4294	[pn: insertion element is1f protein insa] [gn: insa_7]
CONTIG403	21666540_f3_56	1483	7145	729	243	1128	1.8(10)-114	Transposon Tn1525	M12900	or: transposon tn1525 gn: p12 le: 996 re: >1721 di: direct sr: transposon tn1525 dna nt: putative
CONTIG403	14885165_c1_59	1484	7146	297	99	377	6.7(10)-35	Escherichia coli	D93826	hypothetical 11k protein (insertion sequence is1) - escherichia coli this protein is coded by the insertion sequence is1.
CONTIG403	11909633_c1_60	1485	7147	681	227	420	1.8(10)-39	Escherichia coli	b3025	[pn: hypothetical protein] [gn: ygix]
CONTIG403	35556507_c1_71	1486	7148	1413	471	1826	1.8(10)-188	Pseudomonas aeruginosa	Y10528	[PN: cyanide insensitive terminal oxidase] [GN: cioA] [DE: P.aeruginosa cioA and cioB genes.] [LE: 276] [RE: 1742] [DI: direct]

CONTIG403	11891882_c2_73	1487	7149	738	246	1268	2.6(10)-129	Salmonella ordonez	S34451	hypothetical protein (insertion sequence is261) - salmonellaordonez plasmid pip173
CONTIG403	14885165_c2_77	1488	7150	297	99	390	2.7(10)-36	Escherichia coli	D93826	hypothetical 11k protein (insertion sequence is1) -escherichia coli this protein is coded by the insertion sequence is1.
CONTIG403	36501937_c3_89	1489	7151	1386	462	347	1.0(10)-31	Escherichia coli	b3026	[pn:hypothetical protein] [gn:ygy]
CONTIG403	35410666_c3_96	1490	7152	348	116	104	5.7(10)-6	Escherichia coli	Q52331	transcriptional repressor protein korc.
CONTIG403	24109468_c3_99	1491	7153	1041	347	1016	1.3(10)-102	Pseudomonas aeruginosa	Y10528	[PN:cyanide insensitive terminal oxidase] [GN:cioB] [DE:P.aeruginosa cioA and cioB genes.] [LE:1746] [RE:2753] [DI:direct]
CONTIG404	4453427_fl_14	1492	7154	501	167	253	9.1(10)-22	Escherichia coli	b1164	[pn:hypothetical protein]
CONTIG404	34413938_fl_15	1493	7155	381	127	212	2.0(10)-17	Escherichia coli	b1166	[pn:hypothetical protein]
CONTIG404	21509432_f2_40	1494	7156	2184	728	2478	1.5(10)-257	Escherichia coli	b4079	[pn:formate dehydrogenase, formate- hydrogen-lyase-linked, selenocysteine-containing polypeptide] [gn:fdhf]
CONTIG404	4882768_c1_67	1495	7157	687	229	413	1.0(10)-38	Escherichia coli	b4316	[pn:chaperone protein involved in biogenesis of type 1 fimbriae] [gn:fimc]
CONTIG404	6506280_c1_68	1496	7158	2571	857	1485	2.6(10)-152	Escherichia coli	b3144	[pn:hypothetical outer membrane usher protein in agal-mtr intergenic region] [gn:yraj]
CONTIG404	16130091_c1_71	1497	7159	681	227	176	1.3(10)-13	Escherichia coli	b4319	[pn:fing protein precursor] [gn:fimg]
CONTIG404	14485390_c1_77	1498	7160	1284	428	1312	5.5(10)-134	Escherichia coli	b1163	[pn:hypothetical protein]
CONTIG404	35431691_c1_78	1499	7161	852	284	812	5.4(10)-81	Escherichia coli	b1162	[pn:hypothetical protein]
CONTIG404	4008518_c1_79	1500	7162	378	126	563	1.3(10)-54	Escherichia coli	b1684	[pn:hypothetical protein] [gn:ydjc]

CONTIG404	14900762_c1_82	1501	7163	1317	439	1843	3.0(10)-190	Escherichia coli	b1681	[pn:hypothetical protein]
CONTIG404	35335205_c2_87	1502	7164	1065	355	145	1.1(10)-9	Escherichia coli	b0530	[pn:hypothetical protein] [gn:sfma]
CONTIG404	22160268_c2_99	1503	7165	534	178	515	1.6(10)-49	Escherichia coli	b1679	[pn:hypothetical protein]
CONTIG404	12382906_c3_117	1504	7166	846	282	1343	2.8(10)-137	Escherichia coli	b1683	[pn:hypothetical protein]
CONTIG404	12681461_c3_118	1505	7167	702	234	1158	1.2(10)-117	Escherichia coli	b1683	[pn:hypothetical protein]
CONTIG404	4479693_c3_119	1506	7168	756	252	1168	1.0(10)-118	Escherichia coli	b1682	[pn:hypothetical protein]
CONTIG404	15913557_c3_120	1507	7169	1269	423	1797	2.2(10)-185	Escherichia coli	b1680	[pn:hypothetical protein]
CONTIG404	2152177_c3_122	1508	7170	609	203	740	2.2(10)-73	Escherichia coli	b1678	[pn:hypothetical protein]
CONTIG405	33886088_fl_1	1509	7171	495	165	90	0.034	Bacillus subtilis	ybf1	[pn:hypothetical protein]
CONTIG405	11050187_fl_2	1510	7172	990	330	146	7.4(10)-10	Escherichia coli	b2578	[pn:hypothetical 21.2 kd protein in srmb-ung intergenic region]
CONTIG405	23833393_fl_12	1511	7173	810	270	341	4.4(10)-31	Escherichia coli	X02307	or:escherichia coli le:115 re:522 di:complement nt:urf 4
CONTIG405	36414562_fl_21	1512	7174	537	179	456	2.7(10)-43	Escherichia coli	b4144	[pn:hypothetical protein] [gn:yjei]
CONTIG405	6522338_fl_23	1513	7175	843	281	966	2.6(10)-97	Escherichia coli	b4147	[pn:elongation factor p] [gn:efp]
CONTIG405	4346033_f2_33	1514	7176	1221	407	247	4.0(10)-21	Escherichia coli	X02307	or:escherichia coli le:2104 re:2802 di:complement nt:urf 3
CONTIG405	23571887_f2_37	1515	7177	555	185	376	8.5(10)-35	Escherichia coli	b4140	[pn:hypothetical protein]
CONTIG405	12141687_f2_41	1516	7178	390	130	449	1.6(10)-42	Escherichia coli	b4142	[pn:groes protein] [gn:mopb]
CONTIG405	21523957_f3_51	1517	7179	219	73	104	2.2(10)-5	Bacillus subtilis	ybf1	[pn:hypothetical protein]
CONTIG405	24798443_f3_64	1518	7180	1659	553	2236	6.7(10)-232	Escherichia coli	b4143	[pn:60 kd chaperonin] [gn:mopa]
CONTIG405	13104628_f3_71	1519	7181	333	111	338	9.0(10)-31	Escherichia coli	b4148	[pn:suges] [gn:suge]

CONTIG405	3017827_c1_81	1520	7182	1248	416	1588	3.1(10)-163	Escherichia coli	b4141	[pn:hypothetical 44.8 kd protein in aspa-mopb intergenic region] [gn:yjeh]
CONTIG405	4353830_c1_83	1521	7183	1518	506	2253	1.1(10)-233	Escherichia coli	b4139	[pn:aspartate ammonia-lyase] [gn:aspa]
CONTIG405	2037566_c1_87	1522	7184	1722	574	2053	1.7(10)-212	Escherichia coli	b4136	[pn:thiol] [gn:dsbd]
CONTIG405	24005301_c2_98	1523	7185	1041	347	1532	2.7(10)-157	Escherichia coli	b4146	[pn:hypothetical 38.7 kd protein in mopa-efp intergenic region]
CONTIG405	19688916_c2_101	1524	7186	627	209	601	1.2(10)-58	Escherichia coli	b4329	[pn:hypothetical 16.2 kd protein in iada-mcrd intergenic region]
CONTIG405	6511528_c2_102	1525	7187	1674	558	101	0.00259	Mycobacterium tuberculosis	M15467	or:mycobacterium tuberculosis pn:unknown protein le:1194 re:1670 di:complement sr:mycobacterium tuberculosis (strain erdman) dna nt:orf d158; putative
CONTIG405	2986693_c2_109	1526	7188	336	112	426	4.2(10)-40	Escherichia coli	b4137	[pn:c-type cy] [gn:cuta]
CONTIG405	30080456_c2_110	1527	7189	681	227	916	5.0(10)-92	Escherichia coli	b4135	[pn:yjdc]
CONTIG405	26777213_c3_113	1528	7190	966	322	703	1.8(10)-69	Escherichia coli	b4149	[pn:hypothetical 19.9 kd protein in suge-ampc intergenic region]
CONTIG405	16620215_c3_115	1529	7191	639	213	118	1.3(10)-5	Escherichia coli	b1951	[pn:colanic acid capsular biosynthesis activation protein a] [gn:rcsa]
CONTIG405	6291406_c3_118	1530	7192	711	237	762	1.1(10)-75	Escherichia coli	b4330	[pn:hypothetical 24.2 kd protein in iada-mcrd intergenic region]
CONTIG405	10317165_c3_119	1531	7193	1191	397	1216	8.3(10)-124	Escherichia coli	b4328	[pn:isoaspartyl dipeptidase] [gn:iada]
CONTIG405	134683_c3_123	1532	7194	474	158	233	1.2(10)-19	Mycobacterium tuberculosis	M15467	or:mycobacterium tuberculosis pn:unknown protein le:242 re:769 di:complement sr:mycobacterium tuberculosis (strain erdman) dna nt:orf f175; putative
CONTIG405	7214683_c3_129	1533	7195	1317	439	1127	2.2(10)-114	Escherichia coli	b4138	[pn:anaerobic c4-dicarboxylate transporter dcua] [gn:dcua]
CONTIG406	15673342_f1_9	1534	7196	1734	578	2055	1.0(10)-212	Escherichia coli	b1336	[pn:hypothetical protein in ogt 5' region] [gn:ydah]

CONTIG406	5100443_f1_10	1535	7197	525	175	717	6.2(10)-71	Escherichia coli	b1335	[pn:o6-methylguanine-dna-alkyltransferase] [gn:ogt]
CONTIG406	24103168_f1_11	1536	7198	768	256	1255	6.0(10)-128	Escherichia coli	b1334	[pn:fumarate and nitrate reduction regulatory protein] [gn:fnr]
CONTIG406	26613250_f2_21	1537	7199	1470	490	1998	1.1(10)-206	Escherichia coli	b1337	[pn:hypothetical protein]
CONTIG406	25942627_f3_28	1538	7200	1719	573	1220	3.1(10)-124	Escherichia coli	b4355	[pn:methyl-accepting chemotaxis protein i] [gn:tsr]
CONTIG406	22345917_f3_29	1539	7201	1347	449	1668	1.0(10)-171	Escherichia coli	b1338	[pn:hypothetical protein] [gn:ydaaj]
CONTIG406	16503_f3_36	1540	7202	999	333	1547	7.0(10)-159	Escherichia coli	b1333	[pn:35.6 kd protein in tpx-fnr intergenic region] [gn:ydaa]
CONTIG406	22073937_c2_63	1541	7203	1296	432	1811	7.4(10)-187	Escherichia coli	b1602	[pn:pyridine nucleotide transhydrogenase subunit- beta] [gn:pnrb]
CONTIG406	4339527_c2_82	1542	7204	897	299	1093	9.0(10)-111	Escherichia coli	b1406	[pn:hypothetical 30.7 kd protein near the replication terminus] [gn:ydbcl]
CONTIG406	23677087_c3_83	1543	7205	1230	410	1253	9.9(10)-128	Salmonella typhimurium	U94729	[de:salmonella typhimurium oxd-6 operon, putative substrate-binding protein (oxd-6a), putative transmembrane protein (oxd-6), putative transmembrane protein (oxd-6c), putative atpase (oxd-6d), and putative atpase (oxd-6e)]
CONTIG406	3963432_c3_92	1544	7206	927	309	1201	3.2(10)-122	Escherichia coli	b1339	[pn:hypothetical protein]
CONTIG406	29328408_c3_93	1545	7207	651	217	658	1.1(10)-64	Escherichia coli	b1340	[pn:hypothetical protein]
CONTIG407	119052_f1_4	1546	7208	1410	470	971	7.5(10)-98	Bacillus subtilis	sacA	[pn:sucrase-6-phosphate hydrolase] [gn:ipa-50d]
CONTIG407	9978192_f1_6	1547	7209	1602	534	552	1.8(10)-53	Escherichia coli	b1421	[pn:methyl-accepting chemotaxis protein iii] [gn:trg]
CONTIG407	9817713_f2_19	1548	7210	873	291	1222	1.8(10)-124	Klebsiella pneumoniae	P27218	sucrose porin precursor.
CONTIG407	3237531_f2_20	1549	7211	1437	479	918	3.1(10)-92	Bacillus subtilis	sacP	[pn:phosphotransferase system] [gn:ipa-49d]
CONTIG407	34422278_f2_23	1550	7212	1023	341	1420	2.0(10)-145	Klebsiella pneumoniae	P37076	sucrose (scr) operon repressor.

CONTIG407	33797292_f2_25	1551	7213	837	279	688	7.4(10)-68	Escherichia coli	b0604	[pn: hypothetical protein]
CONTIG407	34192591_f3_42	1552	7214	591	197	114	1.3(10)-5	Bacillus subtilis	sigZ	[pn: rna polymerase ecf-type sigma factor]
CONTIG407	4188967_f3_47	1553	7215	894	298	418	3.0(10)-39	Bacillus subtilis	alsR	[pn: transcriptional regulator]
CONTIG407	962827_c1_56	1554	7216	1548	516	294	2.2(10)-24	Methanobacterium thermoautotrophicum	MTH1394	[pn: conserved protein]
CONTIG407	31808341_c2_72	1555	7217	1107	369	136	8.5(10)-7	Bacillus subtilis	ymP	[pn: hypothetical protein]
CONTIG407	22657666_c3_90	1556	7218	369	123	332	3.8(10)-30	Bacillus subtilis	yxD	[pn: hypothetical protein]
CONTIG407	14553381_c3_92	1557	7219	936	312	276	3.3(10)-24	Escherichia coli	b0076	[pn: leuO]
CONTIG407	16253767_c3_95	1558	7220	1434	478	1768	2.7(10)-182	Escherichia coli	b1491	[pn: hypothetical protein]
CONTIG407	1457626_c3_100	1559	7221	318	106	96	0.00067	Enterobacter aerogenes	U67194	or: enterobacter aerogenes pn: traec gn: traec le: 44196 re: 46259 di: complement
CONTIG408	34398586_f1_14	1560	7222	609	203	655	2.2(10)-64	Escherichia coli	b1806	[pn: hypothetical protein]
CONTIG408	6120461_f2_20	1561	7223	1572	524	2053	1.7(10)-212	Escherichia coli	b1816	[pn: hypothetical protein] [gn: yoae]
CONTIG408	665793_f2_34	1562	7224	1206	402	1650	8.5(10)-170	Escherichia coli	b1804	[pn: ribonuclease d] [gn: rmd]
CONTIG408	14464211_f3_46	1563	7225	195	65	261	1.3(10)-22	Escherichia coli	b1811	[pn: hypothetical protein]
CONTIG408	6364088_f3_48	1564	7226	1920	640	2813	4.9(10)-293	Escherichia coli	b1808	[pn: hypothetical protein]
CONTIG408	15660326_f3_49	1565	7227	723	241	931	1.3(10)-93	Escherichia coli	b1807	[pn: hypothetical protein]
CONTIG408	23485885_f3_51	1566	7228	1755	585	2749	2.8(10)-286	Escherichia coli	b1805	[pn: long-chain-fatty-acid-coa ligase] [gn: fadd]
CONTIG408	915878_c1_53	1567	7229	270	90	422	1.1(10)-39	Escherichia coli	b1174	[pn: cell division topological specificity factor] [gn: mine]
CONTIG408	4416061_c1_74	1568	7230	966	322	1392	1.8(10)-142	Escherichia coli	b1817	[pn: pts system, mannose-specific iiab component] [gn: manx]



CONTIG408	20789142_c2_89	1569	7231	615	205	836	1.5(10)-83	Escherichia coli	b1813	[pn:hypothetical 21.4 kd protein in pabb-sdaa intergenic region] [gn:yeab]
CONTIG408	26844406_c3_105	1570	7232	504	168	548	5.0(10)-53	Escherichia coli	b1809	[pn:hypothetical protein]
CONTIG408	16265877_c3_106	1571	7233	1380	460	1671	5.0(10)-172	Escherichia coli	b1812	[pn:para-aminobenzoate synthase component i] [gn:pabb]
CONTIG408	29845761_c3_108	1572	7234	1407	469	2040	4.0(10)-211	Escherichia coli	b1814	[pn:l-serine dehydratase 1] [gn:sdad]
CONTIG408	9784408_c3_109	1573	7235	1728	576	1732	1.7(10)-178	Escherichia coli	b1815	[pn:hypothetical protein]
CONTIG409	23714651_f1_16	1574	7236	573	191	95	0.03799	Thermus thermophilus	Y15464	[de:thermus thermophilus phes, phet genes and 5 orf"s.] [pn:ferredoxin oxidoreductase-like (gamma and alpha) [gn:orf2]
CONTIG409	26352002_f1_17	1575	7237	792	264	245	1.2(10)-20	Escherichia coli	b3442	[pn:hypothetical 44.2 kd protein in gnt-r-ggt intergenic region] [gn:yhhz]
CONTIG409	406255_f2_27	1576	7238	495	165	293	5.2(10)-26	Escherichia coli	b1183	[pn:umud protein] [gn:umud]
CONTIG409	14885165_f3_46	1577	7239	297	99	390	2.7(10)-36	Escherichia coli	D93826	hypothetical 11k protein (insertion sequence is1) -escherichia coli this protein is coded by the insertion sequence is1.
CONTIG409	16486527_f3_48	1578	7240	1272	424	959	1.3(10)-96	Escherichia coli	b1184	[pn:umuc protein] [gn:umuc]
CONTIG409	33460927_f3_53	1579	7241	570	190	194	1.6(10)-15	Escherichia coli	Z68186	or:escherichia coli pn:bfp gn:bfp le:9044 re:9490 di:direct
CONTIG409	14578162_f3_54	1580	7242	228	76	181	1.7(10)-13	Escherichia coli	b3442	[pn:hypothetical 44.2 kd protein in gnt-r-ggt intergenic region] [gn:yhhz]
CONTIG409	4322963_f3_55	1581	7243	516	172	135	2.8(10)-9	Escherichia coli	b3443	[pn:hypothetical protein] [gn:yrra]
CONTIG409	24410340_c1_63	1582	7244	1500	500	214	5.0(10)-15	Plasmid F	M24492	or:plasmid f pn:periplasmic protein gn:trah le:550 re:1629 di:direct sr:plasmid f dna
CONTIG409	6735393_c1_72	1583	7245	285	95	446	3.2(10)-42	Escherichia coli	b4294	[pn:insertion element is1f protein insa] [gn:insa_7]
CONTIG409	15897332_c1_75	1584	7246	1959	653	115	0.00309	Lycopersicon esculentum	Y07861	or:lycopersicon esculentum pn:mfp1 protein gn:mfp1 le:6 re:2099 di:direct sr:tomato

CONTIG409	2047880_c1_76	1585	7247	315	105	369	4.7(10)-34	Yersinia pestis AF053945	[de:yersinia pestis plasmid ppcpl, complete plasmid sequence.] [pn:transposase]
CONTIG409	4901068_c2_80	1586	7248	759	253	347	1.0(10)-31	Escherichia coli	[pn:disulfide interchange protein, precursor] [gn:dsbc]
CONTIG409	34178880_c2_83	1587	7249	909	303	238	3.6(10)-20	Coxiella burnetii	outer membrane protein - coxiella burnetii
CONTIG409	2911290_c2_86	1588	7250	234	78	296	2.6(10)-26	Escherichia coli	hypothetical protein - escherichia coli
CONTIG409	25480326_c3_91	1589	7251	3963	1321	111	0.032	Bombyx mori	sericin lb - silkworm
CONTIG409	25567885_c3_96	1590	7252	525	175	898	4.0(10)-90	Escherichia coli	[pn:insertion element isl protein insb] [gn:insb_2]
CONTIG41	14492812_fl_1	1591	7253	453	151	410	2.1(10)-38	Escherichia coli	[pn:hypothetical 16.0 kd protein in lpxc-seca intergenic region] [gn:yaca]
CONTIG41	26688902_f3_3	1592	7254	213	71	278	2.1(10)-24	Escherichia coli	[pn:udp-3-o-3-hydroxymyristoyl n-acetylglucosamine deacetylase] [gn:lpxc]
CONTIG410	26593890_fl_19	1593	7255	1506	502	523	8.0(10)-58	Bacillus subtilis	[pn:hypothetical protein]
CONTIG410	20587883_f2_38	1594	7256	462	154	293	5.2(10)-26	Erwinia amylovora	or:erwinia amylovora le:10466 re:10885 di:direct nt:orf 12
CONTIG410	26449211_f3_46	1595	7257	963	321	218	5.7(10)-18	Escherichia coli	[pn:asc operon repressor protein] [gn:ascg]
CONTIG410	34070927_c1_67	1596	7258	672	224	155	2.2(10)-11	Escherichia coli	[pn:hypothetical protein] [gn:ycdc]
CONTIG410	24423567_c1_72	1597	7259	3141	1047	2558	5.0(10)-266	Escherichia coli	[pn:acriflavin resistance protein b] [gn:acrb]
CONTIG410	16828927_c1_73	1598	7260	1164	388	433	7.7(10)-41	Bacillus subtilis	[pn:hypothetical protein]
CONTIG410	10948818_c1_76	1599	7261	327	109	182	3.1(10)-14	Bacillus subtilis	[pn:hypothetical protein]
CONTIG410	34416581_c2_80	1600	7262	576	192	146	2.0(10)-10	Escherichia coli	[pn:glucuronide repressor] [gn:uidr]
CONTIG410	3396086_c2_81	1601	7263	1155	385	831	5.2(10)-83	Escherichia coli	[pn:acriflavin resistance protein a precursor] [gn:acra]
CONTIG410	12692683_c3_105	1602	7264	1386	462	744	8.5(10)-74	Escherichia coli	[pn:hypothetical protein] [gn:yicb]

CONTIG410	3016517_c3_113	1603	7265	1284	428	539	4.5(10)-52	Bacillus subtilis	ywbA	[pn:hypothetical protein] [gn:ipa-16d]
CONTIG411	24650302_fl_1	1604	7266	1062	354	1374	1.5(10)-140	Escherichia coli	b0929	[pn:outer membrane protein f precursor] [gn:ompf]
CONTIG411	22520135_fl_15	1605	7267	666	222	648	1.3(10)-63	Escherichia coli	b0920	[pn:hypothetical protein in kdsb-kicb intergenic region] [gn:ycbc]
CONTIG411	24851430_f2_17	1606	7268	1209	403	1930	1.8(10)-199	Escherichia coli	b0928	[pn:aspartate aminotransferase] [gn:aspc]
CONTIG411	35814061_c1_59	1607	7269	816	272	1089	2.3(10)-110	Escherichia coli	b0921	[pn:hypothetical 29.8 kd protein in kdsb-kicb intergenic region] [gn:smta]
CONTIG411	25665880_c1_60	1608	7270	1326	442	2147	1.8(10)-222	Escherichia coli	b0922	[pn:mukf protein] [gn:mukf]
CONTIG411	34274216_c1_61	1609	7271	4464	1488	6395	0	Escherichia coli	b0924	[pn:mukb]
CONTIG411	15686_c1_65	1610	7272	582	194	820	7.5(10)-82	Escherichia coli	b0926	[pn:hypothetical protein] [gn:ycbk]
CONTIG411	30730282_c2_75	1611	7273	801	267	1108	2.2(10)-112	Escherichia coli	b0923	[pn:muke protein] [gn:muke]
CONTIG411	13088555_c2_76	1612	7274	225	75	93	8.3(10)-5	Homo sapiens	S40543	or:homo sapiens pn:low density lipoprotein receptor lc:1 re:177 di:direct sr:human nt:description: low density lipoprotein receptor, ldl cell division protein mukb.
CONTIG411	33839465_c2_79	1613	7275	330	110	136	9.3(10)-8	Escherichia coli	P22523	
CONTIG411	16542168_c3_89	1614	7276	1908	636	2340	6.5(10)-243	Escherichia coli	b0925	[pn:hypothetical protein in mukb 3''' region] [gn:ycbb]
CONTIG411	35683285_c3_90	1615	7277	693	231	981	6.5(10)-99	Escherichia coli	b0927	[pn:hypothetical protein] [gn:ycbl]
CONTIG411	1442568_c3_91	1616	7278	1218	406	393	1.3(10)-36	Pseudomonas aeruginosa	U50396	or:pseudomonas aeruginosa pn:wbpn gn:wbpn lc:22302 re:23693 di:direct
CONTIG412	13087788_fl_13	1617	7279	1392	464	543	1.7(10)-52	Bacillus subtilis	ycnF	[pn:hypothetical protein]
CONTIG412	31344415_fl_14	1618	7280	1586	529	1902	1.7(10)-196	Escherichia coli	b4340	[pn:hypothetical 53.0 kd protein in iada-mcrd intergenic region]
CONTIG412	24431531_f2_15	1619	7281	1275	425	1574	9.5(10)-162	Escherichia coli	b4336	[pn:hypothetical 48.2 kd protein in iada-mcrd intergenic region]
CONTIG412	4957168_f3_36	1620	7282	528	176	183	2.3(10)-14	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr lc:59 re:580 di:direct nt:orf2

CONTIG412	31431505_f3_38	1621	7283	906	302	510	5.4(10)-49	Escherichia coli	b0900	[pn:hypothetical protein] [gn:ycan]
CONTIG412	26432281_c1_50	1622	7284	1050	350	276	3.3(10)-24	Bacillus subtilis	ydfG	[pn:hypothetical protein]
CONTIG412	13783340_c1_57	1623	7285	495	165	208	5.4(10)-17	Escherichia coli	b1642	[pn:hypothetical protein] [gn:slya]
CONTIG412	5119091_c2_74	1624	7286	744	248	373	1.8(10)-34	Bacillus subtilis	ydfF	[pn:hypothetical protein]
CONTIG412	22895156_c3_78	1625	7287	189	63	161	5.2(10)-12	Escherichia coli	b4341	[pn:hypothetical protein] [gn:yjis]
CONTIG412	3407211_c3_84	1626	7288	1449	483	687	9.4(10)-68	Escherichia coli	b0312	[pn:betaine aldehyde dehydrogenase] [gn:betb]
CONTIG412	2081292_c3_85	1627	7289	399	133	96	4.0(10)-5	Bacillus subtilis	yqjZ	[pn:hypothetical protein]
CONTIG412	4883288_c3_87	1628	7290	1077	359	267	3.0(10)-23	Escherichia coli	b4082	[pn:hypothetical 36.9 kd protein in fdhf-phnp intergenic region]
CONTIG413	22370937_f1_4	1629	7291	948	316	1266	4.2(10)-129	Escherichia coli	b2131	[pn:hypothetical 32.6 kd protein in molr-bglx intergenic region] [gn:yehz]
CONTIG413	16986091_f1_5	1630	7292	1170	390	662	4.2(10)-65	Escherichia coli	b2130	[pn:hypothetical abc transporter permease protein yehy] [gn:yehy]
CONTIG413	17058580_f2_17	1631	7293	1104	368	1257	3.7(10)-128	Escherichia coli	b2129	[pn:hypothetical abc transporter in molr-bglx intergenic region] [gn:yehx]
CONTIG413	3227158_f2_24	1632	7294	771	257	1035	1.3(10)-104	Escherichia coli	b2125	[pn:hypothetical 27.9 kd protein in molr-bglx intergenic region]
CONTIG413	14255177_f3_28	1633	7295	285	95	441	1.1(10)-41	Escherichia coli	b2134	[pn:penicillin-binding protein 7 precursor] [gn:pbgp]
CONTIG413	7223781_f3_34	1634	7296	2391	797	3750	0	Escherichia coli	b2132	[pn:periplasmic beta-glucosidase precursor] [gn:bglx]
CONTIG413	30666041_f3_39	1635	7297	741	247	821	6.0(10)-82	Escherichia coli	b2128	[pn:hypothetical 25.5 kd protein in molr-bglx intergenic region] [gn:yehw]
CONTIG413	1431637_f3_40	1636	7298	1704	568	2269	2.2(10)-235	Escherichia coli	b2126	[pn:hypothetical 62.1 kd protein in molr-bglx intergenic region] [gn:yehu]
CONTIG413	14642543_f3_42	1637	7299	525	175	642	5.5(10)-63	Escherichia coli	b2124	[pn:hypothetical 18.0 kd protein in molr-bglx intergenic region] [gn:yehs]

CONTIG413	21565658_c1_45	1638	7300	1182	394	1740	2.5(10)-179	Escherichia coli	b2114	[pn:methionyl-trna synthetase] [gn:metg]
CONTIG413	24397250_c1_48	1639	7301	825	275	890	2.8(10)-89	Escherichia coli	b2127	[pn:hypothetical transcriptional regulator in molr-bglx intergenic region] [gn:yehv]
CONTIG413	16620831_c2_85	1640	7302	1923	641	2678	9.8(10)-279	Escherichia coli	b2133	[pn:d-lactate dehydrogenase] [gn:dlid]
CONTIG413	23847008_c2_86	1641	7303	636	212	488	1.2(10)-46	Alcaligenes faecalis	P31668	phosphinothricin-resistance protein (ptc-resistance protein).
CONTIG414	25829787_fl_1	1642	7304	900	300	1364	1.7(10)-139	Escherichia coli	b4161	[pn:hypothetical 37.7 kd protein in psd-amib intergenic region]
CONTIG414	23567037_fl_9	1643	7305	417	139	472	5.7(10)-45	Escherichia coli	b4152	[pn:fumarate reductase, membrane anchor polypeptide] [gn:frdc]
CONTIG414	3162711_fl_12	1644	7306	1254	418	1245	7.0(10)-127	Escherichia coli	b4150	[pn:beta-lactamase precursor] [gn:ampe]
CONTIG414	2439512_f2_13	1645	7307	1044	348	1600	1.7(10)-164	Escherichia coli	b4160	[pn:phosphatidylserine decarboxylase proenzyme] [gn:psd]
CONTIG414	33206957_f2_21	1646	7308	1842	614	2810	1.0(10)-292	Escherichia coli	b4154	[pn:fumarate reductase flavoprotein subunit] [gn:frda]
CONTIG414	10948336_f2_22	1647	7309	417	139	535	1.2(10)-51	Escherichia coli	b4151	[pn:fumarate reductase, membrane anchor polypeptide] [gn:frdd]
CONTIG414	16879681_f3_26	1648	7310	3483	1161	4253	0	Escherichia coli	b4159	[pn:hypothetical 123.8 kd protein in genx-psd intergenic region]
CONTIG414	32131937_f3_35	1649	7311	810	270	1239	3.0(10)-126	Escherichia coli	b4153	[pn:fumarate reductase iron-sulfur protein] [gn:frdb]
CONTIG414	37576_c1_41	1650	7312	903	301	1516	1.3(10)-155	Enterobacter cloacae	A25686	amp <sup>r</sup> protein - enterobacter cloacae
CONTIG414	26056526_c1_48	1651	7313	981	327	1573	1.2(10)-161	Escherichia coli	b4155	[pn:lysyl-trna synthetase analog] [gn:yjea]
CONTIG414	163931_c2_66	1652	7314	1563	521	1981	7.0(10)-205	Escherichia coli	b4156	[pn:hypothetical 56.3 kd protein in genx-psd intergenic region]
CONTIG415	23572162_fl_1	1653	7315	642	214	883	1.6(10)-88	Serratia marcescens	U59131	or:serratia marcescens le:1362 re:1982 di:direct nt:orf
CONTIG415	4876425_fl_10	1654	7316	306	102	418	3.0(10)-39	Escherichia coli	b2105	[pn:hypothetical protein] [gn:yohl]
CONTIG415	32230063_fl_19	1655	7317	1311	437	172	3.1(10)-9	Saccharomyces cerevisiae	YDL058 W	[pn:intracellular protein transport protein] [gn:usol]
CONTIG415	23928775_f2_26	1656	7318	765	255	1195	1.3(10)-121	Serratia marcescens	U59131	or:serratia marcescens pn:stba gn:stba le:313 re:1350 di:direct

CONTIG415	19572828_f2_27	1657	7319	1416	472	2236	6.7(10)-232	Escherichia coli	S70165	,
CONTIG415	6297128_f2_34	1658	7320	825	275	102	0.016	Serratia marcescens	U60283	or:serratia marcescens pn:restriction methylase gn:trag l le:84 re:1775 di:direct nt:putative restriction methylase
CONTIG415	24744778_f2_39	1659	7321	447	149	237	4.5(10)-20	Escherichia coli	P09183	very hypothetical 20.3 kd protein in dcm 3'region (orf 3).
CONTIG415	6439528_f3_43	1660	7322	255	85	358	6.9(10)-33	Escherichia coli	I77547	hypothetical protein 2 (insertion sequence is903) -escherichia coli
CONTIG415	20984450_f3_49	1661	7323	468	156	618	1.8(10)-60	Escherichia coli	S70162	,
CONTIG415	4875376_f3_58	1662	7324	585	195	265	4.9(10)-23	Haemophilus influenzae	HI1296	[pn:partitioning system protein] [gn:parb]
CONTIG415	21726630_c1_67	1663	7325	1536	512	802	6.2(10)-80	Rhizobium sp.	P55501	hypothetical 57.2 kd protein y4javy4neVy4se,
CONTIG415	31532840_c2_86	1664	7326	1101	367	787	2.3(10)-78	Yersinia pestis	AF053947	[de:yersinia pestis plasmid pmt1, complete plasmid sequence.] [pn:transposase]
CONTIG415	14704677_c2_87	1665	7327	990	330	1597	3.5(10)-164	Escherichia coli	X02527	or:escherichia coli le:199 re:1122 di:direct nt:orf1 (aa 1-307)
CONTIG415	33992155_c3_92	1666	7328	1482	494	1385	1.0(10)-141	Escherichia coli	b1961	[pn:dna-cytosine methyltransferase] [gn:dcn]
CONTIG415	15750433_c3_99	1667	7329	786	262	474	3.5(10)-45	Rhizobium sp.	P55500	putative insertion sequence atp-binding protein y4iqVy4ndVy4sd,
CONTIG415	5117806_c3_101	1668	7330	1131	377	498	3.8(10)-93	Escherichia coli	b2106	[pn:hypothetical protein]
CONTIG416	16260012_f1_8	1669	7331	1026	342	1058	4.5(10)-107	Escherichia coli	b3196	[pn:hypothetical 34.7 kd protein in murz-rpon intergenic region] [gn:yrbg]
CONTIG416	32547880_f1_9	1670	7332	606	202	869	4.9(10)-87	Escherichia coli	b3198	[pn:hypothetical 20.0 kd protein in murz-rpon intergenic region] [gn:yrbj]
CONTIG416	1445325_f1_10	1671	7333	576	192	655	2.2(10)-64	Escherichia coli	b3200	[pn:17.3 kd protein in rpon 5''' region precursor] [gn:yhbh]
CONTIG416	16828525_f1_11	1672	7334	732	244	1175	1.8(10)-119	Escherichia coli	b3201	[pn:probable abc transporter in ntra/rpon 5''' region] [gn:yhbq]
CONTIG416	29928931_f1_12	1673	7335	417	139	339	7.0(10)-31	Escherichia coli	b3203	[pn:probable sigma] [gn:yhbh]

CONTIG416	35556932_f1_13	1674	7336	372	124	438	2.2(10)-41	Escherichia coli	b3206	[pn:phosphocarrier protein npr] [gn:ptso]
CONTIG416	13127067_f2_22	1675	7337	891	297	1242	1.5(10)-126	Escherichia coli	b3187	[pn:octaprenyl pyro] [gn:ispb]
CONTIG416	17540_f2_23	1676	7338	342	114	389	3.6(10)-36	Escherichia coli	b3188	[pn:ner-like protein] [gn:nlp]
CONTIG416	212966_f2_31	1677	7339	1062	354	1444	5.7(10)-148	Escherichia coli	b3197	[pn:hypothetical 35.2 kd protein in murz-rpon intergenic region] [gn:yrbh]
CONTIG416	25678166_f2_36	1678	7340	507	169	733	9.5(10)-75	Escherichia coli	b3204	[pn:enzyme iiantr] [gn:ptsn]
CONTIG416	36524205_f2_37	1679	7341	900	300	1357	9.5(10)-139	Escherichia coli	b3205	[pn:hypothetical protein] [gn:yhbj]
CONTIG416	17070418_f3_58	1680	7342	594	198	770	1.5(10)-76	Escherichia coli	b3199	[pn:hypothetical 21.7 kd protein in murz-rpon intergenic region] [gn:yrbk]
CONTIG416	21756342_f3_60	1681	7343	1473	491	1867	8.5(10)-193	Escherichia coli	b3202	[pn:sigma-n] [gn:rpon]
CONTIG416	34395429_c1_70	1682	7344	833	277	1192	2.8(10)-121	Escherichia coli	b4262	[pn:hypothetical 39.8 kd protein in pepa-gntv intergenic region] [gn:yjgq]
CONTIG416	24412837_c1_73	1683	7345	501	167	617	2.5(10)-60	Escherichia coli	b3210	[pn:aerobic respiration control sensor protein arcb] [gn:arcb]
CONTIG416	35438562_c1_74	1684	7346	759	253	917	4.0(10)-92	Escherichia coli	b3208	[pn:hypothetical 27.3 kd protein in ptso-arcb intergenic region] [gn:yrbm]
CONTIG416	13078568_c1_89	1685	7347	798	266	1236	6.2(10)-126	Escherichia coli	b3194	[pn:hypothetical 27.9 kd protein in murz-rpon intergenic region] [gn:yrbf]
CONTIG416	35750840_c1_90	1686	7348	357	119	371	2.8(10)-34	Escherichia coli	b3191	[pn:hypothetical 14.4 kd protein in murz-rpon intergenic region] [gn:yrbf]
CONTIG416	20875000_c1_91	1687	7349	381	127	412	1.3(10)-38	Escherichia coli	b3190	[pn:hypothetical 9.5 kd protein in murz-rpon intergenic region] [gn:yrbf]
CONTIG416	7930_c2_96	1688	7350	702	234	792	7.0(10)-79	Escherichia coli	b3209	[pn:sigma cross-reacting protein 27a] [gn:yhbl]
CONTIG416	4978140_c2_108	1689	7351	567	189	610	1.3(10)-59	Escherichia coli	b3193	[pn:hypothetical protein] [gn:yrbf]

CONTIG416	24823250_c2_109	1690	7352	654	218	1004	2.3(10)-101	Escherichia coli	b3192	[pn:hypothetical 24.0 kd protein in murz-rpon intergenic region] [gn:yrbcb]
CONTIG416	34381303_c3_112	1691	7353	1977	659	2853	2.7(10)-297	Escherichia coli	b3210	[pn:aerobic respiration control sensor protein archb] [gn:archb]
CONTIG416	14630327_c3_127	1692	7354	813	271	1182	3.2(10)-120	Escherichia coli	b3195	[pn:hypothetical protein] [gn:yrbfb]
CONTIG416	16211468_c3_132	1693	7355	1305	435	1930	1.8(10)-199	Escherichia coli	b3189	[pn:udp-n-acetylglucosamine 1-carboxyvinyltransferase] [gn:mura]
CONTIG417	33376713_fl_3	1694	7356	1020	340	108	1.2(10)-5	coliphage T4	P39506	hypothetical 9.5 kd protein in frd-gp32 intergenic region.
CONTIG417	10976425_f2_30	1695	7357	201	67	94	6.5(10)-5	Escherichia coli	b1565	[pn:hypothetical protein]
CONTIG417	21775383_f2_45	1696	7358	300	100	505	1.8(10)-48	Bacteriophage lambda	J01735	or:bacteriophage lambda le:S25 re:851 di:direct sr:bacteriophage lambda kh100 is5 element nt:small gene
CONTIG417	21877261_f2_46	1697	7359	252	84	327	1.3(10)-29	Escherichia coli	b1371	[pn:hypothetical protein]
CONTIG417	33691012_c1_72	1698	7360	855	285	508	8.8(10)-49	Rhizobium sp.	P55373	putative transposase y4bf,
CONTIG417	4299217_c2_88	1699	7361	642	214	903	1.2(10)-90	Enterobacter agglomerans	B38965	hypothetical protein b (insertion sequence is1222) -enterobacter agglomerans
CONTIG417	2118828_c2_89	1700	7362	1053	351	301	4.2(10)-26	Mycobacterium leprae	Z97369	[PN:hypothetical protein MLCB250.18c] [GN:MLCB250.18c] [DE:Mycobacterium leprae cosmid B250.] [NT:MLCB250.18c, unknown, len: 596 aa; highly similar] [LE:5885] [RE:7675] [DI:complement]
CONTIG417	30204130_c2_90	1701	7363	279	93	109	9.8(10)-6	Escherichia coli	U95365	transposase, is5b,
CONTIG417	34629052_c2_96	1702	7364	690	230	236	5.7(10)-20	Escherichia coli	b1567	[pn:hypothetical protein]
CONTIG417	160080_c2_97	1703	7365	948	316	747	4.0(10)-74	Synechocystis sp.	P74068	[GN:SLL1263] [SR:PCC 6803.] [DE:HYPOTHETICAL 33.3 KD PROTEIN SLL1263] [SP:P74068]



CONTIG417	16837632_c2_98	1704	7366	294	98	473	4.5(10)-45	Escherichia coli	b1563	[pn:hypothetical rele protein] [gn:rele]
CONTIG417	4339135_c3_104	1705	7367	1026	342	1767	3.3(10)-182	Escherichia coli	b1994	[pn:insertion element is5 hypothetical 39.3 kd protein]
CONTIG417	25830_c3_115	1706	7368	1374	458	278	1.0(10)-22	Streptomyces coelicolor	P14707	mini-circle hypothetical 45.7 kd protein.
CONTIG417	33494038_c3_116	1707	7369	243	81	387	5.7(10)-36	Escherichia coli	b1564	[pn:relb protein] [gn:relb]
CONTIG417	628161_c3_117	1708	7370	189	63	127	2.1(10)-8	Escherichia coli	P23587	fmc protein homolog.
CONTIG418	35314451_f1_2	1709	7371	1212	404	1621	1.0(10)-166	Escherichia coli	b2028	[pn:hypothetical protein] [gn:ugd]
CONTIG418	14635416_f1_7	1710	7372	468	156	194	1.6(10)-15	Escherichia coli	D90840	or:escherichia coli gn:yh1028w le:12646 re:12942 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:orf_id:o350#11; similar to [swissprot accession]
CONTIG418	2776938_f1_15	1711	7373	351	117	102	0.00046	Homo sapiens	AC004493	[de:homo sapiens chromosome 16, cosmid clone 373c8 (lanl), complete sequence.] [pn:kiaa0324] [gn:kiaa0324]
CONTIG418	17051316_f2_18	1712	7374	1143	381	1544	1.3(10)-158	Escherichia coli	b2041	[pn:tdp-glucose 4,6-dehydratase] [gn:rftb]
CONTIG418	32229077_f2_19	1713	7375	882	294	1314	3.3(10)-134	Escherichia coli	b2039	[pn:tdp-glucose pyrophosphorylase] [gn:rfta]
CONTIG418	6057958_f2_31	1714	7376	876	292	1133	5.2(10)-115	Escherichia coli	b2016	[pn:hypothetical protein]
CONTIG418	32553165_f3_32	1715	7377	1449	483	2300	1.1(10)-238	Escherichia coli	b2029	[pn:6-phosphogluconate dehydrogenase, decarboxylating] [gn:gnd]
CONTIG418	15113290_f3_39	1716	7378	1143	381	901	2.0(10)-90	Escherichia coli	b2027	[pn:hypothetical protein]
CONTIG418	31739687_c1_53	1717	7379	984	328	1450	1.3(10)-148	Escherichia coli	b2019	[pn:atp phosphoribosyltransferase] [gn:hig]
CONTIG418	34188291_c1_58	1718	7380	1089	363	1666	1.7(10)-171	Escherichia coli	b2022	[pn:histidinol phosphatase] [gn:hishb]
CONTIG418	32424192_c1_59	1719	7381	933	311	1268	2.6(10)-129	Escherichia coli	b2025	[pn:hisf protein] [gn:hisf]

CONTIG418	31922662_c2_70	1720	7382	1140	380	1608	2.3(10)-165	Escherichia coli	b2021	[pn:imidazole] [gn:hisc]
CONTIG418	32678126_c2_75	1721	7383	816	272	915	6.5(10)-92	Escherichia coli	b2024	[pn:phosphoribosylformimino-5-aminoimidazole carboxamide ribotide] [gn:hisa]
CONTIG418	15734716_c2_77	1722	7384	330	110	97	3.1(10)-5	Salmonella typhimurium	X03976	or:salmonella typhimurium le:l re:>173 di:direct nt:chimeric protein of hisf and hisie genes (57 aa)
CONTIG418	16620952_c2_80	1723	7385	1017	339	1466	2.7(10)-150	Escherichia coli	Q04871	hypothetical 37.6 kd protein in cld 5'region (orf2).
CONTIG418	35361063_c3_88	1724	7386	1326	442	1927	3.7(10)-199	Escherichia coli	b2020	[pn:histidinol dehydrogenase]
CONTIG418	2469541_c3_90	1725	7387	621	207	961	8.6(10)-97	Escherichia coli	b2023	[pn:amidotransferase] [gn:hish]
CONTIG418	6445253_c3_91	1726	7388	654	218	897	5.2(10)-90	Escherichia coli	b2026	[pn:phosphoribosyl-amp cyclohydrolase / phosphoribosyl-atp pyrophosphohydrolase] [gn:hisi]
CONTIG419	4350678_f1_1	1727	7389	531	177	93	0.26	Caenorhabditis elegans	L46861	or:caenorhabditis elegans pn:taln le:2 re:7663 di:direct sr:caenorhabditis elegans (strain bristol) (tissue library: whol
CONTIG419	22558287_f1_9	1728	7390	2559	853	2822	5.4(10)-294	Escherichia coli	b0496	[pn:hypothetical protein] [gn:ybbp]
CONTIG419	16458166_f2_24	1729	7391	735	245	814	3.2(10)-81	Escherichia coli	b0490	[pn:hypothetical protein] [gn:ybbi]
CONTIG419	33847266_f2_32	1730	7392	741	247	959	1.3(10)-96	Escherichia coli	b0495	[pn:hypothetical abc transporter] [gn:ybba]
CONTIG419	24791562_f2_42	1731	7393	642	214	581	1.6(10)-56	Escherichia coli	U82664	or:escherichia coli le:l33380 re:l34066 di:direct nt:hypothetical protein
CONTIG419	21525291_f3_54	1732	7394	864	288	909	2.7(10)-91	Escherichia coli	b0491	[pn:hypothetical protein] [gn:ybbm]
CONTIG419	21644141_f3_63	1733	7395	1167	389	99	0.033	Escherichia coli	b1377	[pn:hypothetical protein]
CONTIG419	24833550_f3_70	1734	7396	1581	527	2192	3.1(10)-227	Escherichia coli	b0526	[pn:cysteiny-l-trna synthetase] [gn:cys]
CONTIG419	2750412_c1_77	1735	7397	1554	518	1039	4.7(10)-105	Bacillus subtilis	ptsG	[pn:phosphotransferase system]

CONTIG419	162701_c1_79	1736	7398	564	188	774	5.7(10)-77	Escherichia coli	b0527	[pn:hypothetical protein in cyss-fold intergenic region] [gn:ybcj]
CONTIG419	31719580_c1_84	1737	7399	645	215	735	7.7(10)-73	Escherichia coli	b0523	[pn:phosphoribosylaminoimidazole carboxylase catalytic subunit] [gn:pure]
CONTIG419	1272192_c1_95	1738	7400	297	99	92	0.00011	Entamoeba histolytica	Y14328	[PN:3E1 protein] [DE:Entamoeba histolytica mRNA for 3E1 protein.] [LE:32] [RE:418] [DI:direct]
CONTIG419	31369027_c1_98	1739	7401	813	271	1203	2.0(10)-122	Escherichia coli	b0493	[pn:hypothetical protein] [gn:ybb0]
CONTIG419	13015666_c1_103	1740	7402	531	177	553	1.5(10)-53	Escherichia coli	b0489	[pn:hypothetical protein] [gn:ybbk]
CONTIG419	33848817_c2_104	1741	7403	342	114	324	2.7(10)-29	Escherichia coli	b0528	[pn:hypothetical 7.4 kd protein in cyss-fold intergenic region] [gn:ybcj]
CONTIG419	34173532_c2_126	1742	7404	876	292	1141	7.2(10)-116	Escherichia coli	b0492	[pn:hypothetical protein] [gn:ybbn]
CONTIG419	11228205_c3_134	1743	7405	249	83	330	6.4(10)-30	Escherichia coli	b0529	[pn:methylenetetrahydrofolate dehydrogenase] [gn:fold]
CONTIG419	36048591_c3_138	1744	7406	1173	391	498	1.0(10)-47	Escherichia coli	b1620	[pn:repressor protein] [gn:mali]
CONTIG419	4004186_c3_142	1745	7407	558	186	829	8.5(10)-83	Escherichia coli	b0525	[pn:peptidyl-prolyl cis-trans isomerase b] [gn:ppib]
CONTIG419	14978525_c3_143	1746	7408	726	242	1045	1.1(10)-105	Escherichia coli	b0524	[pn:hypothetical 26.9 kd protein in pure-ppib intergenic region]
CONTIG419	16102281_c3_144	1747	7409	1089	363	1634	4.2(10)-168	Escherichia coli	b0522	[pn:phosphoribosylaminoimidazole carboxylase apase subunit] [gn:purk]
CONTIG419	16113178_c3_145	1748	7410	1155	385	1365	1.3(10)-139	Escherichia coli	b0503	[pn:hypothetical 41.1 kd protein in rhds-gcl intergenic region] [gn:ybbb]
CONTIG419	24039076_c3_150	1749	7411	888	296	990	7.4(10)-100	Escherichia coli	b0494	[pn:acyl-coa thioesterase i] [gn:tesa]
CONTIG420	14629541_c1_70	1750	7412	630	210	722	1.8(10)-71	Escherichia coli	b2051	[pn:hypothetical 18.4 kd protein in cpsb 5''' region] [gn:yefc]
CONTIG420	886252_c1_73	1751	7413	1422	474	2149	1.1(10)-222	Escherichia coli	b2048	[pn:phosphomannomutase] [gn:manb]
CONTIG420	32682807_c1_74	1752	7414	1497	499	2158	1.2(10)-223	Escherichia coli	b2046	[pn:hypothetical protein]
CONTIG420	17004375_c1_75	1753	7415	1296	432	1937	3.2(10)-200	Escherichia coli	b2045	[pn:hypothetical protein] [gn:weak]

CONTIG420	24427167_c2_79	1754	7416	1035	345	1509	7.4(10)-155	Escherichia coli	b2052	[pn:hypothetical 36.1 kd protein in cpsb 5''' region] [gn:yefb]
CONTIG420	20330425_c2_94	1755	7417	936	312	1389	3.8(10)-142	Escherichia coli	b2042	[pn:utp-glucose-1-phosphate uridylyltransferase] [gn:galf]
CONTIG420	24251592_c3_95	1756	7418	1134	378	1888	5.0(10)-195	Escherichia coli	b2053	[pn:gdp-mannose 4,6-dehydratase] [gn:yefa]
CONTIG420	36417061_c3_98	1757	7419	1275	425	1634	4.2(10)-168	Escherichia coli	b2050	[pn:hypothetical 44.9 kd protein in cpsb 5''' region] [gn:yefd]
CONTIG420	21759652_c3_99	1758	7420	1440	480	2174	2.5(10)-225	Escherichia coli	b2049	[pn:mannose-1-phosphate guanylyltransferase] [gn:manc]
CONTIG420	11800807_c3_103	1759	7421	1467	489	2060	3.0(10)-213	Escherichia coli	b2047	[pn:hypothetical protein] [gn:wcaj]
CONTIG420	31505387_c3_109	1760	7422	1392	464	1689	6.2(10)-174	Escherichia coli	b2044	[pn:hypothetical protein] [gn:wcal]
CONTIG420	3158143_c3_110	1761	7423	1398	466	1857	9.8(10)-192	Escherichia coli	b2043	[pn:hypothetical protein] [gn:wcam]
CONTIG420	30672166_c3_111	1762	7424	1011	337	234	1.7(10)-18	Saccharomyces cerevisiae	YGL001C	[pn:putative 3-beta-hydroxysteroid dehydrogenase]
CONTIG421	16991042_f1_3	1763	7425	438	146	91	0.00839	Saccharomyces cerevisiae	YJR151C	[pn:similarity to mucin proteins, ykl224c, stalp] [gn:j2223]
CONTIG421	33331650_f1_5	1764	7426	696	232	907	4.5(10)-91	Escherichia coli	b0046	[pn:hypothetical nadph oxidoreductase in fixc-kefc intergenic region] [gn:yabf]
CONTIG421	6679631_f2_15	1765	7427	288	96	182	3.1(10)-14	Escherichia coli	b2809	[pn:hypothetical protein]
CONTIG421	31646007_f2_22	1766	7428	1977	659	2127	2.3(10)-220	Escherichia coli	b0047	[pn:glutathione-regulated potassium-efflux system protein kefc] [gn:kefc]
CONTIG421	964590_f2_23	1767	7429	513	171	804	3.7(10)-80	Escherichia coli	b0048	[pn:dihydrofolate reductase type i] [gn:fola]
CONTIG421	20996093_f3_33	1768	7430	834	278	1308	1.5(10)-133	Escherichia coli	b0032	[pn:carbamoyl-phosphate synthase small chain] [gn:carc]
CONTIG421	25524180_f3_34	1769	7431	3243	1081	5312	0	Escherichia coli	b0033	[pn:carbamoyl-phosphate synthase large chain] [gn:carb]
CONTIG421	34180387_c1_57	1770	7432	1299	433	1816	2.2(10)-187	Escherichia coli	b0053	[pn:survival protein sura precursor] [gn:sura]
CONTIG421	33866441_c1_66	1771	7433	501	167	98	0.0011	Helicobacter pylori	AC000108	or:helicobacter pylori pn:orf33 le:34041 re:34685 di:direct
CONTIG421	2371025_c2_75	1772	7434	1437	479	2091	1.6(10)-216	Escherichia coli	b0054	[pn:organic solvent tolerance protein precursor] [gn:imp]

CONTIG421	14881910_c2_76	1773	7435	828	276	1320	7.9(10)-135	Escherichia coli	b0051	[pn:dimethyladenosine transferase] [gn:ksga]
CONTIG421	25431533_c2_77	1774	7436	381	127	506	1.3(10)-48	Escherichia coli	b0050	[pn:apag protein] [gn:apag]
CONTIG421	24083442_c3_91	1775	7437	1050	350	1384	1.3(10)-141	Escherichia coli	b0052	[pn:pyridoxal phosphate biosynthetic protein pda] [gn:pdxa]
CONTIG421	31664075_c3_92	1776	7438	969	323	1366	1.1(10)-139	Escherichia coli	b0049	[pn:bis] [gn:apah]
CONTIG421	35285415_c3_97	1777	7439	471	157	100	0.0076	Caenorhabditis elegans	AF022974	[de:caenorhabditis elegans cosmid f26g5.] [gn:f26g5.9] [nt:contains similarity to c3hc4-type zinc
CONTIG421	23626562_c3_100	1778	7440	3102	1034	441	4.5(10)-65	Escherichia coli	b1509	[pn:hypothetical protein]
CONTIG422	10971967_fl_3	1779	7441	1221	407	2054	1.3(10)-212	Escherichia coli	b2290	[pn:hypothetical protein]
CONTIG422	21759631_fl_4	1780	7442	603	201	950	1.3(10)-95	Escherichia coli	b2291	[pn:hypothetical protein]
CONTIG422	2931592_fl_21	1781	7443	435	145	183	4.5(10)-14	Escherichia coli	b3438	[pn:gtmkr operon regulator] [gn:gnt]
CONTIG422	25833401_fl_22	1782	7444	624	208	262	1.0(10)-22	Escherichia coli	b3438	[pn:gtmkr operon regulator] [gn:gnt]
CONTIG422	33203141_f2_34	1783	7445	1239	413	1902	1.7(10)-196	Escherichia coli	b2296	[pn:acetate kinase] [gn:acka]
CONTIG422	31728532_f2_35	1784	7446	2145	715	3368	0	Escherichia coli	b2297	[pn:phosphate acetyltransferase] [gn:pta]
CONTIG422	31678191_c1_71	1785	7447	636	212	795	3.3(10)-79	Escherichia coli	b2299	[pn:hypothetical protein]
CONTIG422	2786715_c1_75	1786	7448	306	102	94	6.5(10)-5	Escherichia coli	b4194	[pn:hypothetical 10.9 kd protein in aidb-rpsf intergenic region] [gn:yft]
CONTIG422	26620418_c1_76	1787	7449	1404	468	435	4.7(10)-41	Escherichia coli	b4193	[pn:hypothetical 52.9 kd protein in aidb-rpsf intergenic region] [gn:yfs]
CONTIG422	16683451_c1_77	1788	7450	1008	336	462	6.5(10)-44	Methanococcus jannaschii	MJ0679	[pn:transketolase <sup>mtm</sup> ]
CONTIG422	4429018_c1_84	1789	7451	465	155	738	3.7(10)-73	Escherichia coli	b2295	[pn:hypothetical protein]
CONTIG422	26681415_c2_108	1790	7452	678	226	703	1.8(10)-69	Escherichia coli	b2293	[pn:hypothetical protein]
CONTIG422	6039811_c2_109	1791	7453	1845	615	2338	1.1(10)-242	Escherichia coli	b2292	[pn:hypothetical protein] [gn:yfs]

CONTIG422	5111062_c2_111	1792	7454	894	298	1222	1.8(10)-124	Escherichia coli	b2289	[pn:nadh dehydrogenase operon transcriptional regulator] [gn:lrha]
CONTIG422	2931576_c3_112	1793	7455	555	185	835	2.0(10)-83	Escherichia coli	b2300	[pn:hypothetical protein] [gn:yfce]
CONTIG422	23853437_c3_114	1794	7456	447	149	238	3.6(10)-20	Escherichia coli	b4195	[pn:hypothetical phosphotransferase enzyme ii] [gn:ptxa]
CONTIG422	24507291_c3_117	1795	7457	858	286	462	6.5(10)-44	Methanococcus jannaschii	MJ0681	[pn:transketolase <sup>III</sup> ]
CONTIG422	13725686_c3_122	1796	7458	549	183	797	2.1(10)-79	Escherichia coli	b2294	[pn:hypothetical protein]
CONTIG423	36125431_fl_7	1797	7459	216	72	105	4.5(10)-6	Bacteriophage phi-80	P17651	adsorption-inhibiting cor protein.
CONTIG423	24222277_f2_27	1798	7460	1383	461	1955	4.0(10)-202	Escherichia coli	b1784	[pn:hypothetical protein]
CONTIG423	7151578_f3_45	1799	7461	264	88	429	3.7(10)-40	Escherichia coli	b1783	[pn:hypothetical protein]
CONTIG423	30522566_c1_70	1800	7462	3363	1121	248	6.0(10)-20	Salmonella typhimurium	AF007380	[PN:lambda phage H tail component homolog] [DE:Salmonella typhimurium lambda phage K tail component homolog gene,partial cds, lambda phage L tail component homolog, copper-zincsuperoxide dismutase (sodC), attachment and invasion prote
CONTIG423	6286331_c1_74	1801	7463	4020	1340	1029	2.7(10)-115	Bacteriophage lambda	P03749	host specificity protein j.
CONTIG423	665907_c2_89	1802	7464	762	254	329	8.1(10)-30	Yersinia pestis	AF053947	[de:yersinia pestis plasmid pmt1, complete plasmid sequence.]
CONTIG423	36534812_c2_91	1803	7465	609	203	128	2.3(10)-8	Bacteriophage lambda	P03730	[pn:phage lambda minor tail protein i homolog]
CONTIG423	20916702_c2_100	1804	7466	672	224	93	0.066	Bacillus subtilis	yhaU	tail assembly protein i.
CONTIG423	26584831_c2_102	1805	7467	360	120	91	0.00013	Salmonella typhimurium	P23831	[pn:hypothetical protein]
CONTIG423	22948562_c3_106	1806	7468	1122	374	581	1.6(10)-56	Bacteriophage HK97	P49861	sama protein (ec 3.4.21.-).
CONTIG423										major capsid protein precursor (gp5) (head protein).

CONTIG423	31728755_c3_112	1807	7469	387	129	136	2.2(10)-9	Bacteriophage lambda	P03737	minor tail protein m.
CONTIG423	25524180_c3_114	1808	7470	723	241	387	5.7(10)-36	Coxiella burnetii	Y15898	[de:coxiella burnetii plasmid qprs dna.] [pn:hypothetical protein] [gn:orf 248]
CONTIG424	33641631_fl_6	1809	7471	1398	466	1791	9.6(10)-185	Escherichia coli	b1002	[pn:glucose-1-phosphatase precursor] [gn:agp]
CONTIG424	2846890_fl_8	1810	7472	195	65	210	3.2(10)-17	Escherichia coli	b1259	[pn:hypothetical protein in tonb-trpa intergenic region] [gn:ycig]
CONTIG424	33807708_fl_15	1811	7473	666	222	908	3.6(10)-91	Escherichia coli	b1013	[pn:hypothetical protein] [gn:ycdc]
CONTIG424	9776952_fl_19	1812	7474	651	217	136	2.2(10)-9	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:59 re:580 di:direct nt:orf2
CONTIG424	9964202_f2_33	1813	7475	534	178	151	5.9(10)-11	Helicobacter pylori	HP0571	[pn:conserved hypothetical integral membrane protein]
CONTIG424	33235452_f2_54	1814	7476	1509	503	2172	4.0(10)-225	Escherichia coli	b1015	[pn:sodium/proline symporter] [gn:putp]
CONTIG424	23862882_f3_62	1815	7477	975	325	151	1.3(10)-8	Bacillus subtilis	yoaV	[pn:hypothetical protein]
CONTIG424	22266038_c1_80	1816	7478	1092	364	1129	1.3(10)-114	Escherichia coli	b2393	[pn:nucleoside permease nupc] [gn:nupc]
CONTIG424	13067881_c1_86	1817	7479	324	108	91	0.00479	Escherichia coli	f53597	proline dehydrogenase (ec 1.5.99.8) - escherichia coli
CONTIG424	32290750_c1_93	1818	7480	441	147	596	4.2(10)-58	Escherichia coli	b1010	[pn:hypothetical protein]
CONTIG424	32425751_c1_100	1819	7481	1116	372	870	3.7(10)-87	Providencia stuartii	U23806	or:providencia stuartii le:343 re:1413 di:direct nt:extended orf of mgte gene; transcription from this
CONTIG424	14455001_c1_101	1820	7482	537	179	97	0.0038	Bacillus subtilis	yxjL	[pn:hypothetical protein]
CONTIG424	4803205_c2_105	1821	7483	4092	1364	5754	0	Escherichia coli	b1014	[pn:proline oxidase] [gn:puta]
CONTIG424	23564416_c2_106	1822	7484	447	149	207	6.9(10)-17	Sinorhizobium meliloti	P42879	hypothetical 15.0 kd protein in ureb- urec intergenic region (orf5).
CONTIG424	31813125_c2_110	1823	7485	783	261	1104	6.0(10)-112	Escherichia coli	b1011	[pn:hypothetical protein]
CONTIG424	32619542_c2_111	1824	7486	858	286	1022	3.0(10)-103	Escherichia coli	b1009	[pn:hypothetical protein]

CONTIG424	16614702_c2_112	1825	7487	591	197	874	1.3(10)-87	Escherichia coli	b1008	[pn:hypothetical protein]
CONTIG424	16835915_c2_120	1826	7488	603	201	807	1.8(10)-80	Escherichia coli	b1004	[pn:trp repressor binding protein] [gn:wrbai]
CONTIG424	34645256_c2_121	1827	7489	246	82	366	9.8(10)-34	Escherichia coli	b1003	[pn:hypothetical 8.5 kd protein in agn 3'''region] [gn:ycgj]
CONTIG424	32661281_c3_132	1828	7490	183	61	117	8.1(10)-6	Escherichia coli	I53597	proline dehydrogenase (ec 1.5.99.8) - escherichia coli
CONTIG424	23625786_c3_138	1829	7491	1179	393	1701	3.2(10)-175	Escherichia coli	b1012	[pn:hypothetical protein]
CONTIG424	4332318_c3_141	1830	7492	504	168	707	7.2(10)-70	Escherichia coli	b1007	[pn:hypothetical protein]
CONTIG424	33601030_c3_142	1831	7493	1344	448	1835	2.1(10)-189	Escherichia coli	b1006	[pn:hypothetical protein] [gn:ycdg]
CONTIG425	22659426_fl_1	1832	7494	2682	894	3783	0	Escherichia coli	b1834	[pn:hypothetical protein]
CONTIG425	11176451_fl_4	1833	7495	285	95	350	4.9(10)-32	Escherichia coli	b1836	[pn:hypothetical protein]
CONTIG425	35599132_fl_5	1834	7496	1104	368	524	1.8(10)-50	Escherichia coli	b0877	[pn:hypothetical protein]
CONTIG425	7159682_fl_10	1835	7497	273	91	333	3.1(10)-30	Escherichia coli	b1842	[pn:dna polymerase iii, theta subunit] [gn:hole]
CONTIG425	26050762_fl_11	1836	7498	684	228	332	2.0(10)-41	Escherichia coli	b1843	[pn:hypothetical protein]
CONTIG425	32635763_fl_12	1837	7499	678	226	1052	2.0(10)-106	Escherichia coli	b1844	[pn:hypothetical protein]
CONTIG425	4803751_f3_44	1838	7500	1314	438	1773	7.7(10)-183	Escherichia coli	b1833	[pn:hypothetical protein]
CONTIG425	29781327_f3_46	1839	7501	1464	488	2052	2.1(10)-212	Escherichia coli	b1835	[pn:hypothetical protein]
CONTIG425	24349067_f3_54	1840	7502	1275	425	1599	2.2(10)-164	Escherichia coli	b1849	[pn:phosphoribosylglycinamide formyltransferase 2] [gn:put]



CONTIG425	33494665_c1_58	1841	7503	285	95	91	0.0027	Triticum aestivum	JN0690	glutenin, high-molecular-weight bx7 chain precursor - wheat the main wheat storage proteins are divided into two groups: the glutenins, composed of high- and low-molecular weight families, and the gliadins, composed of alpha or beta, gamma and omega fam
CONTIG425	3158387_c1_62	1842	7504	876	292	545	1.1(10)-52	Escherichia coli	b1840	[pn:hypothetical protein]
CONTIG425	7228508_c2_77	1843	7505	687	229	529	5.2(10)-51	Escherichia coli	b1846	[pn:hypothetical 23.7 kd protein in purt 5'''region] [gn:yebe]
CONTIG425	32225388_c2_78	1844	7506	2103	701	2793	6.4(10)-291	Escherichia coli	b1845	[pn:protease ii] [gn:ptrb]
CONTIG425	24415886_c2_82	1845	7507	342	114	444	5.2(10)-42	Escherichia coli	b1839	[pn:hypothetical protein]
CONTIG425	24025302_c2_93	1846	7508	501	167	754	7.5(10)-75	Escherichia coli	b1832	[pn:hypothetical protein]
CONTIG425	196055_c2_94	1847	7509	516	172	325	2.2(10)-29	Haemophilus influenzae	HI1670	[pn:conjunctive transfer co-repressor] [gn:fino]
CONTIG425	4976566_c3_104	1848	7510	390	130	272	9.0(10)-24	Escherichia coli	b1841	[pn:hypothetical protein]
CONTIG425	24897717_c3_110	1849	7511	777	259	704	1.5(10)-69	Escherichia coli	b1838	[pn:hypothetical protein] [gn:prpa]
CONTIG426	26460813_f1_27	1850	7512	216	72	178	8.1(10)-14	Escherichia coli	b3537	[pn:hypothetical protein] [gn:yhit]
CONTIG426	25987562_f2_49	1851	7513	1587	529	2040	4.0(10)-211	Escherichia coli	b3536	[pn:hypothetical 59.4 kd protein in deta-dppf intergenic region]
CONTIG426	31666653_f2_50	1852	7514	1713	571	2368	7.0(10)-246	Escherichia coli	b3538	[pn:hypothetical 62.0 kd protein in deta-dppf intergenic region]
CONTIG426	19556892_c1_85	1853	7515	1014	338	1364	1.7(10)-139	Escherichia coli	b3973	[pn:bifunctional protein] [gn:bira]
CONTIG426	16691707_c1_92	1854	7516	2643	881	3852	0	Escherichia coli	b3533	[pn:hypothetical 101.6 kd protein in deta-dppf intergenic region]
CONTIG426	35657127_c1_98	1855	7517	3507	1169	4218	0	Escherichia coli	b3530	[pn:hypothetical 125.7 kd protein in deta-dppf intergenic region]
CONTIG426	23455028_c1_99	1856	7518	2100	700	2387	6.7(10)-248	Escherichia coli	b3529	[pn:hypothetical 73.1 kd protein in deta-dppf intergenic region]

CONTIG426	16598131_c1_100	1857	7519	567	189	570	2.3(10)-55	Escherichia coli	b3528	[pn:c4-dicarboxylate transport protein] [gn:dcta]
CONTIG426	33719693_c2_102	1858	7520	1176	392	1474	3.7(10)-151	Escherichia coli	b3972	[pn:udp-n-acetylenolpyruvoylglucosamine reductase] [gn:murb]
CONTIG426	14551878_c2_107	1859	7521	783	261	897	5.2(10)-90	Escherichia coli	b3534	[pn:hypothetical protein] [gn:yhiq]
CONTIG426	21769375_c2_111	1860	7522	2541	847	2874	0	Escherichia coli	b3532	[pn:hypothetical 86.0 kd protein in dcta-dppf intergenic region]
CONTIG426	24650458_c2_112	1861	7523	1113	371	1417	4.2(10)-145	Escherichia coli	b3531	[pn:hypothetical 41.7 kd protein in dcta-dppf intergenic region]
CONTIG426	32319840_c3_124	1862	7524	249	83	245	6.5(10)-21	Escherichia coli	b3535	[pn:hypothetical protein] [gn:yhjr]
CONTIG426	32035966_c3_137	1863	7525	348	116	94	0.00095	Pseudomonas sp.	D10769	or:pseudomonas sp. pn:maltopectase forming amylase le:717 re:2561 di:direct sr:pseudomonas sp. (strain ko-8940) (library: lambda [47] dna
CONTIG427	4152178_f1_8	1864	7526	1020	340	1216	8.3(10)-124	Escherichia coli	b1320	[pn:hypothetical protein]
CONTIG427	4823265_f2_20	1865	7527	807	269	1160	7.0(10)-118	Escherichia coli	b1326	[pn:hypothetical protein] [gn:ycj]
CONTIG427	24265676_f2_22	1866	7528	531	177	727	5.5(10)-72	Escherichia coli	b1324	[pn:thiol peroxidase] [gn:tpx]
CONTIG427	31681349_c1_59	1867	7529	838	279	1209	4.5(10)-123	Escherichia coli	b1313	[pn:hypothetical protein]
CONTIG427	5350202_c1_60	1868	7530	1077	359	1412	1.3(10)-144	Escherichia coli	b1315	[pn:hypothetical protein]
CONTIG427	7314416_c1_66	1869	7531	765	255	596	4.2(10)-58	Escherichia coli	b1317	[pn:hypothetical protein]
CONTIG427	32842_c1_68	1870	7532	927	309	990	7.4(10)-100	Escherichia coli	b1319	[pn:hypothetical protein] [gn:ompg]
CONTIG427	10634640_c1_76	1871	7533	1032	344	1162	4.4(10)-118	Escherichia coli	b1325	[pn:hypothetical protein]
CONTIG427	5165892_c2_81	1872	7534	858	286	1201	3.2(10)-122	Escherichia coli	b1314	[pn:hypothetical protein]
CONTIG427	35236466_c2_84	1873	7535	1125	375	1378	5.5(10)-141	Escherichia coli	b1318	[pn:hypothetical protein]

CONTIG427	14885377_c2_89	1874	7536	1059	353	1256	4.7(10)-128	Escherichia coli	b1322	[pn:hypothetical protein in pspe-tyrr intergenic region] [gn:ycjif]
CONTIG427	3411330_c2_90	1875	7537	1638	546	2210	3.8(10)-229	Escherichia coli	b1323	[pn:transcriptional regulatory protein tyr] [gn:tyrr]
CONTIG427	24505167_c3_96	1876	7538	2310	770	2548	5.9(10)-265	Escherichia coli	b1316	[pn:hypothetical protein]
CONTIG427	5213467_c3_100	1877	7539	1407	469	2285	4.4(10)-237	Escherichia coli	b1321	[pn:hypothetical protein]
CONTIG427	4157968_c3_108	1878	7540	1626	542	2510	6.2(10)-261	Escherichia coli	b1329	[pn:hypothetical protein]
CONTIG428	16932050_f1_22	1879	7541	1098	366	538	5.7(10)-52	Bacillus subtilis	yesQ	[pn:hypothetical protein]
CONTIG428	13867963_f1_23	1880	7542	1140	380	818	1.2(10)-81	Bacillus subtilis	msmX	[pn:multiple sugar-binding transport atp-binding protein] [gn:yxkg]
CONTIG428	4807327_f2_30	1881	7543	1023	341	1503	3.2(10)-154	Escherichia coli	b2837	[pn:galactose operon repressor] [gn:galr]
CONTIG428	32149135_f2_38	1882	7544	957	319	1322	4.7(10)-135	Escherichia coli	b2839	[pn:transcriptional activator protein lysr] [gn:lysr]
CONTIG428	24427336_f2_41	1883	7545	336	112	218	4.7(10)-18	Erwinia chrysanthemi	Q05527	pectin degradation protein kdgl
CONTIG428	4345932_f2_47	1884	7546	1350	450	327	1.3(10)-29	Bacillus subtilis	yesO	[pn:hypothetical protein]
CONTIG428	2468768_f3_59	1885	7547	1119	373	692	2.7(10)-68	Escherichia coli	b2714	[pn:asc operon repressor protein] [gn:ascg]
CONTIG428	24642553_f3_65	1886	7548	897	299	775	4.5(10)-77	Bacillus subtilis	yesP	[pn:hypothetical protein]
CONTIG428	24026430_c1_81	1887	7549	1425	475	609	1.7(10)-59	Bacillus subtilis	licC	[pn:phosphotransferase system] [gn:celb]
CONTIG428	21587562_c2_104	1888	7550	1494	498	958	1.8(10)-96	Bacillus subtilis	ydhP	[pn:hypothetical protein]
CONTIG428	14257826_c2_105	1889	7551	795	265	560	2.7(10)-54	Haemophilus influenzae	HI0054	[pn:uxu operon regulator] [gn:uxur]
CONTIG428	4962950_c2_110	1890	7552	2322	774	3240	0	Escherichia coli	b2836	[pn:2-acylglycerophosphoethanolamine acyltransferase / acyl-acyl carrier protein synthetase] [gn:aas]
CONTIG428	9943775_c3_117	1891	7553	1215	405	1077	4.4(10)-109	Escherichia coli	b0587	[pn:ferric enterobactin transport protein fepe] [gn:fepe]

CONTIG428	34179511_c3_118	1892	7554	1185	395	895	8.5(10)-90	Escherichia coli	b2840	[pn:hypothetical 25.2 kd protein in lysr-arae intergenic region] [gn:ygea]
CONTIG428	4376958_c3_120	1893	7555	1365	455	1906	6.2(10)-197	Escherichia coli	b2838	[pn:diaminopimelate decarboxylase] [gn:lysa]
CONTIG428	1413592_c3_136	1894	7556	930	310	1155	2.3(10)-117	Escherichia coli	b2835	[pn:hypothetical protein in muth-aas intergenic region] [gn:yged]
CONTIG429	34664655_fl_2	1895	7557	540	180	420	1.8(10)-39	Escherichia coli	b0419	[pn:hypothetical protein] [gn:yajo]
CONTIG429	3414166_fl_3	1896	7558	363	121	225	1.1(10)-18	Escherichia coli	b0419	[pn:hypothetical protein] [gn:yajo]
CONTIG429	19964657_fl_22	1897	7559	609	203	217	6.0(10)-18	Bacillus subtilis	yyaR	[pn:hypothetical protein]
CONTIG429	428316_fl_23	1898	7560	591	197	461	8.4(10)-44	Bacillus subtilis	yhxD	[pn:hypothetical protein]
CONTIG429	21955468_f2_24	1899	7561	1209	403	325	2.2(10)-29	Bacillus subtilis	iolS	[pn:hypothetical protein] [gn:yxbf]
CONTIG429	2819755_f2_25	1900	7562	1074	358	480	8.0(10)-46	Synechocystis sp.	S76674	[PN:hypothetical protein] [OR:Synechocystis sp.] [SR:PCC 6803,, PCC 6803] [SR:PCC 6803,, ]
CONTIG429	4890901_f2_26	1901	7563	582	194	291	8.6(10)-26	Escherichia coli	b0046	[pn:hypothetical nadph oxidoreductase in fixc-kefc intergenic region] [gn:yabf]
CONTIG429	6750016_f2_31	1902	7564	642	214	215	9.8(10)-18	Bacillus subtilis	ywrO	[pn:hypothetical protein]
CONTIG429	24222287_f3_54	1903	7565	762	254	91	0.00052	Escherichia coli	b1518	[pn:hypothetical protein]
CONTIG429	5353556_f3_55	1904	7566	819	273	137	4.0(10)-9	Methanobacterium thermoautotrophicum	MTH234	[pn:gamma-carboxymuconolactone decarboxylase]
CONTIG429	30156668_f3_58	1905	7567	1491	497	597	3.2(10)-58	Escherichia coli	b1828	[pn:hypothetical protein]
CONTIG429	30329136_c1_90	1906	7568	549	183	398	4.0(10)-37	Mycobacterium tuberculosis	Z95210	unknown,, mtcy21c12.13, len
CONTIG429	33867840_c2_103	1907	7569	2484	828	965	1.3(10)-117	Arthrobacter sp.	S65769	maltooligosyl trehalose synthase - arthrobacter sp. (straining36)
CONTIG429	14730330_c2_104	1908	7570	2091	697	1423	9.5(10)-146	Escherichia coli	b3431	[pn:glycogen operon protein glgx] [gn:glgx]

CONTIG429	10551037_c2_112	1909	7571	963	321	168	2.7(10)-10	Escherichia coli	b0076	[pn:leuo]
CONTIG429	20157792_c2_113	1910	7572	294	98	129	1.3(10)-8	Mycobacterium tuberculosis	Z95210	unknown, mtcy21c12.12, mtcy21c12.12, len
CONTIG429	33601702_c3_120	1911	7573	1788	596	826	1.8(10)-82	Sulfolobus solfataricus	S73087	[pn:alpha-amylase, precursor]
CONTIG43	34018765_f2_1	1912	7574	663	221	1034	1.6(10)-104	Escherichia coli	b3320	[pn:50s ribosomal subunit protein 3]
CONTIG43	36517717_f3_3	1913	7575	368	123	510	5.4(10)-49	Escherichia coli	b3319	[pn:50s ribosomal subunit protein 4]
CONTIG430	25570176_f1_3	1914	7576	921	307	1276	3.6(10)-130	Escherichia coli	b0428	[pn:cytochrome o ubiquinol oxidase c subunit] [gn:cyoc]
CONTIG430	7089132_f1_4	1915	7577	1374	458	1801	8.5(10)-186	Escherichia coli	b0427	[pn:hypothetical protein] [gn:yajr]
CONTIG430	10835913_f1_18	1916	7578	615	205	677	1.1(10)-66	Escherichia coli	b0412	[pn:hypothetical 21.8 kd protein in tsx-ribd intergenic region] [gn:yaji]
CONTIG430	24869658_f2_19	1917	7579	2031	677	3197	0	Escherichia coli	b0431	[pn:cytochrome o ubiquinol oxidase subunit i] [gn:cyob]
CONTIG430	20801331_f2_20	1918	7580	423	141	297	2.0(10)-26	Escherichia coli	b0429	[pn:cytochrome o ubiquinol oxidase c subunit] [gn:cyod]
CONTIG430	5196068_f2_23	1919	7581	921	307	1376	9.1(10)-141	Escherichia coli	b0425	[pn:apba protein] [gn:apba]
CONTIG430	235707_f2_26	1920	7582	1005	335	1531	3.5(10)-157	Escherichia coli	b0419	[pn:hypothetical protein] [gn:yajo]
CONTIG430	14587812_f3_37	1921	7583	618	206	937	3.0(10)-94	Escherichia coli	b0430	[pn:cytochrome o ubiquinol oxidase subunit iii] [gn:cyoc]
CONTIG430	34648537_f3_44	1922	7584	609	203	905	7.5(10)-91	Escherichia coli	b0424	[pn:thij protein] [gn:thij]
CONTIG430	12922775_f3_47	1923	7585	249	83	357	8.8(10)-33	Escherichia coli	b0422	[pn:exodeoxyribonuclease small subunit] [gn:xseb]
CONTIG430	10157952_f3_48	1924	7586	900	300	1234	1.0(10)-125	Escherichia coli	b0421	[pn:geranyltransferase]
CONTIG430	2147132_f3_49	1925	7587	1887	629	2952	0	Escherichia coli	b0420	[pn:hypothetical protein] [gn:yajp]
CONTIG430	5161562_c1_60	1926	7588	1209	403	1658	1.2(10)-170	Escherichia coli	b0414	[pn:riboflavin biosynthesis protein ribd] [gn:ribd]
CONTIG430	4182842_c1_61	1927	7589	480	160	580	2.1(10)-56	Escherichia coli	b0415	[pn:probable riboflavin synthase beta chain] [gn:ribh]

CONTIG430	23671875_c1_62	1928	7590	441	147	661	5.4(10)-65	Escherichia coli	b0416	[pn:n utilization substance protein b] [gn:nusb]
CONTIG430	19781883_c2_99	1929	7591	555	185	775	4.5(10)-77	Escherichia coli	b0426	[pn:hypothetical protein] [gn:yajq]
CONTIG430	31430461_c3_108	1930	7592	462	154	723	1.3(10)-71	Escherichia coli	b0413	[pn:hypothetical 17.2 kd protein in tsx-ribd intergenic region] [gn:ybad]
CONTIG430	30212766_c3_109	1931	7593	975	325	1449	1.7(10)-148	Escherichia coli	b0417	[pn:hypothetical protein] [gn:thil]
CONTIG430	34005008_c3_116	1932	7594	1473	491	2182	3.6(10)-226	Escherichia coli	b0423	[pn:hypothetical protein] [gn:yajk]
CONTIG431	32236592_f1_4	1933	7595	2160	720	3307	0	Escherichia coli	b2675	[pn:ribonucleoside-diphosphate reductase 2 alpha chain] [gn:nrde]
CONTIG431	5320443_f1_5	1934	7596	969	323	1465	3.3(10)-150	Escherichia coli	b2676	[pn:ribonucleoside-diphosphate reductase 2 beta chain] [gn:nrdi]
CONTIG431	2067336_f1_7	1935	7597	1221	407	1908	3.8(10)-197	Escherichia coli	b2677	[pn:glycine betaine/l-proline transport atp-binding protein prov]
CONTIG431	10343930_f1_11	1936	7598	1212	404	954	4.7(10)-96	Escherichia coli	b2681	[pn:hypothetical protein]
CONTIG431	35604561_f1_14	1937	7599	1224	408	1632	6.7(10)-168	Escherichia coli	b2685	[pn:multidrug resistance protein a] [gn:emra]
CONTIG431	113337_f2_25	1938	7600	486	162	676	1.3(10)-66	Escherichia coli	b2670	[pn:hypothetical protein]
CONTIG431	31767930_f2_27	1939	7601	546	182	633	5.0(10)-62	Escherichia coli	b2674	[pn:hypothetical protein] [gn:nrdj]
CONTIG431	36604152_f2_31	1940	7602	1083	361	1416	5.2(10)-145	Escherichia coli	b2678	[pn:glycine betaine/l-proline transport system permease protein p] [gn:prow]
CONTIG431	5194693_f2_32	1941	7603	1005	335	1470	1.0(10)-150	Escherichia coli	b2679	[pn:glycine betaine-binding periplasmic protein precursor] [gn:prox]
CONTIG431	13007717_f2_34	1942	7604	1581	527	2347	1.2(10)-243	Escherichia coli	b2686	[pn:multidrug resistance protein b] [gn:emrb]
CONTIG431	22383582_f3_45	1943	7605	354	118	278	2.1(10)-24	Escherichia coli	b2672	[pn:hypothetical protein] [gn:ygam]
CONTIG431	32461077_f3_46	1944	7606	255	85	325	2.2(10)-29	Escherichia coli	b2673	[pn:hypothetical protein] [gn:nrdh]
CONTIG431	35166511_f3_57	1945	7607	627	209	872	2.3(10)-87	Escherichia coli	b2684	[pn:transcriptional repressor mpra] [gn:mpra]

CONTIG431	14070762_f3_59	1946	7608	1695	565	1454	5.0(10)-149	Saccharomyces cerevisiae	YNL104C	[pn:2-isopropylmalate synthase] [gn:leu4]
CONTIG431	26046955_c1_66	1947	7609	435	145	559	3.5(10)-54	Escherichia coli	b2690	[pn:hypothetical protein]
CONTIG431	35355165_c1_67	1948	7610	462	154	576	5.5(10)-56	Escherichia coli	b2689	[pn:hypothetical protein]
CONTIG431	23681890_c1_69	1949	7611	885	295	301	7.5(10)-27	Escherichia coli	b1790	[pn:hypothetical protein]
CONTIG431	4117193_c1_89	1950	7612	354	118	527	8.5(10)-51	Escherichia coli	b2671	[pn:hypothetical 13.1 kd protein in stpa-nrde intergenic region] [gn:ygac]
CONTIG431	33838307_c2_94	1951	7613	315	105	297	2.0(10)-26	Escherichia coli	b2690	[pn:hypothetical protein]
CONTIG431	1275266_c2_97	1952	7614	519	173	845	1.7(10)-84	Escherichia coli	b2687	[pn:hypothetical protein in emrb 3"] region] [gn:ygag]
CONTIG431	25390686_c3_117	1953	7615	1572	524	2378	6.0(10)-247	Escherichia coli	b2688	[pn:glutamate--cysteine ligase] [gn:gsha]
CONTIG432	24425931_f1_4	1954	7616	357	119	296	2.6(10)-26	Escherichia coli	b4023	[pn:hypothetical 10.5 kd protein in pepe-lysc intergenic region]
CONTIG432	5917286_f2_24	1955	7617	1362	454	1766	4.2(10)-182	Escherichia coli	b4024	[pn:lysine-sensitive aspartokinase iii] [gn:lysc]
CONTIG432	5162807_f2_25	1956	7618	1020	340	559	3.5(10)-54	Bacillus subtilis	yocS	[pn:hypothetical protein]
CONTIG432	30198511_f2_37	1957	7619	933	311	1323	3.7(10)-135	Escherichia coli	b4018	[pn:acetate operon repressor] [gn:iclr]
CONTIG432	34663402_f3_48	1958	7620	318	106	220	2.8(10)-18	Haemophilus influenzae	HI1419	[pn:hypothetical protein]
CONTIG432	22692827_f3_49	1959	7621	294	98	179	6.4(10)-14	Haemophilus influenzae	HI1420	[pn:hypothetical protein]
CONTIG432	7218818_c1_67	1960	7622	942	314	1568	4.0(10)-161	Escherichia coli	b4013	[pn:homoserine o-succinyltransferase] [gn:meta]
CONTIG432	17051906_c1_71	1961	7623	1365	455	2134	4.4(10)-221	Escherichia coli	b4015	[pn:isocitrate lyase] [gn:acea]
CONTIG432	12203130_c1_72	1962	7624	1803	601	2698	7.5(10)-281	Escherichia coli	b4016	[pn:isocitrate dehydrogenase kinase/phosphatase] [gn:acek]
CONTIG432	36415708_c1_75	1963	7625	3702	1234	5941	0	Escherichia coli	b4019	[pn:b12-dependent homocysteine-n5-methyltetrahydrofolate transmethyase] [gn:meth]

CONTIG432	2207187_c1_78	1964	7626	954	318	1276	3.6(10)-130	Escherichia coli	b4022	[pn:hypothetical 32.5 kd protein in pepe-lysc intergenic region]
CONTIG432	13708462_c2_87	1965	7627	225	75	124	5.2(10)-7	Escherichia coli	M18974	or:escherichia coli gn:acek le:472 re:2205 di:direct sr:e.coli (strain k12) dna nt:isocitrate dehydrogenase kinase/phosphatase
CONTIG432	86625_c3_104	1966	7628	1629	543	2464	4.7(10)-256	Escherichia coli	b4014	[pn:malate synthase a] [gn:aceb]
CONTIG432	12930443_c3_113	1967	7629	1647	549	2060	3.0(10)-213	Escherichia coli	b4020	[pn:hypothetical 59.5 kd protein in meth-pepe intergenic region] [gn:yjbb]
CONTIG433	783_f1_7	1968	7630	432	144	631	8.0(10)-62	Escherichia coli	b3167	[gn:ribosome-binding factor a] [gn:rbfa]
CONTIG433	14901591_f1_9	1969	7631	525	175	413	1.0(10)-38	Escherichia coli	b3165	[pn:30s ribosomal subunit protein s15] [gn:rpso]
CONTIG433	33364211_f1_20	1970	7632	555	185	762	1.1(10)-75	Escherichia coli	b3157	[pn:hypothetical 19.7 kd protein in soha-mtr intergenic region] [gn:yhbt]
CONTIG433	13677326_f2_28	1971	7633	492	164	740	2.2(10)-73	Escherichia coli	b3170	[pn:hypothetical 16.8 kd protein in nusa-mety intergenic region] [gn:yhbc]
CONTIG433	32694807_f2_34	1972	7634	1011	337	1485	2.6(10)-152	Escherichia coli	b3163	[pn:hypothetical 33.6 kd protein in dead-pnp intergenic region] [gn:yhbm]
CONTIG433	13129803_f2_35	1973	7635	1935	645	2085	3.2(10)-256	Escherichia coli	b3162	[pn:dead] [gn:dead]
CONTIG433	35656427_f2_42	1974	7636	723	241	867	8.0(10)-87	Escherichia coli	b3152	[pn:hypothetical 24.8 kd protein in agai-mtr intergenic region] [gn:yrar]
CONTIG433	13086018_f3_47	1975	7637	1518	506	2280	1.5(10)-236	Escherichia coli	b3169	[pn:n utilization substance protein a] [gn:nusa]
CONTIG433	36069132_f3_48	1976	7638	2709	903	2853	0	Escherichia coli	b3168	[pn:protein chain initiation factor 2] [gn:infb]
CONTIG433	32317906_f3_49	1977	7639	978	326	1389	3.8(10)-142	Escherichia coli	b3166	[pn:tma pseudouridine 55 synthase] [gn:trub]
CONTIG433	30367705_f3_50	1978	7640	2220	740	3047	0	Escherichia coli	b3164	[pn:polynucleotide phosphorylase] [gn:pnp]
CONTIG433	16833293_f3_55	1979	7641	1251	417	1821	6.4(10)-188	Escherichia coli	b3161	[pn:tryptophan-specific permease] [gn:mtr]
CONTIG433	5103407_f3_60	1980	7642	507	169	753	9.5(10)-75	Escherichia coli	b3156	[pn:hypothetical protein] [gn:yhbs]



CONTIG433	24742838_f3_61	1981	7643	447	149	607	2.7(10)-59	Escherichia coli	b3154	[pn:hypothetical 16.8 kd protein in soha-mtr intergenic region]
CONTIG433	36422036_f3_64	1982	7644	738	246	757	3.6(10)-75	Escherichia coli	b3151	[pn:hypothetical 37.3 kd protein in agai-mtr intergenic region] [gn:yraa]
CONTIG433	6745840_c1_68	1983	7645	363	121	349	6.2(10)-32	Escherichia coli	b3155	[pn:hypothetical 11.3 kd protein in soha-mtr intergenic region]
CONTIG433	22902158_c1_72	1984	7646	1038	346	1437	3.2(10)-147	Escherichia coli	b3159	[pn:hypothetical 33.2 kd protein in soha-mtr intergenic region]
CONTIG433	14875383_c1_73	1985	7647	1014	338	1451	1.0(10)-148	Escherichia coli	b3160	[pn:hypothetical 37.1 kd protein in soha-mtr intergenic region] [gn:yhbw]
CONTIG433	7071032_c2_91	1986	7648	456	152	98	0.00479	Caenorhabditis elegans	Z93395	[de:caenorhabditis elegans cosmid zc101, complete sequence.] [pn:zc101.1] [nt:similar to low-density lipoprotein receptor]
CONTIG433	24023542_c2_92	1987	7649	540	180	794	4.2(10)-79	Escherichia coli	b3153	[pn:hypothetical 20.3 kd protein in soha-mtr intergenic region]
CONTIG433	12001058_c2_128	1988	7650	1125	375	1780	1.3(10)-183	Escherichia coli	b3172	[pn:argininosuccinate synthetase] [gn:argg]
CONTIG433	3337752_c3_133	1989	7651	999	333	1494	2.8(10)-153	Escherichia coli	b3158	[pn:putative protease in soha-mtr intergenic region] [gn:yhbu]
CONTIG433	21675430_c3_166	1990	7652	321	107	406	5.7(10)-38	Escherichia coli	b3172	[pn:argininosuccinate synthetase] [gn:argg]
CONTIG434	16053507_f1_2	1991	7653	474	158	481	6.4(10)-46	Escherichia coli	b1104	[pn:hypothetical protein]
CONTIG434	4876342_f1_9	1992	7654	270	90	314	3.2(10)-28	Escherichia coli	b1112	[pn:hypothetical protein]
CONTIG434	25520382_f1_20	1993	7655	1269	423	1490	7.5(10)-153	Escherichia coli	b1118	[pn:hypothetical protein]
CONTIG434	35330125_f1_21	1994	7656	966	322	1287	2.5(10)-131	Escherichia coli	b1119	[pn:hypothetical protein] [gn:ycfx]
CONTIG434	23478157_f2_28	1995	7657	651	217	757	3.6(10)-75	Escherichia coli	b1105	[pn:hypothetical protein]
CONTIG434	11072040_f2_29	1996	7658	1068	356	1580	2.2(10)-162	Escherichia coli	b1107	[pn:hypothetical protein]
CONTIG434	7323500_f2_30	1997	7659	1320	440	1937	3.2(10)-200	Escherichia coli	b1109	[pn:nadh dehydrogenase] [gn:ndh]
CONTIG434	640966_f2_40	1998	7660	1944	648	1372	2.3(10)-140	Escherichia coli	b1116	[pn:hypothetical protein]

CONTIG434	7226712_f3_49	1999	7661	372	124	584	7.7(10)-57	Escherichia coli	b1103	[pn:hypothetical protein in flue-ndh intergenic region] [gn:ycff]
CONTIG434	33605040_f3_52	2000	7662	864	288	879	4.2(10)-88	Escherichia coli	b1106	[pn:hypothetical protein]
CONTIG434	15049141_f3_53	2001	7663	555	185	938	2.3(10)-94	Escherichia coli	b1108	[pn:hypothetical protein]
CONTIG434	421905_f3_55	2002	7664	612	204	697	8.1(10)-69	Escherichia coli	b1110	[pn:hypothetical 18.9 kd protein in ndh-mfd intergenic region] [gn:ycf]
CONTIG434	14973833_f3_63	2003	7665	900	300	1044	1.3(10)-105	Escherichia coli	b1120	[pn:hypothetical protein]
CONTIG434	29329666_f3_70	2004	7666	279	93	251	3.0(10)-21	Escherichia coli	b1127	[pn:peptidase t] [gn:pept]
CONTIG434	10195287_c1_74	2005	7667	1059	353	1637	2.0(10)-168	Escherichia coli	b1123	[pn:spermidine/putrescine-binding periplasmic protein precursor] [gn:potd]
CONTIG434	1195910_c1_84	2006	7668	1155	385	1221	2.3(10)-124	Escherichia coli	b1115	[pn:hypothetical protein]
CONTIG434	12362887_c2_99	2007	7669	1212	404	1650	8.5(10)-170	Escherichia coli	b1126	[pn:spermidine/putrescine transport atp-binding protein pota] [gn:pot]
CONTIG434	16689813_c2_101	2008	7670	807	269	943	7.0(10)-95	Escherichia coli	b1124	[pn:spermidine/putrescine transport system permease protein potc] [gn:potc]
CONTIG434	6483817_c2_108	2009	7671	3513	1171	5326	0	Escherichia coli	b1114	[pn:transcription-repair coupling factor] [gn:mfd]
CONTIG434	14460050_c2_109	2010	7672	1020	340	1377	7.2(10)-141	Escherichia coli	b1113	[pn:hypothetical protein]
CONTIG434	34657312_c3_120	2011	7673	882	294	1258	2.8(10)-128	Escherichia coli	b1125	[pn:spermidine/putrescine transport system permease protein potb] [gn:potb]
CONTIG434	24018801_c3_140	2012	7674	669	223	875	1.1(10)-87	Escherichia coli	b1111	[pn:hypothetical protein]
CONTIG435	21774136_fl_3	2013	7675	846	282	205	1.1(10)-16	Bacillus subtilis	ylsO	[pn:hypothetical protein]
CONTIG435	5214541_fl_9	2014	7676	2742	914	1145	2.7(10)-116	Escherichia coli	b2592	[pn:clpb protein] [gn:clpb]
CONTIG435	26681587_fl_10	2015	7677	1437	479	159	4.0(10)-8	Saccharomyces cerevisiae	YOL045 W	[pn:similarity to ser/thr protein kinase]
CONTIG435	25782568_fl_12	2016	7678	789	263	139	9.4(10)-8	Serratia liquefaciens	P18954	phb protein precursor.

CONTIG435	32303567_fl_14	2017	7679	1194	398	223	1.6(10)-15	Escherichia coli	AF044503	[de:escherichia coli strain ec11 unknown (498), hcp gene, complete cds;and rhsg accessory genetic element vrg protein, core component anddsorf-g1 genes, complete cds.] [pn:vrg protein]
CONTIG435	22832555_f2_15	2018	7680	1797	599	96	0.04399	Bos taurus	U92535	neuronal axonal membrane protein,,nap-22 homolog
CONTIG435	16588251_f3_48	2019	7681	453	151	326	1.7(10)-29	Escherichia coli	b2055	[pn:hypothetical protein] [gn:wcae]
CONTIG436	292882_f2_49	2020	7682	1164	388	353	2.2(10)-32	Azorhizobium caulinodans	S52856	arae-like protein - azorhizobium
CONTIG436	95380_f2_50	2021	7683	1089	363	955	3.7(10)-96	Bacillus subtilis	gap	caulinodans
CONTIG436	11198311_f2_51	2022	7684	1725	575	2264	7.2(10)-235	Escherichia coli	b1197	[pn:glyceraldehyde-3-phosphate dehydrogenase]
CONTIG436	29383457_c1_83	2023	7685	1725	575	391	2.8(10)-63	Escherichia coli	b0150	[pn:periplasmic trehalase precursor] [gn:trea]
CONTIG436	36035686_c1_93	2024	7686	3666	1222	170	3.3(10)-21	Legionella pneumophila	Y15044	[pn:ferrichrome-iron receptor precursor] [gn:fhua]
CONTIG436	14511687_c2_100	2025	7687	270	90	103	0.00012	Escherichia coli	b0150	[de:legionella pneumophila 22kb dna fragment from icm gene cluster.] [pn:icmf protein] [gn:icmf]
CONTIG436	4957906_c2_110	2026	7688	1269	423	165	4.2(10)-10	Bacillus subtilis	motB	[pn:ferrichrome-iron receptor precursor] [gn:fhua]
CONTIG437	24870905_fl_1	2027	7689	1032	344	766	4.0(10)-76	Escherichia coli	b3826	[pn:motility protein b] [gn:mot]
CONTIG437	10995841_fl_11	2028	7690	660	220	828	1.1(10)-82	Escherichia coli	b3834	[pn:yigJ]
CONTIG437	19956550_fl_15	2029	7691	282	94	288	1.8(10)-25	Escherichia coli	b3836	[pn:hypothetical 22.3 kd protein in udp-rfah intergenic region] [gn:yigp]
CONTIG437	25791025_fl_16	2030	7692	546	182	472	5.7(10)-45	Escherichia coli	b3838	[pn:hypothetical 11.3 kd protein in udp-rfah intergenic region]
CONTIG437	2112687_fl_22	2031	7693	520	174	686	1.2(10)-67	Escherichia coli	b3554	[pn:hypothetical 15.6 kd protein in udp-rfah intergenic region]
CONTIG437	4569713_f2_23	2032	7694	912	304	1307	1.8(10)-133	Escherichia coli	b3827	[pn:hypothetical 30.2 kd protein in bise-cspa intergenic region] [gn:yiaf]
CONTIG437										[pn:hypothetical 33.7 kd protein in pldb-metr intergenic region] [gn:yigm]

CONTIG437	197802_f2_25	2033	7695	2289	763	3781	0	Escherichia coli	b3829	[pn:5-methyltetrahydropteroylriglutamate-homocysteine methyltransferase] [gn:mete]
CONTIG437	50892_f2_28	2034	7696	810	270	1241	1.8(10)-126	Escherichia coli	b3831	[pn:uridine phosphorylase] [gn:udp]
CONTIG437	12694067_f2_32	2035	7697	831	277	758	2.7(10)-75	Escherichia coli	b3840	[pn:yigw]
CONTIG437	33751042_f2_33	2036	7698	1515	505	2515	1.8(10)-261	Escherichia coli	b3843	[pn:hypothetical 55.3 kd protein in rfah-rfe intergenic region] [gn:yigc]
CONTIG437	22519382_f2_34	2037	7699	723	241	1083	1.0(10)-109	Escherichia coli	b3844	[pn:flavin reductase] [gn:ubib]
CONTIG437	36017000_f2_35	2038	7700	1638	546	654	3.0(10)-64	Streptomyces lincolnensis	S69834	,
CONTIG437	25595458_f3_37	2039	7701	228	76	334	2.3(10)-30	Escherichia coli	b3825	[pn:lysophospholipase 12] [gn:pldb]
CONTIG437	2032755_f3_47	2040	7702	1602	534	2022	3.2(10)-209	Escherichia coli	b3832	[pn:hypothetical 54.7 kd protein in udp 3''' region precursor] [gn:yign]
CONTIG437	10444800_f3_48	2041	7703	771	257	1228	4.4(10)-125	Escherichia coli	b3833	[pn:hypothetical 28.1 kd protein in udp-rfah intergenic region] [gn:yigo]
CONTIG437	22062915_f3_49	2042	7704	1677	559	2532	2.8(10)-263	Escherichia coli	b3835	[pn:hypothetical 63.2 kd protein in udp-rfah intergenic region]
CONTIG437	4491042_f3_50	2043	7705	780	260	1000	6.4(10)-101	Escherichia coli	b3839	[pn:hypothetical 29.0 kd protein in udp-rfah intergenic region] [gn:yigu]
CONTIG437	6102030_c1_58	2044	7706	318	106	310	8.4(10)-28	Escherichia coli	b3555	[pn:hypothetical 11.0 kd protein in bise-cspa intergenic region]
CONTIG437	22128816_c1_65	2045	7707	546	182	678	8.5(10)-67	Escherichia coli	b3842	[pn:transcriptional activator] [gn:rfah]
CONTIG437	19971006_c1_79	2046	7708	861	287	850	5.0(10)-85	Escherichia coli	b3830	[pn:hypothetical protein]
CONTIG437	16830043_c2_110	2047	7709	444	148	157	4.9(10)-11	Escherichia coli	b3830	[pn:hypothetical protein]
CONTIG437	4297842_c2_114	2048	7710	963	321	1466	2.7(10)-150	Escherichia coli	b3828	[pn:trans-activator of mete and meth] [gn:metr]
CONTIG438	12601687_f1_3	2049	7711	1506	502	566	6.2(10)-55	Bacillus subtilis	yhdl	[pn:hypothetical protein]
CONTIG438	32204502_f1_13	2050	7712	666	222	750	2.0(10)-74	Escherichia coli	b0464	[pn:potential acraB operon repressor] [gn:acrr]

CONTIG438	13719682_fl_18	2051	7713	618	206	889	3.7(10)-89	Escherichia coli	b0469	[pn:adenine phosphoribosyltransferase] [gn:apt]
CONTIG438	34257891_fl_20	2052	7714	354	118	324	2.7(10)-29	Escherichia coli	b0471	[pn:hypothetical 12.0 kd protein in dnax-recr intergenic region] [gn:ybab]
CONTIG438	11816261_f2_52	2053	7715	1974	658	2630	1.2(10)-273	Escherichia coli	b0470	[pn:dna polymerase iii subunits gamma and tau] [gn:dnax]
CONTIG438	22554702_f2_54	2054	7716	1875	625	2946	0	Escherichia coli	b0473	[pn:heat shock protein htpg] [gn:htpg]
CONTIG438	10750312_f2_55	2055	7717	678	226	1060	2.7(10)-107	Escherichia coli	b0474	[pn:adenylate kinase] [gn:adk]
CONTIG438	22517057_f2_57	2056	7718	1410	470	2087	4.2(10)-216	Escherichia coli	b0477	[pn:inosine-guanosine kinase] [gn:gsk]
CONTIG438	12210058_f3_78	2057	7719	3483	1161	3978	0	Escherichia coli	b0465	[pn:hypothetical protein] [gn:aefa]
CONTIG438	268831_f3_79	2058	7720	405	135	359	5.4(10)-33	Escherichia coli	b0468	[pn:hypothetical 14.8 kd protein in pric-apt intergenic region] [gn:yban]
CONTIG438	1988513_f3_82	2059	7721	618	206	906	5.9(10)-91	Escherichia coli	b0472	[pn:recombination protein recr] [gn:recr]
CONTIG438	16970952_f3_86	2060	7722	963	321	1377	7.2(10)-141	Escherichia coli	b0475	[pn:ferrochelatase] [gn:hemh]
CONTIG438	16604206_f3_87	2061	7723	270	90	109	8.3(10)-6	Escherichia coli	JU0314	hypothetical 34.6k protein (visa 3 region) - escherichiacoli
CONTIG438	33636577_f3_89	2062	7724	792	264	115	0.0014	Saccharomyces cerevisiae	YIR019C	[pn:extracellular alpha-1,4-glucan glucosidase] [gn:sta1]
CONTIG438	20369535_c1_96	2063	7725	312	104	90	0.0011	Homo sapiens	P29966	myristoylated alanine-rich c-kinase substrate (marcks) (protein kinase c substrate, 80 kd protein, light chain) (pkcs1) (80k-l protein).
CONTIG438	29901058_c1_103	2064	7726	519	173	277	2.6(10)-24	Escherichia coli	U82664	or:escherichia coli le:73500 re:73877 di:complement nt:hypothetical
CONTIG438	110457_c1_109	2065	7727	555	185	486	1.8(10)-46	Escherichia coli	b0467	[pn:primosomal replication protein n] [gn:pric]
CONTIG438	3229836_c1_110	2066	7728	189	63	168	9.4(10)-13	Escherichia coli	b0466	[pn:hypothetical 6.0 kd protein in accr-pric intergenic region]
CONTIG438	33851510_c1_116	2067	7729	1266	422	1539	4.9(10)-158	Escherichia coli	b0463	[pn:acriflavin resistance protein a precursor] [gn:acra]

CONTIG438	5085917_c1_120	2068	7730	201	67	103	7.2(10)-6	Escherichia coli	U82664	or:escherichia coli pn:hha protein gn:hha le:59538 re:5957 di:complement
CONTIG438	33673537_c1_121	2069	7731	573	191	761	1.3(10)-75	Escherichia coli	b0459	[pn:hypothetical protein] [gn:ylad]
CONTIG438	20100019_c2_126	2070	7732	278	92	277	2.6(10)-24	Escherichia coli	b0479	[pn:fosmidomycin resistance protein] [gn:fsr]
CONTIG438	23540913_c2_159	2071	7733	3114	1038	4072	0	Escherichia coli	b0462	[pn:acriflavin resistance protein b] [gn:acrb]
CONTIG438	31437757_c3_164	2072	7734	1740	580	2165	2.2(10)-224	Escherichia coli	b0478	[pn:hypothetical protein in gsk 3'''region] [gn:ybal]
CONTIG438	34431890_c3_188	2073	7735	414	138	633	5.0(10)-62	Escherichia coli	b0461	[pn:hypothetical 14.6 kd protein in hha-acrb intergenic region] [gn:ybai]
CONTIG438	3914143_c3_189	2074	7736	228	76	357	8.8(10)-33	Escherichia coli	b0460	[pn:haemolysin expression modulating protein] [gn:hha]
CONTIG438	4882717_c3_190	2075	7737	477	159	510	5.4(10)-49	Escherichia coli	b0458	[pn:hypothetical protein] [gn:ylac]
CONTIG438	2164755_c3_194	2076	7738	744	248	635	3.1(10)-62	Saccharomyces cerevisiae	P40586	hypothetical 27.4 kd protein in hyr1 3'region.
CONTIG439	22395066_fl_2	2077	7739	798	266	1095	5.5(10)-111	Escherichia coli	b0763	[pn:24 aa signal peptide] [gn:moda]
CONTIG439	836680_fl_7	2078	7740	1008	336	1444	5.7(10)-148	Escherichia coli	b0767	[pn:hypothetical 36.3 kd protein in modc-bioa intergenic region] [gn:ybhc]
CONTIG439	22297577_fl_11	2079	7741	1257	419	730	2.6(10)-72	Bacillus subtilis	hutI	[pn:imidazolonepropionase] [gn:ee57b]
CONTIG439	10599075_fl_13	2080	7742	888	296	988	1.2(10)-99	Klebsiella aerogenes	P12380	histidine utilization repressor.
CONTIG439	34161325_fl_15	2081	7743	1581	527	1849	6.9(10)-191	Pseudomonas putida	A35251	histidine ammonia-lyase (ec 4.3.1.3)
CONTIG439	31820791_fl_19	2082	7744	1194	398	1163	3.3(10)-118	Escherichia coli	b0776	pseudomonas putida [pn:8-amino-7-oxononanoate synthase] [gn:biof]
CONTIG439	26845457_f2_29	2083	7745	1077	359	1508	9.5(10)-155	Escherichia coli	b0765	[pn:molybdenum transport atp-binding protein mode] [gn:mode]
CONTIG439	32441655_f2_30	2084	7746	354	118	202	2.2(10)-16	Escherichia coli	U27192	or:escherichia coli pn:modd gn:modd le:5360 re:6055 di:direct
CONTIG439	29398902_f2_36	2085	7747	1761	587	1861	3.7(10)-192	Bacillus subtilis	hutU	[pn:urocanate hydratase] [gn:ee57a]

CONTIG439	14945407_f2_46	2086	7748	1080	360	1602	1.0(10)-164	Escherichia coli	b0775	[pn:biotin synthetase] [gn:biob]
CONTIG439	22913166_f2_49	2087	7749	810	270	741	1.8(10)-73	Escherichia coli	b0777	[pn:biotin synthesis protein bioc] [gn:bioc]
CONTIG439	5334457_f3_56	2088	7750	717	239	713	1.7(10)-70	Escherichia coli	b0764	[pn:molybdenum transport system permease protein modb] [gn:modb]
CONTIG439	5164052_f3_59	2089	7751	270	90	159	8.4(10)-12	Escherichia coli	D90715	or:escherichia coli pn:putative molybdenum transport protein modd gn:modd le:4270 re:4410 di:direct sr:escherichia coli(strain:k12) dna, clone:kohara clone #180
CONTIG439	32033513_f3_63	2090	7752	999	333	643	4.2(10)-63	Klebsiella aerogenes	P19452	formiminoglutamase (ec 3.5.3.8) (formiminoglutamate hydrolase) (histidine utilization protein g) (fragment).
CONTIG439	14320833_f3_76	2091	7753	837	279	858	7.2(10)-86	Escherichia coli	b0778	[pn:dethiobiotin synthetase] [gn:biob]
CONTIG439	12582512_f3_77	2092	7754	351	117	364	8.5(10)-33	Escherichia coli	b0779	[pn:excision nuclease abc subunit b] [gn:uvrb]
CONTIG439	23531503_c1_78	2093	7755	828	276	275	4.2(10)-24	Escherichia coli	b3454	[pn:high-affinity branched-chain amino acid transport atp-binding] [gn:livf]
CONTIG439	25970953_c1_87	2094	7756	498	166	705	1.2(10)-69	Escherichia coli	b0773	[pn:hypothetical 17.1 kd protein in bioa 5'''region] [gn:ybbh]
CONTIG439	2945417_c1_101	2095	7757	1137	379	1440	1.5(10)-147	Escherichia coli	b0772	[pn:hypothetical protein in bioa 5'''region] [gn:ybhc]
CONTIG439	31306950_c2_119	2096	7758	1350	450	1866	1.1(10)-192	Escherichia coli	b0774	[pn:adenosylmethionine-8-amino-7-oxonanoate aminotransferase] [gn:bioa]
CONTIG439	16601510_c2_135	2097	7759	501	167	258	6.2(10)-22	Escherichia coli	b0772	[pn:hypothetical protein in bioa 5'''region] [gn:ybhc]
CONTIG439	31694127_c3_162	2098	7760	903	301	1093	9.0(10)-111	Escherichia coli	b0766	[pn:hypothetical 30.2 kd protein in mode-bioa intergenic region] [gn:ybha]
CONTIG44	29432965_f3_3	2099	7761	555	185	220	2.7(10)-17	Escherichia coli	b3966	[pn:vitamin b12 receptor precursor] [gn:btub]
CONTIG440	29298500_f1_1	2100	7762	258	86	400	2.3(10)-37	Escherichia coli	b3637	[pn:50s ribosomal subunit protein 128] [gn:rpmb]

CONTIG440	24814713_f1_28	2101	7763	775	259	257	3.5(10)-22	Haemophilus influenzae	HI0653	[pn:pir]
CONTIG440	23633567_f3_46	2102	7764	183	61	246	5.0(10)-21	Escherichia coli	b3636	[pn:50s ribosomal subunit protein 133] [gn:rpmg]
CONTIG440	4141380_f3_47	2103	7765	876	292	1324	3.0(10)-135	Escherichia coli	b3635	[pn:formamidopyrimidine-dna glycosylase] [gn:mutm]
CONTIG440	31525267_f3_67	2104	7766	1395	465	1853	2.6(10)-191	Escherichia coli	b3617	[pn:2-amino-3-ketobutyrate coenzyme a ligase] [gn:kb1]
CONTIG440	5127068_f3_68	2105	7767	1038	346	1563	1.3(10)-160	Escherichia coli	b3616	[pn:threonine 3-dehydrogenase] [gn:tdh]
CONTIG440	31901006_c1_81	2106	7768	1137	379	1636	2.6(10)-168	Escherichia coli	b3620	[pn:adp-heptose-lps heptosyltransferase ii] [gn:rfa]
CONTIG440	7290966_c1_82	2107	7769	987	329	1267	3.2(10)-129	Escherichia coli	b3621	[pn:lipopolysaccharide heptosyltransferase-1] [gn:rfa]
CONTIG440	19806531_c1_83	2108	7770	1113	371	164	1.3(10)-9	Helicobacter pylori	HP1191	[pn:adp-heptose-lps heptosyltransferase ii] [gn:rfa]
CONTIG440	25910910_c1_92	2109	7771	1086	362	716	8.0(10)-71	Escherichia coli	b3632	[pn:lipopolysaccharide core biosynthesis protein rfa] [gn:rfa]
CONTIG440	12892041_c1_95	2110	7772	1047	349	606	3.6(10)-59	Escherichia coli	b3615	[pn:hypothetical 40.5 kd protein in secb-tdh 5''' region] [gn:yibd]
CONTIG440	16893762_c1_96	2111	7773	1311	437	1727	5.9(10)-178	Escherichia coli	b3633	[pn:3-deoxy-d-manno-octulosonic acid transferase] [gn:kdtA]
CONTIG440	9979011_c2_99	2112	7774	1221	407	198	3.2(10)-13	Methanococcus jannaschii	MJ1059	[pn:capsular polysaccharide biosynthesis protein m]
CONTIG440	20603766_c2_103	2113	7775	975	325	1601	1.3(10)-164	Escherichia coli	b3619	[pn:adp-1-glycero-d-mannoheptose-6-epimerase] [gn:rfa]
CONTIG440	26594043_c2_110	2114	7776	1218	406	219	2.2(10)-16	Escherichia coli	b3622	[pn:rfa]
CONTIG440	23860312_c2_114	2115	7777	1152	384	192	1.3(10)-12	Methanococcus jannaschii	MJ1059	[pn:capsular polysaccharide biosynthesis protein m]
CONTIG440	4691452_c3_124	2116	7778	576	192	103	0.0038	Homo sapiens	Z34277	or:homo sapiens pn:mucin gn:muc5ac le:<1 re:>1431 di:direct
CONTIG440	32285666_c3_130	2117	7779	930	310	126	1.5(10)-5	Haemophilus influenzae	HI1578	[pn:glycosyl transferase] [gn:lgtd]
CONTIG440	2205040_c3_132	2118	7780	1161	387	224	1.8(10)-16	Escherichia coli	b3631	[pn:lipopolysaccharide core biosynthesis protein rfa] [gn:rfa]
CONTIG440	10672642_c3_136	2119	7781	849	283	637	1.8(10)-62	Haemophilus influenzae	HI0653	[pn:pir]



CONTIG440	14337783_c3_137	2120	7782	483	161	685	1.5(10)-67	Escherichia coli	b3634	[pn:lipopolysaccharide core biosynthesis protein kdtb] [gn:kdtb]
CONTIG441	31447255_fl_8	2121	7783	402	134	280	1.3(10)-24	Escherichia coli	b2824	[pn:hypothetical 13.5 kd protein in ppdc-ppdb intergenic region] [gn:ygdh]
CONTIG441	25645890_fl_15	2122	7784	2907	969	4170	0	Escherichia coli	b2821	[pn:protease iii precursor] [gn:ptr]
CONTIG441	19585327_fl_22	2123	7785	1338	446	1912	1.5(10)-197	Escherichia coli	b2817	[pn:hypothetical protein]
CONTIG441	33790961_f2_25	2124	7786	879	293	1386	8.0(10)-142	Escherichia coli	b2828	[pn:prolipoprotein diacylglycerol transferase] [gn:lg]
CONTIG441	24644068_f2_26	2125	7787	801	267	1401	2.1(10)-143	Escherichia coli	b2827	[pn:thymidylate synthetase] [gn:thya]
CONTIG441	34645790_f2_28	2126	7788	579	193	495	2.1(10)-47	Escherichia coli	b2825	[pn:prepilin peptidase dependent protein b precursor] [gn:ppdb]
CONTIG441	33828506_f2_36	2127	7789	1851	617	2175	2.0(10)-225	Escherichia coli	b2819	[pn:exonuclease v alpha-subunit] [gn:reed]
CONTIG441	477291_f3_40	2128	7790	360	120	541	2.7(10)-52	Escherichia coli	b2830	[pn:hypothetical protein] [gn:ygdh]
CONTIG441	4869633_f3_41	2129	7791	2259	753	3278	0	Escherichia coli	b2829	[pn:phosphoenolpyruvate-protein phosphotransferase] [gn:ptsp]
CONTIG441	281892_f3_46	2130	7792	516	172	410	2.1(10)-38	Escherichia coli	b2826	[pn:prepilin peptidase dependent protein a precursor] [gn:ppda]
CONTIG441	16182818_f3_48	2131	7793	336	112	239	2.7(10)-20	Escherichia coli	b2823	[pn:prepilin peptidase dependent protein c precursor] [gn:ppdc]
CONTIG441	1305455_f3_49	2132	7794	3387	1129	4626	0	Escherichia coli	b2822	[pn:exonuclease v subunit] [gn:recc]
CONTIG441	14883438_f3_51	2133	7795	3549	1183	4526	0	Escherichia coli	b2820	[pn:exonuclease v subunit] [gn:recc]
CONTIG441	23550057_c3_124	2134	7796	1353	451	2120	1.3(10)-219	Escherichia coli	b2818	[pn:amino-acid acetyltransferase] [gn:arga]
CONTIG442	35750680_fl_1	2135	7797	726	242	1115	4.2(10)-113	Escherichia coli	b2777	[pn:hypothetical protein] [gn:ygef]
CONTIG442	33463508_fl_10	2136	7798	1752	584	2661	6.2(10)-277	Escherichia coli	b2763	[pn:nadph hemoprotein alpha subunit] [gn:cys]
CONTIG442	7242681_fl_11	2137	7799	750	250	1236	6.2(10)-126	Escherichia coli	b2762	[pn:3'-phosphoadenosine 5'-phosphosulfate sulfotransferase] [gn:cys]

CONTIG442	6769537_f1_18	2138	7800	375	125	403	1.2(10)-37	Escherichia coli	b2748	[pn:hypothetical protein]
CONTIG442	25664010_f1_19	2139	7801	558	186	578	3.3(10)-56	Escherichia coli	b2746	[pn:hypothetical 16.9 kd protein in sure-cysc intergenic region] [gn:ygbg]
CONTIG442	19806956_f1_20	2140	7802	843	281	1240	2.3(10)-126	Escherichia coli	b2744	[pn:stationary-phase survival protein sure] [gn:sure]
CONTIG442	11760791_f1_21	2141	7803	199	67	109	3.0(10)-6	Escherichia coli	b2743	[pn:l-isoaspartyl protein carboxyl methyltransferase type ii]
CONTIG442	36017966_f2_22	2142	7804	642	214	274	5.5(10)-24	Escherichia coli	b3223	[pn:hypothetical protein] [gn:yhcj]
CONTIG442	14880191_f2_29	2143	7805	1812	604	2693	2.5(10)-280	Escherichia coli	b2764	[pn:nadph flavoprotein beta subunit] [gn:cysj]
CONTIG442	16109558_f2_41	2144	7806	651	217	876	8.9(10)-88	Escherichia coli	b2750	[pn:adenosine 5-phosphosulfate kinase] [gn:cysc]
CONTIG442	17052176_f2_42	2145	7807	810	270	1063	1.3(10)-107	Escherichia coli	b2747	[pn:hypothetical protein]
CONTIG442	17010938_f3_44	2146	7808	273	91	197	5.0(10)-15	Escherichia coli	b1101	[pn:pts system, glucose-specific iibc component] [gn:ptsg]
CONTIG442	16679686_f3_50	2147	7809	2397	799	150	7.2(10)-7	Bacillus subtilis	yesS	[pn:hypothetical protein]
CONTIG442	11753301_f3_57	2148	7810	1443	481	975	5.9(10)-107	Escherichia coli	b3368	[pn:siroheme synthase] [gn:cysg]
CONTIG442	3963443_f3_58	2149	7811	918	306	1473	4.7(10)-151	Escherichia coli	b2752	[pn:atp sulfurylase] [gn:cysd]
CONTIG442	4426068_f3_59	2150	7812	1434	478	2131	9.0(10)-221	Escherichia coli	b2751	[pn:atp sulfurylase] [gn:cysn]
CONTIG442	14119653_f3_61	2151	7813	348	116	407	4.4(10)-38	Escherichia coli	b2749	[pn:hypothetical protein in sure-cysc intergenic region] [gn:ygbg]
CONTIG442	16298917_f3_64	2152	7814	1080	360	1563	1.3(10)-160	Escherichia coli	b2745	[pn:hypothetical protein] [gn:ybgo]
CONTIG442	35596015_f3_65	2153	7815	363	121	453	5.9(10)-43	Escherichia coli	b2743	[pn:l-isoaspartyl protein carboxyl methyltransferase type ii]
CONTIG442	4018943_c2_107	2154	7816	1053	351	1448	2.2(10)-148	Escherichia coli	b2753	[pn:iap] [gn:iap]
CONTIG442	4142201_c2_120	2155	7817	1293	431	573	1.1(10)-55	Bacillus subtilis	yhaA	[pn:hypothetical protein]
CONTIG442	9866650_c2_121	2156	7818	1425	475	271	2.2(10)-21	Bacillus subtilis	mmr	[pn:methylenomycin a resistance protein]

CONTIG442	400675_c3_148	2157	7819	534	178	634	3.8(10)-62	Escherichia coli	b2765	[pn:hypothetical protein]
CONTIG443	17050011_fl_1	2158	7820	324	108	555	9.1(10)-54	Plasmid R478	L38824	or:plasmid r478 gn:tera lc:751 re:1776 di:direct sr:plasmid r478 dna nt:putative
CONTIG443	3932668_fl_13	2159	7821	306	102	97	3.1(10)-5	Escherichia coli	b1508	[pn:hipb protein] [gn:hipb]
CONTIG443	23870438_fl_15	2160	7822	828	276	93	0.065	Rattus norvegicus	S15674	cell surface glycoprotein ox47 precursor - rat
CONTIG443	25567885_fl_18	2161	7823	525	175	898	4.0(10)-90	Escherichia coli	b0264	[pn:insertion element is1 protein insb] [gn:insb_2]
CONTIG443	16914082_f2_23	2162	7824	588	196	749	2.5(10)-74	Plasmid R478	L38824	or:plasmid r478 gn:terb lc:1799 re:2254 di:direct sr:plasmid r478 dna nt:putative
CONTIG443	12268751_f2_25	2163	7825	633	211	429	2.1(10)-40	Bacillus subtilis	yceD	[pn:hypothetical protein]
CONTIG443	33632805_f2_26	2164	7826	1263	421	2036	1.1(10)-210	Plasmid R478	L38824	or:plasmid r478 gn:terf lc:5017 re:6258 di:direct sr:plasmid r478 dna nt:putative
CONTIG443	22676377_f2_38	2165	7827	1359	453	832	4.0(10)-83	Escherichia coli	b1507	[pn:hipa protein] [gn:hipa]
CONTIG443	6735393_f2_43	2166	7828	285	95	446	3.2(10)-42	Escherichia coli	b4294	[pn:insertion element is1f protein insa] [gn:insa_7]
CONTIG443	9822137_f3_46	2167	7829	1044	348	1530	4.4(10)-157	Plasmid R478	L38824	or:plasmid r478 gn:terc lc:2277 re:3317 di:direct sr:plasmid r478 dna nt:putative
CONTIG443	23547182_f3_47	2168	7830	606	202	549	4.0(10)-53	Bacillus subtilis	yceD	[pn:hypothetical protein]
CONTIG443	2911290_f3_65	2169	7831	234	78	370	3.7(10)-34	Escherichia coli	S40546	hypothetical protein - escherichia coli
CONTIG443	23947168_c1_88	2170	7832	2043	681	328	2.0(10)-26	Bacillus subtilis	yjcD	[pn:hypothetical protein]
CONTIG443	16125182_c2_94	2171	7833	2448	816	3998	0	Escherichia coli	P08504	transposase for transposon tn2501.
CONTIG443	31284383_c2_95	2172	7834	1026	342	1577	4.5(10)-162	Enterobacter aerogenes	U67194	or:enterobacter aerogenes pn:tnpa gn:tnpa lc:10303 re:11307 di:complement

CONTIG443	14885165_c2_97	2173	7835	297	99	390	2.7(10)-36	Escherichia coli	D93826	hypothetical 11k protein (insertion sequence is) -escherichia coli this protein is coded by the insertion sequence isl.
CONTIG443	22313533_c2_113	2174	7836	1083	361	100	0.027	Saccharomyces cerevisiae	P25040	hypothetical protein in ifm1 3'region (fragment).
CONTIG444	6341_fl_13	2175	7837	2628	876	4054	0	Escherichia coli	b3806	[pn:adenylate cyclase] [gn:cyaal]
CONTIG444	25886466_fl_14	2176	7838	888	296	1326	1.8(10)-135	Escherichia coli	b3809	[pn:diaminopimelate epimerase] [gn:dapf]
CONTIG444	10677291_fl_17	2177	7839	750	250	1051	2.5(10)-106	Escherichia coli	b3812	[pn:hypothetical 27.1 kd protein in xerc-uvrd intergenic region]
CONTIG444	14656965_fl_23	2178	7840	1023	341	1516	1.3(10)-155	Escherichia coli	b3816	[pn:magnesium and cobalt transport protein cora] [gn:cora]
CONTIG444	15097887_fl_26	2179	7841	924	308	1440	1.5(10)-147	Escherichia coli	b3821	[pn:phospholipase a1 precursor] [gn:plda]
CONTIG444	1368832_fl_27	2180	7842	1896	632	2840	6.7(10)-296	Escherichia coli	b3822	[pn:dna-dependent atpase, dna helicase] [gn:recq]
CONTIG444	36619627_fl_30	2181	7843	471	157	703	1.8(10)-69	Escherichia coli	b3825	[pn:lysophospholipase 12] [gn:pldb]
CONTIG444	11932643_f2_46	2182	7844	459	153	206	8.8(10)-17	Escherichia coli	P39166	hypothetical 7.2 kd protein in cyay-dapf intergenic region.
CONTIG444	14877083_f2_48	2183	7845	909	303	1308	1.5(10)-133	Escherichia coli	b3811	[pn:integrase/recombinase xerc] [gn:xerc]
CONTIG444	35158586_f2_50	2184	7846	2217	739	3493	0	Escherichia coli	b3813	[pn:dna helicase ii] [gn:uvrd]
CONTIG444	829026_f2_57	2185	7847	651	217	446	3.2(10)-42	Escherichia coli	b3823	[pn:hypothetical 13.3 kd protein in recq 3''' region] [gn:yigi]
CONTIG444	11875466_f3_72	2186	7848	723	241	992	4.5(10)-100	Escherichia coli	b3810	[pn:hypothetical 26.7 kd protein in dapf-xerc intergenic region]
CONTIG444	5198312_c1_89	2187	7849	645	215	547	6.5(10)-53	Escherichia coli	b3824	[pn:hypothetical 15.4 kd protein in recq-pldb intergenic region]
CONTIG444	22520812_c1_120	2188	7850	1254	418	1123	5.9(10)-114	Escherichia coli	b3803	[pn:uroporphyrinogen iii methylase] [gn:hmx]
CONTIG444	11744501_c2_126	2189	7851	555	185	727	5.5(10)-72	Escherichia coli	b3820	[pn:hypothetical 17.1 kd protein in rard-plda intergenic region] [gn:yigi]
CONTIG444	12109716_c2_127	2190	7852	909	303	1197	8.5(10)-122	Escherichia coli	b3819	[pn:rard protein] [gn:rard]

CONTIG444	12242841_c2_141	2191	7853	756	252	924	7.2(10)-93	Escherichia coli	b3804	[pn:uroporphyrinogen iii synthase] [gn:hemd]
CONTIG444	15735641_c3_169	2192	7854	360	120	431	1.3(10)-40	Escherichia coli	b3807	[pn:cyay protein] [gn:cyay]
CONTIG444	4728191_c3_173	2193	7855	966	322	1447	2.7(10)-148	Escherichia coli	b3805	[pn:porphobilinogen deaminase] [gn:hmc]
CONTIG444	16145837_c3_177	2194	7856	1224	408	1682	3.3(10)-173	Escherichia coli	b3802	[pn:hemy protein] [gn:hemy]
CONTIG444	6282962_c3_178	2195	7857	1224	408	189	2.1(10)-13	Erwinia herbicola	Q01334	hypothetical 29.9 kd protein in crte 3'region (orf3).
CONTIG445	23634781_fl_4	2196	7858	3081	1027	485	1.6(10)-52	Serratia marcescens	JC5568	[pn:serine proteinase h1, precursor] [gn:ssp-h1]
CONTIG445	10052331_fl_15	2197	7859	819	273	737	4.7(10)-73	Escherichia coli	b2805	[pn:l-fucose operon activator] [gn:fucl]
CONTIG445	24868825_fl_24	2198	7860	348	116	128	1.6(10)-8	Escherichia coli	b3004	[pn:hypothetical protein]
CONTIG445	32035208_f2_31	2199	7861	2013	671	1321	2.0(10)-186	Escherichia coli	b1023	[pn:hypothetical protein] [gn:yedr]
CONTIG445	9948592_f2_34	2200	7862	447	149	208	5.4(10)-17	Yersinia pestis	U22837	or:yersinia pestis pn:hms gn:hms le:7025 re:7492 di:direct nt:hypothetical and essential protein; pi 6.68; 17.5
CONTIG445	16613762_f2_39	2201	7863	1212	404	1574	9.5(10)-162	Escherichia coli	b3001	[pn:hypothetical protein]
CONTIG445	6410751_f3_44	2202	7864	2424	808	1529	5.5(10)-157	Escherichia coli	b1024	[pn:hypothetical protein] [gn:yods]
CONTIG445	24304207_f3_48	2203	7865	1434	478	1459	1.5(10)-149	Escherichia coli	b1022	[pn:hypothetical protein] [gn:yedq]
CONTIG445	441008_f3_51	2204	7866	537	179	91	0.049	Klebsiella pneumoniae	P06218	nitrogen regulation protein ntrb (ec 2.7.3.-)
CONTIG445	7241576_f3_53	2205	7867	981	327	1332	4.2(10)-136	Escherichia coli	b2989	[pn:hypothetical protein]
CONTIG445	25990684_c1_61	2206	7868	281	93	274	5.5(10)-24	Escherichia coli	b3005	[pn:biopolymer transport exbd protein] [gn:exbd]
CONTIG445	32604711_c1_81	2207	7869	531	177	91	0.03599	Neisseria gonorrhoeae	SI6613	[PN:opacity protein opaB]
CONTIG445	11854837_c1_85	2208	7870	645	215	198	6.2(10)-16	Haemophilus influenzae	HI0726	[pn:nitrate/nitrite response regulator protein] [gn:narp]

CONTIG445	33615888_c2_87	2209	7871	528	176	749	2.5(10)-74	Escherichia coli	b3002	[pn:hypothetical protein] [gn:yqha]
CONTIG445	14659377_c2_90	2210	7872	1677	559	746	5.2(10)-74	Escherichia coli	b1421	[pn:methyl-accepting chemotaxis protein iii] [gn:trg]
CONTIG445	4477253_c2_94	2211	7873	1872	624	2696	1.2(10)-280	Escherichia coli	b2988	[pn:glutathionylspermidine synthetase/amidase] [gn:gsp]
CONTIG445	24862907_c3_110	2212	7874	390	130	157	5.7(10)-11	Pseudomonas sp.	P18896	increased glyphosate resistance protein.
CONTIG446	14222077_fl_7	2213	7875	1062	354	1563	1.3(10)-160	Escherichia coli	b4269	[pn:hypothetical zinc-type alcohol dehydrogenase- like protein] [gn:yjgb]
CONTIG446	35711006_f2_24	2214	7876	2880	960	4657	0	Escherichia coli	b4258	[pn:valyl-trna synthetase] [gn:vals]
CONTIG446	475468_f2_25	2215	7877	597	199	422	1.1(10)-39	Escherichia coli	b4256	[pn:hypothetical protein]
CONTIG446	21992143_f3_40	2216	7878	603	201	708	5.5(10)-70	Escherichia coli	b4259	[pn:dna polymerase iii chi subunit] [gn:holec]
CONTIG446	16172193_f3_50	2217	7879	1572	524	2108	2.5(10)-218	Escherichia coli	b4263	[pn:hypothetical 54.3 kd protein in pepa-gntv intergenic region]
CONTIG446	36381892_f3_52	2218	7880	4503	1501	7200	0	Escherichia coli	b3212	[pn:glutamate synthase] [gn:gltb]
CONTIG446	5250318_f3_53	2219	7881	1428	476	2377	7.7(10)-247	Escherichia coli	b3213	[pn:glutamate synthase] [gn:gltd]
CONTIG446	29304682_c1_71	2220	7882	1665	555	1540	3.7(10)-158	Citrobacter freundii	P45510	dihydroxyacetone kinase (ec 2.7.1.29) (glycerone kinase).
CONTIG446	33682087_c2_90	2221	7883	1032	344	1515	1.7(10)-155	Escherichia coli	b3211	[pn:hypothetical 34.6 kd protein in arcB-gltB intergenic region]
CONTIG446	21954086_c2_91	2222	7884	948	316	181	5.0(10)-12	Pseudomonas aeruginosa	D86947	[de:pseudomonas aeruginosa gene for chemotactic transducer, completecds.] [pn:hydrophilic protein] [nt:orf1]
CONTIG446	24415936_c3_102	2223	7885	675	225	244	8.3(10)-21	Escherichia coli	b3219	[pn:hypothetical protein] [gn:yhcf]
CONTIG447	15125800_fl_1	2224	7886	540	180	801	7.7(10)-80	Escherichia coli	b0812	[pn:dna protection during starvation protein] [gn:dps]
CONTIG447	25604712_fl_3	2225	7887	741	247	1104	6.0(10)-112	Escherichia coli	b0809	[pn:glutamine transport atp-binding protein glnQ] [gn:glnQ]
CONTIG447	26285806_fl_4	2226	7888	2328	776	2857	1.1(10)-297	Escherichia coli	b0808	[pn:hypothetical protein]

CONTIG447	12586088_f1_6	2227	7889	300	100	285	3.7(10)-25	Escherichia coli	b0806	[pn:hypothetical protein]
CONTIG447	861407_f1_9	2228	7890	2583	861	1504	2.5(10)-154	Escherichia coli	b3144	[pn:hypothetical outer membrane usher protein in agal-mtr intergenic region] [gn:yrai]
CONTIG447	4488453_f1_11	2229	7891	513	171	197	7.9(10)-16	Escherichia coli	b4319	[pn:fimg protein precursor] [gn:fimg]
CONTIG447	13161462_f1_24	2230	7892	1224	408	1395	8.9(10)-143	Escherichia coli	b0793	[pn:hypothetical protein]
CONTIG447	25964387_f2_25	2231	7893	816	272	1210	3.6(10)-123	Escherichia coli	b0811	[pn:glutamine-binding periplasmic protein precursor] [gn:glnh]
CONTIG447	16853916_f2_26	2232	7894	729	243	744	8.5(10)-74	Escherichia coli	b0810	[pn:glutamine transport system permease protein glnp] [gn:glnp]
CONTIG447	36033312_f2_38	2233	7895	582	194	209	4.2(10)-17	Escherichia coli	b4314	[pn:type 1 fimbrial subunit] [gn:fima]
CONTIG447	3909688_f2_39	2234	7896	684	228	408	3.5(10)-38	Escherichia coli	b3143	[pn:hypothetical 25.7 kd fimbrial chaperone in agal- mtr intergeni] [gn:yrai]
CONTIG447	24494091_f2_46	2235	7897	1065	355	114	7.4(10)-5	Escherichia coli	b0942	[pn:hypothetical protein]
CONTIG447	15720380_f2_56	2236	7898	1035	345	1194	1.8(10)-121	Escherichia coli	b0795	[pn:hypothetical protein]
CONTIG447	26736057_f3_73	2237	7899	285	95	401	1.8(10)-37	Escherichia coli	b0803	[pn:hypothetical 9.8 kd protein in ding/rarb 3'''region] [gn:ybii]
CONTIG447	14650256_f3_74	2238	7900	264	88	193	2.1(10)-15	Escherichia coli	b0802	[pn:hypothetical 8.6 kd protein in ding/rarb 3'''region] [gn:ybii]
CONTIG447	23437800_f3_89	2239	7901	696	232	874	1.3(10)-87	Escherichia coli	b0796	[pn:hypothetical transcriptional regulator in moae-rhle intergenic region] [gn:ybih]
CONTIG447	24900957_f3_92	2240	7902	1755	585	2686	1.3(10)-279	Escherichia coli	b0794	[pn:hypothetical protein]
CONTIG447	32713215_c2_138	2241	7903	1407	469	1801	8.5(10)-186	Escherichia coli	b0797	[pn:putative atp-dependent ma helicase] [gn:rhle]
CONTIG447	5177157_c3_156	2242	7904	309	103	378	5.2(10)-35	Entamoeba histolytica	Y14328	[PN:3E1 protein] [DE:Entamoeba histolytica mRNA for 3E1 protein.] [LE:32] [RE:418] [DI:direct]

CONTIG447	29933406_c3_157	2243	7905	435	145	95	0.05299	Caenorhabditis elegans	U39852	or:caenorhabditis elegans gn:k10c2.1 le:join(2542 re:2988,309 di:complement sr:caenorhabditis elegans strain=bristol n2 nt:coded for by c. elegans cdna cm20c9; coded for by
CONTIG447	136417_c3_164	2244	7906	2256	752	3206	0	Escherichia coli	b0799	[pn:probable atp-dependent helicase ding] [gn:ding]
CONTIG447	15829811_c3_172	2245	7907	996	332	1122	7.5(10)-114	Escherichia coli	b0800	[pn:hypothetical 35.0 kd protein in ding/rarb 3'''region] [gn:ybib]
CONTIG447	26206586_c3_174	2246	7908	1026	342	1341	4.7(10)-137	Escherichia coli	b0807	[pn:hypothetical protein]
CONTIG448	2510268_fl_7	2247	7909	444	148	137	1.8(10)-9	Escherichia coli	b4347	[pn:hypothetical 14.6 kd protein in mcrb-hsds intergenic region] [gn:yjiw]
CONTIG448	26251502_fl_14	2248	7910	735	245	96	0.01499	Escherichia coli	b4365	[pn:hypothetical 27.0 kd protein in dnat-hold intergenic region]
CONTIG448	35807665_fl_15	2249	7911	771	257	385	9.5(10)-36	Escherichia coli	b0939	[pn:hypothetical protein]
CONTIG448	24408336_fl_21	2250	7912	1110	370	155	9.6(10)-11	Escherichia coli	b4314	[pn:type 1 fimbrial subunit] [gn:fima]
CONTIG448	17007156_fl_25	2251	7913	1209	403	97	0.05899	Helicobacter pylori	HP1079	[pn:h]
CONTIG448	25432962_f2_37	2252	7914	498	166	164	2.5(10)-12	Vibrio cholerae	S81006	or:vibrio cholerae pn:hcp gn:hcp le:690 re:1208 di:direct sr:vibrio cholerae o17 nt:28 kda secreted hydrophilic protein; this sequence
CONTIG448	12766056_f2_42	2253	7915	645	215	193	2.1(10)-15	Escherichia coli	b4314	[pn:type 1 fimbrial subunit] [gn:fima]
CONTIG448	32599061_f2_45	2254	7916	693	231	240	2.2(10)-20	Escherichia coli	b0943	[pn:hypothetical protein]
CONTIG448	473413_f2_50	2255	7917	753	251	481	6.4(10)-46	Escherichia coli	b2127	[pn:hypothetical transcriptional regulator in molr-bglx intergenic region] [gn:yehv]
CONTIG448	31250052_f2_51	2256	7918	474	158	354	1.8(10)-32	Escherichia coli	b1970	[pn:hypothetical protein]
CONTIG448	7245763_f2_52	2257	7919	978	326	405	7.2(10)-38	Escherichia coli	b0707	[pn:hypothetical 20.2 kd protein in phrb 5'''region] [gn:ybga]
CONTIG448	5963256_f3_72	2258	7920	252	84	96	0.0004	Volvox carteri	S22697	extensin - volvox carteri (fragment)



CONTIG448	36568816_f3_74	2259	7921	2598	866	1593	9.3(10)-164	Escherichia coli	b0940	[pn:hypothetical protein]
CONTIG448	4494031_c1_99	2260	7922	783	261	644	3.3(10)-63	Bacillus subtilis	yddR	[pn:hypothetical protein]
CONTIG448	1055337_c1_127	2261	7923	504	168	349	6.2(10)-32	Escherichia coli	b3657	[pn:hypothetical 51.0 kd protein in glts-selc intergenic region] [gn:yicj]
CONTIG448	24659636_c2_131	2262	7924	1257	419	180	3.7(10)-11	Escherichia coli	P22519	colicin v secretion protein cvaa.
CONTIG448	26758592_c2_133	2263	7925	1005	335	146	8.3(10)-8	Haemophilus influenzae	H11052	[pn:arac-like transcription regulator]
CONTIG448	21960077_c2_149	2264	7926	855	285	274	5.5(10)-24	Escherichia coli	b2847	[pn:hypothetical protein]
CONTIG448	6132827_c2_150	2265	7927	990	330	152	1.8(10)-8	Bacillus subtilis	yfiF	[pn:hypothetical protein]
CONTIG448	22272552_c2_151	2266	7928	492	164	231	2.0(10)-19	Escherichia coli	b4012	[pn:hypothetical 16.4 kd protein in rrfc-meta intergenic region]
CONTIG448	16135200_c2_155	2267	7929	1161	387	1062	1.7(10)-107	Escherichia coli	b3657	[pn:hypothetical 51.0 kd protein in glts-selc intergenic region] [gn:yicj]
CONTIG448	2506660_c3_162	2268	7930	1200	400	374	4.0(10)-34	Bacillus subtilis	yknV	[pn:hypothetical protein]
CONTIG448	6929702_c3_177	2269	7931	540	180	614	5.0(10)-60	Escherichia coli	b4149	[pn:hypothetical 19.9 kd protein in sugc-ampc intergenic region]
CONTIG448	33334717_c3_184	2270	7932	492	164	123	5.5(10)-8	Escherichia coli	b2848	[pn:hypothetical protein]
CONTIG448	9900640_c3_190	2271	7933	435	145	541	2.7(10)-52	Escherichia coli	b3657	[pn:hypothetical 51.0 kd protein in glts-selc intergenic region] [gn:yicj]
CONTIG448	34582963_c3_191	2272	7934	2379	793	3661	0	Escherichia coli	b3656	[pn:hypothetical 88.1 kd protein in glts-selc intergenic region] [gn:yicj]
CONTIG449	14843818_f2_51	2273	7935	1017	339	921	1.5(10)-92	Escherichia coli	b2321	[pn:div protein] [gn:div]
CONTIG449	7235901_f2_77	2274	7936	1017	339	818	9.4(10)-126	Escherichia coli	b2344	[pn:long-chain fatty acid transport protein precursor] [gn:fadI]
CONTIG449	16065700_f3_78	2275	7937	183	61	110	1.3(10)-6	Escherichia coli	D90862	or:escherichia coli pn:glycine-rich cell wall structural protein gn:dedd le:14996 re:15160 di:direct sr:escherichia coli (strain:k12) dna, clone lib:kohara lambda minise nt:similar to [swissprot accession number p27483]

CONTIG449	16285206_f3_89	2276	7938	303	101	95	0.00025	Rattus norvegicus	S24169	mucin - rat
CONTIG449	6431430_f3_95	2277	7939	2172	724	2644	3.8(10)-275	Escherichia coli	b2324	[pn:hypothetical protein]
CONTIG449	24415941_f3_102	2278	7940	555	185	900	2.5(10)-90	Escherichia coli	b2331	[pn:hypothetical protein]
CONTIG449	30729686_c1_117	2279	7941	501	167	745	6.7(10)-74	Escherichia coli	b2340	[pn:hypothetical protein]
CONTIG449	111308_c1_121	2280	7942	822	274	1148	1.3(10)-116	Escherichia coli	b2327	[pn:hypothetical 28.6 kd protein in mepa 5'''region] [gn:yfca]
CONTIG449	19558455_c1_123	2281	7943	366	122	369	4.7(10)-34	Escherichia coli	b2325	[pn:hypothetical protein]
CONTIG449	15882827_c1_127	2282	7944	1218	406	1807	2.0(10)-186	Escherichia coli	b2323	[pn:3-oxoacyl-acyl-carrier-protein synthase i] [gn:fabb]
CONTIG449	897890_c1_138	2283	7945	1002	334	1330	6.9(10)-136	Escherichia coli	b2316	[pn:acetyl-coa carboxylase beta subunit] [gn:accd]
CONTIG449	21913887_c2_142	2284	7946	372	124	96	0.00025	Escherichia coli	D90864	or:escherichia coli pn:mitochondrial trifunctional enzyme beta subunit gn:hadhb le:13075 re:14127 di:complement sr:escherichia coli (strain:k12) dna, clone_jib:kohara lambda minise nt:similar to [swissprot accession number [pn:hypothetical protein]
CONTIG449	14665927_c2_145	2285	7947	2151	717	2730	3.0(10)-284	Escherichia coli	b2341	[pn:hypothetical protein]
CONTIG449	16850312_c2_148	2286	7948	216	72	174	1.2(10)-12	Escherichia coli	b2330	[pn:hypothetical adenine-specific methylase in aroc 3'''region] [gn:yfcb]
CONTIG449	22048513_c2_149	2287	7949	975	325	1123	5.9(10)-114	Escherichia coli	b2328	[pn:penicillin-insensitive murein endopeptidase precursor] [gn:mepa]
CONTIG449	6767933_c2_162	2288	7950	912	304	1251	1.6(10)-127	Escherichia coli	b2318	[pn:pseudouridylylate synthase i] [gn:trua]
CONTIG449	4588311_c2_166	2289	7951	1281	427	1748	3.5(10)-180	Escherichia coli	b2315	[pn:folypolyglutamate synthase] [gn:fole]
CONTIG449	23712758_c3_168	2290	7952	288	96	411	1.7(10)-38	Escherichia coli	b2343	[pn:hypothetical protein]
CONTIG449	24088515_c3_169	2291	7953	1344	448	1984	3.3(10)-205	Escherichia coli	b2342	[pn:hypothetical protein]

CONTIG449	14976381_c3_172	2292	7954	1023	341	1544	1.3(10)-158	Escherichia coli	b2330	[pn:hypothetical adenine-specific methylase in aroc 3'''region] [gn:yfcb]
CONTIG449	3411530_c3_173	2293	7955	1146	382	1557	6.0(10)-160	Escherichia coli	b2329	[pn:chorismate synthase] [gn:aroc]
CONTIG449	10751892_c3_176	2294	7956	540	180	134	7.0(10)-9	Escherichia coli	D90863	or:escherichia coli le:14845 re:15579 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [swissprot accession number]
CONTIG449	15085211_c3_177	2295	7957	588	196	914	8.3(10)-92	Escherichia coli	b2326	[pn:hypothetical protein]
CONTIG449	33470092_c3_183	2296	7958	1239	413	1402	1.6(10)-143	Escherichia coli	b2322	[pn:hypothetical protein]
CONTIG449	2348128_c3_185	2297	7959	1167	389	1630	1.1(10)-167	Escherichia coli	b2320	[pn:erythronate-4-phosphate dehydrogenase] [gn:pdxb]
CONTIG449	10972782_c3_186	2298	7960	1020	340	1386	8.0(10)-142	Escherichia coli	b2319	[pn:-l protein] [gn:usg]
CONTIG449	24037543_c3_187	2299	7961	672	224	756	4.5(10)-75	Escherichia coli	b2317	[pn:deda protein] [gn:deda]
CONTIG449	36588465_c3_190	2300	7962	339	113	251	1.5(10)-21	Escherichia coli	b2314	[pn:dedd protein] [gn:dedd]
CONTIG45	20132828_f3_3	2301	7963	771	257	301	8.0(10)-31	Escherichia coli	b2412	[pn:hypothetical protein] [gn:zipa]
CONTIG450	16897627_f1_1	2302	7964	813	271	1166	1.6(10)-118	Escherichia coli	b4364	[pn:hypothetical 30.5 kd protein in dna1-hold intergenic region]
CONTIG450	24067318_f1_12	2303	7965	1479	493	2489	1.0(10)-258	Escherichia coli	Z37980	or:escherichia coli pn:5-carboxy-2-hydroxymuconate semialdehyde gn:hpaec le:2137 re:3603 di:direct
CONTIG450	32052067_f1_16	2304	7966	330	110	90	0.0015	Zea mays	U28017	or:zea mays pn:globulin1 gn:gbl1 le:join(421 re:922,1010 di:direct sr:maize nt:allele gbl1-hb; a null allele caused by the
CONTIG450	35713343_f1_30	2305	7967	342	114	344	2.1(10)-31	Escherichia coli	b4353	[pn:hypothetical 7.7 kd protein in mrr-tsr intergenic region] [gn:yjix]
CONTIG450	5897876_f2_34	2306	7968	489	163	320	7.2(10)-29	Bacillus subtilis	ykmA	[pn:hypothetical protein]

CONTIG450	12714138_f2_35	2307	7969	444	148	593	8.5(10)-58	Acinetobacter calcoaceticus	Y09102	or:acinetobacter calcoaceticus pn:unknown protein le:621 re:1052 di:direct nt:orf2
CONTIG450	34469090_f2_36	2308	7970	756	252	1072	1.5(10)-108	Escherichia coli	b4361	[pn:dna replication protein dnac] [gn:dnac]
CONTIG450	24640902_f2_37	2309	7971	489	163	348	7.9(10)-32	Escherichia coli	b4360	[pn:hypothetical 17.5 kd protein in mdob-dnac intergenic region] [gn:yjia]
CONTIG450	22861262_f2_39	2310	7972	531	177	113	6.2(10)-7	Bacillus subtilis	yvbK	[pn:hypothetical protein]
CONTIG450	31835762_f2_43	2311	7973	1314	438	1646	2.2(10)-169	Escherichia coli	Z37980	or:escherichia coli pn:5-oxo-1,2,5- tricarboxylic-3-penten acid gn:hpag le:851 re:2140 di:direct
CONTIG450	4120187_f2_48	2312	7974	819	273	1318	1.3(10)-134	Escherichia coli	AF036583	[de:escherichia coli 2-oxo-hept-4-ene- 1,7-dioate hydratase (hpeg) gene,complete cds.] [pn:2-oxo-hept- 4-ene-1,7-dioate hydratase] [gn:hpeg] [nt:ohed hydratase]
CONTIG450	4891633_f2_59	2313	7975	540	180	764	6.5(10)-76	Escherichia coli	C55349	4-hydroxyphenylacetate 3- monooxygenase (ec 1.14.13.3) smallchain - escherichia coli (atcc 11105)
CONTIG450	2456431_f2_61	2314	7976	2319	773	2860	5.0(10)-298	Escherichia coli	b4354	[pn:hypothetical 77.9 kd protein in mirr-lsr intergenic region] [gn:yjiv]
CONTIG450	24632126_f2_62	2315	7977	699	233	721	2.3(10)-71	Escherichia coli	b4352	[pn:f2] [gn:yjia]
CONTIG450	34156635_f3_65	2316	7978	588	196	500	6.2(10)-48	Escherichia coli	b4363	[pn:p14 protein] [gn:yjib]
CONTIG450	3955313_f3_67	2317	7979	555	185	677	1.1(10)-66	Escherichia coli	b4362	[pn:primosomal protein i] [gn:dnat]
CONTIG450	31906558_f3_69	2318	7980	2334	778	3278	0	Escherichia coli	b4359	[pn:phosphoglycerol transferase i] [gn:mdob]
CONTIG450	35820818_f3_79	2319	7981	1140	380	1460	1.2(10)-149	Escherichia coli	Z37980	or:escherichia coli pn:homoproteocatechuate dioxygenase gn:hpac le:3605 re:4456 di:direct
CONTIG450	3956693_f3_80	2320	7982	390	130	572	1.5(10)-55	Escherichia coli	S10709	5-carboxymethyl-2- hydroxymuconate isomerase - escherichiacoli

CONTIG450	12776580_f3_82	2321	7983	1017	339	703	1.8(10)-69	Escherichia coli	b2245	[pn:hypothetical protein]
CONTIG450	6066957_f3_83	2322	7984	1365	455	1480	8.8(10)-152	Escherichia coli	Z37980	or:escherichia coli pn:hypothetical 4-hydroxyphenylacetate permease gn:hpax le:6734 re:8110 di:direct
CONTIG450	4035443_f3_84	2323	7985	942	314	1372	2.3(10)-140	Escherichia coli	Z37980	or:escherichia coli pn:regulator of the 4hpa-hydroxylase operon gn:hpaa le:8120 re:9007 di:direct
CONTIG450	24737630_f3_85	2324	7986	1686	562	2653	4.4(10)-276	Escherichia coli	B55349	4-hydroxyphenylacetate 3-monoxygenase (ec 1.14.13.3) largechain - escherichia coli (atcc 11105)
CONTIG450	3228125_c1_108	2325	7987	1704	568	141	2.1(10)-8	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:<1 re:588 di:direct
CONTIG450	12698757_c1_111	2326	7988	1788	596	1893	1.5(10)-195	Escherichia coli	b4355	[pn:methyl-accepting chemotaxis protein i] [gn:tsr]
CONTIG450	24901515_c2_138	2327	7989	456	152	617	2.5(10)-60	Escherichia coli	Q07095	homoprotocatechuate degradative operon repressor.
CONTIG451	5911250_f1_1	2328	7990	825	275	1103	7.7(10)-112	Escherichia coli	b3528	[pn:c4-dicarboxylate transport protein] [gn:dcta]
CONTIG451	21542930_f1_2	2329	7991	1503	501	1973	5.0(10)-204	Escherichia coli	b3527	[pn:53.1 kd protein in kdkg-dcta intergenic region precursor] [gn:yhjj]
CONTIG451	25986526_f1_5	2330	7992	822	274	998	1.0(10)-100	Escherichia coli	b3525	[pn:hypothetical 29.7 kd protein in tref-kdkg intergenic region]
CONTIG451	25665877_f1_11	2331	7993	762	254	360	4.2(10)-33	Bacillus subtilis	fabG	[pn:3-oxoacyl-acyl-carrier protein reductase] [gn:ylpf]
CONTIG451	10272250_f1_12	2332	7994	687	229	481	6.4(10)-46	Escherichia coli	b3520	[pn:hypothetical transcriptional regulator in tref- kdkg intergenic region] [gn:yhjb]
CONTIG451	207877_f1_21	2333	7995	2229	743	3076	0	Escherichia coli	b3498	[pn:oligopeptidase a] [gn:prlc]
CONTIG451	5328530_f1_22	2334	7996	462	154	551	2.3(10)-53	Escherichia coli	b3494	[pn:hypothetical 13.0 kd protein in pit-uspa intergenic region] [gn:yhio]
CONTIG451	5864762_f1_26	2335	7997	1287	429	1518	8.1(10)-156	Escherichia coli	b3492	[pn:hypothetical 43.8 kd protein in rhsb-pit intergenic region] [gn:yhin]
CONTIG451	7166656_f2_54	2336	7998	861	287	1222	1.8(10)-124	Escherichia coli	b3497	[pn:hypothetical protein in uspa-prlc intergenic region] [gn:yhiq]

CONTIG451	4338393_f2_60	2337	7999	852	284	286	2.8(10)-25	Thiobacillus ferrooxidans	AF032884	[de:thiobacillus ferrooxidans n-acetylglucosamine-1-phosphateuridytransferase (glmu) gene, partial cds; glucosamine synthase(glms)"] [pn:transposition complex] [gn:tnsa]
CONTIG451	4958318_f2_61	2338	8000	945	315	292	3.2(10)-25	Thiobacillus ferrooxidans	AF032884	[de:thiobacillus ferrooxidans n-acetylglucosamine-1-phosphateuridytransferase (glmu) gene, partial cds; glucosamine synthase(glms) and recg (recg) genes, complete cds; and transposon tn5468,complete sequence"] [pn:tnsc]
CONTIG451	53382_f3_70	2339	8001	2085	695	2730	3.0(10)-284	Escherichia coli	b3524	[pn:hypothetical 75.1 kd protein in tref-kdkg intergenic region]
CONTIG451	26601457_f3_95	2340	8002	1701	567	466	6.5(10)-44	Thiobacillus ferrooxidans	AF032884	[de:thiobacillus ferrooxidans n-acetylglucosamine-1-phosphateuridytransferase (glmu) gene, partial cds; glucosamine synthase(glms) and recg (recg) genes, complete cds; and transposon tn5468,complete"] [pn:tnsb] [gn:tnsb]
CONTIG451	3157016_c1_104	2341	8003	1506	502	2144	3.7(10)-222	Escherichia coli	b3493	[pn:pita] [gn:pita]
CONTIG451	3055383_c1_109	2342	8004	1443	481	2148	1.3(10)-222	Escherichia coli	b3500	[pn:glutathione oxidoreductase] [gn:gor]
CONTIG451	3939063_c1_120	2343	8005	1332	444	1671	5.0(10)-172	Escherichia coli	b3523	[pn:hypothetical metabolite transport protein in tref-kdkg intergenic region] [gn:yhje]
CONTIG451	23714693_c2_132	2344	8006	453	151	724	1.1(10)-71	Escherichia coli	b3495	[pn:universal stress protein a] [gn:uspa]
CONTIG451	35678462_c2_140	2345	8007	1707	569	2503	3.3(10)-260	Escherichia coli	b3519	[pn:probable cytoplasmic trehalase] [gn:tref]
CONTIG451	3242338_c2_144	2346	8008	351	117	141	6.7(10)-10	Salmonella typhimurium	X67137	or:salmonella typhimurium pn:gp19 protein gn:gene 19 le:578 re:1018 di:direct
CONTIG451	11034506_c2_148	2347	8009	675	225	643	4.2(10)-63	Escherichia coli	b3522	[pn:hypothetical 37.9 kd protein in tref-kdkg intergenic region]

CONTIG451	598951_c2_149	2348	8010	468	156	559	3.5(10)-54	Escherichia coli	b3522	[pn:hypothetical 37.9 kd protein in tref-kdkg intergenic region]
CONTIG451	24333507_c3_174	2349	8011	882	294	1353	2.5(10)-138	Escherichia coli	b3499	[pn:hypothetical 31.9 kd protein in prlc-gor intergenic region] [gn:yhir]
CONTIG451	23594561_c3_182	2350	8012	573	191	156	1.8(10)-11	Bacillus subtilis	ynaD	[pn:hypothetical protein]
CONTIG451	3991556_c3_184	2351	8013	909	303	1012	3.3(10)-102	Escherichia coli	b3521	[pn:hypothetical transcriptional regulator in tref- kdkg intergenic region] [gn:yhjc]
CONTIG451	24254075_c3_191	2352	8014	990	330	1340	6.0(10)-137	Escherichia coli	b3526	[pn:2-dehydro-3-deoxygluconokinase] [gn:kdkg]
CONTIG452	21900187_fl_9	2353	8015	318	106	1115	1.1(10)-6	Pseudomonas aeruginosa	JQ0133	hypothetical 26.4k protein - pseudomonas aeruginosa
CONTIG452	11020661_f2_35	2354	8016	702	234	746	5.2(10)-74	Escherichia coli	b0396	[pn:araj protein precursor] [gn:araj]
CONTIG452	4870402_f2_51	2355	8017	909	303	314	3.2(10)-28	Escherichia coli	b3521	[pn:hypothetical transcriptional regulator in tref- kdkg intergenic region] [gn:yhjc]
CONTIG452	34648408_f2_52	2356	8018	447	149	642	5.5(10)-63	Escherichia coli	b1482	[pn:osmotically inducible protein c] [gn:osmc]
CONTIG452	14636666_f2_54	2357	8019	795	265	568	3.7(10)-55	Bacillus subtilis	ykvO	[pn:hypothetical protein]
CONTIG452	12163286_f2_62	2358	8020	378	126	107	2.3(10)-5	Xenopus laevis	P17437	skin secretory protein xp2 precursor (apeg protein).
CONTIG452	36565837_f2_68	2359	8021	1506	502	104	0.0066	Bacillus subtilis	y4pA	[pn:hypothetical protein]
CONTIG452	36429028_f2_69	2360	8022	1215	405	691	3.6(10)-68	Escherichia coli	b1163	[pn:hypothetical protein]
CONTIG452	1306533_f3_73	2361	8023	513	171	143	4.2(10)-10	Escherichia coli	b4178	[pn:hypothetical 15.6 kd protein in pura-vacb intergenic region] [gn:yjeb]
CONTIG452	35290831_c1_106	2362	8024	1536	512	947	2.7(10)-95	Escherichia coli	b1485	[pn:hypothetical protein]
CONTIG452	17053875_c1_109	2363	8025	987	329	1160	7.0(10)-118	Escherichia coli	b1483	[pn:hypothetical protein]
CONTIG452	5292305_c1_111	2364	8026	921	307	404	9.1(10)-38	Escherichia coli	b3521	[pn:hypothetical transcriptional regulator in tref- kdkg intergenic region] [gn:yhjc]

CONTIG452	14945160_c1_115	2365	8027	1785	595	2635	3.5(10)-274	Escherichia coli	b1479	[pn:nad-linked malic enzyme] [gn:sfca]
CONTIG452	34413433_c1_116	2366	8028	1716	572	455	4.9(10)-52	Bacillus subtilis	ydif	[pn:hypothetical protein]
CONTIG452	22010303_c2_128	2367	8029	1050	350	1476	2.2(10)-151	Escherichia coli	b1486	[pn:hypothetical protein]
CONTIG452	35836461_c3_147	2368	8030	867	289	126	1.1(10)-5	Escherichia coli	b1853	[pn:hypothetical 32.0 kd protein in pyka-zwf intergenic region] [gn:yebk]
CONTIG452	16692842_c3_148	2369	8031	585	195	662	4.2(10)-65	Escherichia coli	b1488	[pn:hypothetical protein]
CONTIG452	12386275_c3_149	2370	8032	1548	516	2261	1.5(10)-234	Escherichia coli	b1487	[pn:hypothetical protein]
CONTIG452	32656630_c3_153	2371	8033	1044	348	1241	1.8(10)-126	Escherichia coli	b1484	[pn:hypothetical protein]
CONTIG452	5953808_c3_161	2372	8034	930	310	94	0.03699	Bacillus subtilis	yobT	[pn:hypothetical protein]
CONTIG452	41502_c3_163	2373	8035	267	89	307	1.7(10)-27	Escherichia coli	b1481	[pn:hypothetical protein]
CONTIG452	22464212_c3_167	2374	8036	1023	341	1349	6.7(10)-138	Escherichia coli	b1478	[pn:hypothetical protein]
CONTIG452	36125916_c3_168	2375	8037	1221	407	558	4.4(10)-54	Mycobacterium tuberculosis	Z96073	dehydrogenase, mcy1619.02, mcy1619.02, probable dehydrogenase, len
CONTIG453	22132932_f3_71	2376	8038	2112	704	1761	1.5(10)-181	Escherichia coli	b0779	[pn:excision nuclease abc subunit b] [gn:uvrb]
CONTIG453	35335183_f3_75	2377	8039	2709	903	133	6.7(10)-5	Archaeoglobus fulgidus	H69378	[pn:purine ntpase homolog]
CONTIG453	11722916_c1_80	2378	8040	441	147	293	2.5(10)-25	Rhizobium sp.	P50360	hypothetical 29.3 kd protein in region 2 of sym plasmid (no1265).
CONTIG453	2047880_c1_81	2379	8041	1050	350	1581	1.7(10)-162	Yersinia pestis	AF053945	[de:yersinia pestis plasmid ppcpl, complete plasmid sequence.] [pn:transposase]
CONTIG453	5286516_c1_88	2380	8042	243	81	283	6.0(10)-25	Salmonella dublin	S22685	vacc protein - salmonella dublin virulence plasmid
CONTIG453	2932082_c1_90	2381	8043	762	254	391	2.2(10)-36	Saccharomyces cerevisiae	P40586	hypothetical 27.4 kd protein in hyr1 3'region.
CONTIG453	4726577_c1_91	2382	8044	267	89	145	2.6(10)-10	Escherichia coli	b1892	[pn:flagellar transcriptional activator flhA] [gn:flhA]



CONTIG453	10744011_c1_95	2383	8045	1113	371	1628	1.8(10)-167	Escherichia coli	b0356	[pn:alcohol--acetaldehyde dehydrogenase] [gn:adhc]
CONTIG453	33644826_c1_97	2384	8046	1362	454	91	0.53	Helicobacter pylori	HP0870	[pn:flagellar hook flge] [gn:flge]
CONTIG453	26255200_c1_98	2385	8047	372	124	203	1.8(10)-16	Haemophilus influenzae	HI0184	[pn:hypothetical protein]
CONTIG453	24306507_c2_130	2386	8048	2655	885	1136	5.4(10)-163	Bacillus subtilis	uvrA	[pn:exonuclease abc]
CONTIG453	33697188_c2_132	2387	8049	279	93	412	1.3(10)-38	Escherichia coli	b0357	[pn:hypothetical protein]
CONTIG453	22672286_c3_135	2388	8050	804	268	1260	1.8(10)-128	Yersinia pestis	AF053945	[de:yersinia pestis plasmid ppcpl, complete plasmid sequence.] [pn:transposase]
CONTIG453	24025381_c3_144	2389	8051	219	73	271	1.1(10)-23	Salmonella dublin	S22686	vagD protein - salmonella dublin virulence plasmid
CONTIG453	10727336_c3_147	2390	8052	288	96	249	2.3(10)-21	Escherichia coli	b0357	[pn:hypothetical protein]
CONTIG453	19767826_c3_149	2391	8053	621	207	683	2.5(10)-67	Escherichia coli	b0355	[pn:hypothetical protein] [gn:yaim]
CONTIG454	15828957_f1_4	2392	8054	633	211	731	2.1(10)-72	Escherichia coli	b3249	[pn:rod shape-determining protein mred] [gn:mred]
CONTIG454	3984837_f1_5	2393	8055	603	201	864	1.7(10)-86	Escherichia coli	b3248	[pn:hypothetical 21.5 kd protein in cafa-mred intergenic region] [gn:yhde]
CONTIG454	24627202_f1_10	2394	8056	1446	482	2135	3.3(10)-221	Escherichia coli	b3244	[pn:tlld protein] [gn:tlld]
CONTIG454	10179131_f1_11	2395	8057	231	77	366	9.8(10)-34	Escherichia coli	b3242	[pn:hypothetical protein] [gn:yhcr]
CONTIG454	36042040_f1_14	2396	8058	2031	677	2286	3.3(10)-237	Escherichia coli	b3240	[pn:hypothetical 73.6 kd protein in argi-cafa intergenic region] [gn:yhcp]
CONTIG454	23614003_f1_17	2397	8059	963	321	1415	6.7(10)-145	Escherichia coli	b3236	[pn:malate dehydrogenase] [gn:mdh]
CONTIG454	13087756_f2_33	2398	8060	1020	340	1207	7.4(10)-123	Escherichia coli	b3250	[pn:rod shape-determining protein mrec] [gn:mrec]
CONTIG454	14453433_f2_35	2399	8061	1506	502	2182	3.6(10)-226	Escherichia coli	b3247	[pn:cytoplasmic axial filament protein] [gn:cafa]
CONTIG454	16026457_f2_36	2400	8062	3849	1283	3981	0	Escherichia coli	b3245	[pn:hypothetical 107.7 kd protein in argi-cafa intergenic region]

CONTIG454	30208260_f2_41	2401	8063	960	320	1415	6.7(10)-145	Escherichia coli	b3241	[pn:hypothetical 34.8 kd protein in argi-cafa intergenic region] [gn:yhcq]
CONTIG454	15647540_f3_55	2402	8064	1983	661	2669	8.8(10)-278	Escherichia coli	b3252	[pn:hypothetical 73.3 kd protein in mreB-accb intergenic region] [gn:yhda]
CONTIG454	32230311_f3_56	2403	8065	1170	390	1827	1.5(10)-188	Escherichia coli	b3251	[pn:rod shape-determining protein mreB] [gn:mreb]
CONTIG454	35282183_f3_70	2404	8066	1674	558	1376	9.1(10)-141	Escherichia coli	b2661	[pn:succinate-semialdehyde dehydrogenase] [gn:gabd]
CONTIG454	22917932_f3_71	2405	8067	315	105	302	5.9(10)-27	Escherichia coli	b3239	[pn:hypothetical protein] [gn:yhco]
CONTIG454	23650765_c1_77	2406	8068	1125	375	1305	3.1(10)-133	Escherichia coli	b3235	[pn:protease precursor] [gn:degS]
CONTIG454	16265886_c1_110	2407	8069	1062	354	1419	2.6(10)-145	Escherichia coli	b1971	[pn:hypothetical protein]
CONTIG454	14455203_c1_111	2408	8070	243	81	309	1.1(10)-27	Escherichia coli	b1972	[pn:hypothetical protein]
CONTIG454	6070136_c2_112	2409	8071	1425	475	1622	7.9(10)-167	Escherichia coli	b3234	[pn:protease precursor] [gn:degQ]
CONTIG454	34244001_c2_116	2410	8072	219	73	124	4.2(10)-8	Escherichia coli	b3237	[pn:arginine repressor] [gn:argR]
CONTIG454	4144568_c2_146	2411	8073	984	328	1280	1.3(10)-130	Escherichia coli	b3253	[pn:hypothetical 34.7 kd protein in mreB-accb intergenic region] [gn:yhdh]
CONTIG454	12585313_c3_149	2412	8074	474	158	540	3.6(10)-52	Escherichia coli	b3233	[pn:hypothetical 15.2 kd protein in rplM-hhoA intergenic region] [gn:yhcb]
CONTIG454	24220842_c3_150	2413	8075	438	146	632	6.4(10)-62	Escherichia coli	b3237	[pn:arginine repressor] [gn:argR]
CONTIG454	24644052_c3_151	2414	8076	270	90	176	1.3(10)-13	Escherichia coli	b3238	[pn:hypothetical protein] [gn:yhcn]
CONTIG454	13683312_c3_155	2415	8077	537	179	164	2.5(10)-12	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:car
CONTIG454	32244052_c3_161	2416	8078	930	310	1440	1.5(10)-147	Escherichia coli	b3243	le:59 re:580 di:direct nt:orf2
CONTIG455	23642302_f1_27	2417	8079	1731	577	2126	3.1(10)-220	Escherichia coli	b0445	[pn:hypothetical protein] [gn:yhcs]
										[pn:hypothetical 65.0 kd protein in hupB-cof intergenic region]

CONTIG455	32228380_fl_28	2418	8080	699	233	1133	5.2(10)-115	Escherichia coli	b0444	[pn:hypothetical protein] [gn:ybax]
CONTIG455	7686_f2_62	2419	8081	1107	369	976	2.2(10)-98	Mycobacterium tuberculosis	AL022121	[de:mycobacterium tuberculosis sequence v025.] [pn:putative lyase] [gn:mtv025.032] [nt:mtv025.032,
CONTIG455	32458412_f3_76	2420	8082	1566	522	1507	1.2(10)-154	Escherichia coli	b0457	[pn:hypothetical protein] [gn:ylab]
CONTIG455	24254782_f3_85	2421	8083	1539	513	284	4.5(10)-25	Escherichia coli	b1439	[pn:hypothetical protein]
CONTIG455	36066375_f3_87	2422	8084	330	110	484	3.1(10)-46	Escherichia coli	b0454	[pn:hypothetical protein] [gn:ybaz]
CONTIG455	13751263_f3_88	2423	8085	942	314	1283	6.5(10)-131	Escherichia coli	b0452	[pn:acyl-coa thioesterase ii] [gn:tesb]
CONTIG455	31439416_c1_105	2424	8086	504	168	550	3.1(10)-53	Escherichia coli	b0443	[pn:hypothetical protein] [gn:ybau]
CONTIG455	24500877_c1_106	2425	8087	363	121	128	1.6(10)-8	Haemophilus influenzae	HI1420	[pn:hypothetical protein]
CONTIG455	3913580_c1_115	2426	8088	621	207	794	4.2(10)-79	Escherichia coli	b0447	[pn:hypothetical protein]
CONTIG455	4875126_c1_116	2427	8089	1785	595	2281	1.2(10)-236	Escherichia coli	b0448	[pn:mdla]
CONTIG455	35582912_c1_126	2428	8090	582	194	627	2.2(10)-61	Escherichia coli	b0453	[pn:hypothetical protein] [gn:ybay]
CONTIG455	29877090_c1_134	2429	8091	1122	374	279	1.1(10)-29	Haemophilus influenzae	HI1635	[pn:purine nucleotide synthesis repressor protein] [gn:purr]
CONTIG455	4140_c2_140	2430	8092	2034	678	2534	1.8(10)-263	Escherichia coli	b0441	[pn:hypothetical protein] [gn:ybau]
CONTIG455	5120650_c2_148	2431	8093	2046	682	2330	7.4(10)-242	Escherichia coli	b0449	[pn:mdlb]
CONTIG455	6536566_c2_150	2432	8094	1338	446	1687	1.0(10)-173	Escherichia coli	b0451	[pn:hypothetical protein in mdl-tesb intergenic region] [gn:amb]
CONTIG455	4509562_c2_157	2433	8095	678	226	309	1.1(10)-27	Haemophilus influenzae	HI0522	[pn:sp]
CONTIG455	24078333_c2_160	2434	8096	3111	1037	3614	0	Escherichia coli	b0344	[pn:beta-galactosidase] [gn:lacz]
CONTIG455	12698806_c3_164	2435	8097	522	174	827	1.3(10)-82	Escherichia coli	b0439	[pn:lon protease] [gn:lon]
CONTIG455	33860768_c3_165	2436	8098	360	120	292	6.7(10)-26	Escherichia coli	b0440	[pn:dna-binding protein hu-beta] [gn:hupb]

CONTIG455	24344391_c3_166	2437	8099	501	167	339	7.0(10)-31	Escherichia coli	b0442	[pn:hypothetical protein] [gn:ybav]
CONTIG455	34257212_c3_168	2438	8100	381	127	130	1.0(10)-8	Haemophilus influenzae	H11419	[pn:hypothetical protein]
CONTIG455	4470206_c3_170	2439	8101	843	281	1105	4.7(10)-112	Escherichia coli	b0446	[pn:cof protein] [gn:cof]
CONTIG455	886451_c3_177	2440	8102	375	125	540	3.6(10)-52	Escherichia coli	b0450	[pn:nitrogen regulatory protein p-ii] [gn:glnk]
CONTIG456	17069787_f1_4	2441	8103	1119	373	1812	5.7(10)-187	Escherichia coli	b2806	[pn:hypothetical 41.9 kd protein in fucr-gcva intergenic region] [gn:ygde]
CONTIG456	14339808_f1_12	2442	8104	462	154	381	2.5(10)-35	Escherichia coli	b2792	[pn:hypothetical protein]
CONTIG456	24726386_f1_18	2443	8105	1383	461	2148	1.3(10)-222	Escherichia coli	b2788	[pn:hypothetical protein] [gn:ygy]
CONTIG456	14704837_f1_19	2444	8106	1350	450	2115	4.5(10)-219	Escherichia coli	b2787	[pn:hypothetical protein] [gn:ygcx]
CONTIG456	1440751_f1_29	2445	8107	1653	551	2559	4.0(10)-266	Escherichia coli	b2780	[pn:ctp synthase] [gn:pyrg]
CONTIG456	960000_f1_31	2446	8108	1350	450	749	2.5(10)-74	Escherichia coli	b1621	[pn:pts system, maltose and glucose-specific ii abc component] [gn:malx]
CONTIG456	24480282_f2_49	2447	8109	1389	463	2172	4.0(10)-225	Escherichia coli	b2789	[pn:hypothetical protein]
CONTIG456	35347826_f2_61	2448	8110	1566	522	1626	3.0(10)-167	Escherichia coli	b2785	[pn:hypothetical rna methyltransferase in rela-bara intergenic region] [gn:ygca]
CONTIG456	5886253_f2_62	2449	8111	2280	760	3496	0	Escherichia coli	b2784	[pn:gtp pyrophosphokinase] [gn:rela]
CONTIG456	16208537_f2_66	2450	8112	1371	457	1985	2.7(10)-205	Escherichia coli	b2779	[pn:lase] [gn:eno]
CONTIG456	22147011_f3_71	2451	8113	936	312	1509	7.4(10)-155	Escherichia coli	b2808	[pn:regulatory protein for glycine cleavage pathway] [gn:gcva]
CONTIG456	15868766_f3_72	2452	8114	435	145	613	6.5(10)-60	Escherichia coli	b2807	[pn:hypothetical 14.3 kd protein in fucr-gcva intergenic region] [gn:ygdd]
CONTIG456	33750181_f3_83	2453	8115	606	202	679	6.7(10)-67	Escherichia coli	b2793	[pn:syd] [gn:syd]
CONTIG456	14880066_f3_85	2454	8116	792	264	1145	2.7(10)-116	Escherichia coli	b2791	[pn:hypothetical protein] [gn:yqcb]

CONTIG456	22870800_f3_86	2455	8117	465	155	622	7.2(10)-61	Escherichia coli	b2790	[pn:hypothetical protein]
CONTIG456	3933193_f3_89	2456	8118	1155	385	1206	9.5(10)-123	Escherichia coli	b3124	[pn:hypothetical 42.1 kd protein in rnpb-soha intergenic region] [gn:yhad]
CONTIG456	16145293_f3_99	2457	8119	834	278	1014	2.1(10)-102	Escherichia coli	b2781	[pn:mazg protein] [gn:mazg]
CONTIG456	563568_c1_137	2458	8120	1374	458	2203	2.1(10)-228	Escherichia coli	b2795	[pn:hypothetical protein in sdac 5'''region] [gn:ygdh]
CONTIG456	4298457_c1_138	2459	8121	1371	457	1818	1.3(10)-187	Escherichia coli	b2796	[pn:putative serine transporter] [gn:sdac]
CONTIG456	34664182_c1_140	2460	8122	774	258	1130	1.1(10)-114	Escherichia coli	b2798	[pn:potential 5'''-3''' nuclease] [gn:exo]
CONTIG456	4563193_c2_147	2461	8123	834	278	323	3.5(10)-29	Haemophilus influenzae	H10143	[pn:gb]
CONTIG456	19619037_c2_175	2462	8124	846	282	1306	2.3(10)-133	Escherichia coli	b2794	[pn:hypothetical protein] [gn:yqcd]
CONTIG456	13707307_c3_188	2463	8125	1299	433	200	3.0(10)-13	Schizosaccharomyces pombe	L37084	or:schizosaccharomyces pombe pn:phosphopyruvate hydratase ec:4.2.1.11 le:2 re:1342 di:complement sr:schizosaccharomyces pombe cdna to mna
CONTIG456	25831336_c3_191	2464	8126	1398	466	175	5.2(10)-12	Escherichia coli	A30374	hypothetical 77k protein (spot 3' region) - escherichia coli
CONTIG456	4379716_c3_195	2465	8127	2769	923	3912	0	Escherichia coli	b2786	[pn:sensor protein bara] [gn:bara]
CONTIG456	32314042_c3_212	2466	8128	1383	461	2082	1.3(10)-215	Escherichia coli	b2797	[pn:l-serine dehydratase 2] [gn:sdab]
CONTIG457	47891_f1_26	2467	8129	501	167	597	3.2(10)-58	Escherichia coli	b3355	[pn:phosphoribulokinase] [gn:prkb]
CONTIG457	35792657_f1_36	2468	8130	822	274	262	1.0(10)-22	Escherichia coli	U82664	or:escherichia coli le:133380 re:134066 di:direct nt:hypothetical protein
CONTIG457	26759702_f2_37	2469	8131	222	74	321	5.7(10)-29	Escherichia coli	b3348	[pn:slyx protein] [gn:slyx]
CONTIG457	24412918_f2_38	2470	8132	654	218	539	4.5(10)-52	Escherichia coli	B49988	hypothetical protein 159 - escherichia coli

CONTIG457	3947187_f2_64	2471	8133	222	74	318	1.2(10)-28	Escherichia coli	b3354	[pn:hypothetical 8.5 kd protein in kifb-prkb intergenic region]
CONTIG457	568755_f2_65	2472	8134	531	177	653	3.7(10)-64	Escherichia coli	b3355	[pn:phosphoribulokinase] [gn:prkb]
CONTIG457	7067711_f2_68	2473	8135	639	213	1070	2.5(10)-108	Escherichia coli	b3357	[pn:cyclic amp receptor protein] [gn:crp]
CONTIG457	24308275_f2_69	2474	8136	2145	715	2846	1.6(10)-296	Escherichia coli	b3358	[pn:hypothetical 79.5 kd protein in crp-argd intergenic region] [gn:yhfk]
CONTIG457	4484686_f3_78	2475	8137	1938	646	2854	2.2(10)-297	Escherichia coli	b3352	[pn:hypothetical abc transporter in kifb-prkb intergenic region]
CONTIG457	34660411_f3_79	2476	8138	948	316	1171	4.9(10)-119	Klebsiella pneumoniae	U95087	[PN:MdcR] [GN:mdcR] [FN:regulates expression of malonate decarboxylase] [DE:Klebsiella pneumoniae malonate decarboxylase gene cluster (mdcA,mdcB,mdcC,mdcD,mdcE,mdcF,mdcG,mdcH,mdcR) genes, completecds.] [NT:transcriptional regul
CONTIG457	34103250_f3_91	2477	8139	1050	350	1317	1.6(10)-134	Escherichia coli	b3353	[pn:hypothetical 38.5 kd protein in kifb-prkb intergenic region]
CONTIG457	132825_f3_92	2478	8140	318	106	137	9.3(10)-9	Escherichia coli	U18997	or:escherichia coli le:265033 re:266106 di:direct nt:orf_o357; codon usage statistics poor where o72
CONTIG457	21678790_c1_100	2479	8141	573	191	737	4.7(10)-73	Escherichia coli	b3363	[pn:peptidyl-prolyl cis-trans isomerase a] [gn:ppia]
CONTIG457	21925080_c1_101	2480	8142	690	230	542	2.2(10)-52	Escherichia coli	b3361	[pn:cell filamentation protein] [gn:fic]
CONTIG457	16933387_c1_102	2481	8143	594	198	804	3.7(10)-80	Escherichia coli	b3360	[pn:para-aminobenzoate synthetase] [gn:paba]
CONTIG457	21485675_c1_103	2482	8144	1266	422	1917	4.2(10)-198	Escherichia coli	b3359	[pn:acetylornithine delta-aminotransferase] [gn:argd]
CONTIG457	22689078_c1_106	2483	8145	411	137	641	7.0(10)-63	Escherichia coli	b3356	[pn:hypothetical protein] [gn:yhfa]

CONTIG457	26019656_c1_111	2484	8146	885	295	985	2.5(10)-99	Klebsiella pneumoniae	U95087	[PN:MdcB] [GN:mdcB] [FN:involved in biosynthesis of the prosthetic] [DE:Klebsiella pneumoniae malonate decarboxylase gene cluster (mdcA,mdcB,mdcC,mdcD,mdcE,mdcF,mdcG,mdcH,mdcR) genes, completecds.] [NT:similar to CitG proteins]
CONTIG457	3954838_c1_112	2485	8147	309	103	366	9.8(10)-34	Klebsiella pneumoniae	U95087	[PN:MdcC] [GN:mdcC] [DE:Klebsiella pneumoniae malonate decarboxylase gene cluster (mdcA,mdcB,mdcC,mdcD,mdcE,mdcF,mdcG,mdcH,mdcR) genes, completecds.] [NT:acyl carrier protein; delta subunit of malonate] [LE:2803] [RE:3102] [DI:
CONTIG457	29297917_c1_113	2486	8148	828	276	1026	1.1(10)-103	Klebsiella pneumoniae	U95087	[PN:MdcE] [GN:mdcE] [DE:Klebsiella pneumoniae malonate decarboxylase gene cluster (mdcA,mdcB,mdcC,mdcD,mdcE,mdcF,mdcG,mdcH,mdcR) genes, completecds.] [NT:decarboxylase subunit; gamma subunit of malonate] [LE:3928] [RE:4728] [DI
CONTIG457	19650250_c1_114	2487	8149	963	321	1138	1.5(10)-115	Klebsiella pneumoniae	U95087	[PN:MdcF] [GN:mdcF] [FN:putative malonate transporter] [DE:Klebsiella pneumoniae malonate decarboxylase gene cluster (mdcA,mdcB,mdcC,mdcD,mdcE,mdcF,mdcG,mdcH,mdcR) genes, completecds.] [NT:encodes ten hydrophobic domains] [LE:4
CONTIG457	35832883_c1_122	2488	8150	1950	650	2271	1.3(10)-235	Escherichia coli	b3350	[pn:glutathione-regulated potassium-efflux system protein] [gn:kfb]

CONTIG457	1449027_c2_138	2489	8151	1671	557	2587	4.2(10)-269	Klebsiella pneumoniae	U95087	[PN:MdcA] [GN:mdcA] [DE:Klebsiella pneumoniae malonate decarboxylase gene cluster (mdcA,mdcB,mdcC,mdcD,mdcE,mdcF,mdcG,mdcH,mdcR) genes, completecds.] [NT:acyl carrier protein transferase; alpha subunit of] [LE:288] [RE:1943] [D
CONTIG457	25650277_c2_140	2490	8152	837	279	1180	5.4(10)-120	Klebsiella pneumoniae	U95087	[PN:MdcD] [GN:mdcD] [DE:Klebsiella pneumoniae malonate decarboxylase gene cluster (mdcA,mdcB,mdcC,mdcD,mdcE,mdcF,mdcG,mdcH,mdcR) genes, completecds.] [NT:decarboxylase subunit; beta subunit of malonate] [LE:3095] [RE:3928] [DI:
CONTIG457	10444826_c2_142	2491	8153	1059	353	1145	2.7(10)-116	Klebsiella pneumoniae	U56096	or:klebsiella pneumoniae pn:mdcg gn:mdcg le:6354 re:7337 di:direct nt:similar to malonyl coa-acyl carrier protein
CONTIG457	2625166_c2_147	2492	8154	792	264	330	6.4(10)-30	Pseudomonas aeruginosa	JQ0133	hypothetical 26.4k protein - pseudomonas aeruginosa
CONTIG457	14960906_c2_148	2493	8155	597	199	882	2.0(10)-88	Escherichia coli	b3351	[pn:hypothetical nadph oxidoreductase in kifb-prkb intergenic region] [gn:yher]
CONTIG457	14461081_c3_160	2494	8156	564	188	537	7.4(10)-52	Erwinia chrysanthemi	X89443	[GN:ORF3] [DE:E.chrysanthemi DNA for crp gene.] [LE:1200] [RE:1874] [DI:complement]
CONTIG457	3928590_c3_161	2495	8157	198	66	145	2.6(10)-10	Erwinia chrysanthemi	X89443	[GN:ORF3] [DE:E.chrysanthemi DNA for crp gene.] [LE:1200] [RE:1874] [DI:complement]
CONTIG457	15752042_c3_173	2496	8158	696	232	643	4.2(10)-63	Klebsiella pneumoniae	U95087	[PN:MdcG] [GN:mdcG] [FN:involved in formation of the holo-acyl carrier] [DE:Klebsiella pneumoniae malonate decarboxylase gene cluster (mdcA,mdcB,mdcC,mdcD,mdcE,mdcF,mdcG,mdcH,mdcR) genes, completecds.] [LE:5828] [RE:6445] [DI:d



CONTIG457	33984378_c3_186	2497	8159	672	224	742	1.3(10)-73	Escherichia coli	b3349	[pn:probable fkbp-type peptidyl-prolyl cis-trans isomerase] [gn:slvd]
CONTIG458	36150466_f1_9	2498	8160	423	141	320	7.2(10)-29	Escherichia coli	b1926	[pn:flagellar protein flit] [gn:flit]
CONTIG458	12601516_f1_16	2499	8161	1713	571	1997	1.3(10)-206	Escherichia coli	b1938	[pn:flagellar basal-body m-ring protein] [gn:flif]
CONTIG458	2847202_f1_19	2500	8162	1215	405	430	1.0(10)-49	Escherichia coli	b1943	[pn:hook-length control protein] [gn:flif]
CONTIG458	13933302_f1_22	2501	8163	441	147	267	3.0(10)-23	Escherichia coli	b1947	[pn:flagellar protein flio] [gn:flio]
CONTIG458	12362590_f1_24	2502	8164	387	129	292	6.7(10)-26	Escherichia coli	b1949	[pn:flagellar biosynthetic protein flit] [gn:flit]
CONTIG458	3166591_f1_25	2503	8165	792	264	919	2.5(10)-92	Escherichia coli	b1950	[pn:flagellar biosynthetic protein flir] [gn:flir]
CONTIG458	4423218_f1_26	2504	8166	633	211	916	5.0(10)-92	Escherichia coli	b1951	[pn:colanic acid capsular biosynthesis activation protein a] [gn:rcsa]
CONTIG458	3962502_f1_29	2505	8167	927	309	821	6.0(10)-82	Escherichia coli	b1955	[pn:hypothetical protein]
CONTIG458	4100318_f2_42	2506	8168	1422	474	1613	7.0(10)-166	Escherichia coli	b1924	[pn:flagellar hook associated protein 2] [gn:ftid]
CONTIG458	33631457_f2_43	2507	8169	438	146	481	6.4(10)-46	Escherichia coli	b1925	[pn:flagellar protein flis] [gn:flis]
CONTIG458	911533_f2_51	2508	8170	1077	359	1487	1.6(10)-152	Escherichia coli	b1939	[pn:flagellar motor switch protein flig] [gn:flig]
CONTIG458	35242718_f2_53	2509	8171	1443	481	2199	5.5(10)-228	Escherichia coli	b1941	[pn:flagellum-specific atp synthase] [gn:flit]
CONTIG458	34620911_f2_54	2510	8172	465	155	635	3.1(10)-62	Escherichia coli	b1942	[pn:flagellar flit protein] [gn:flit]
CONTIG458	35727283_f2_56	2511	8173	528	176	552	1.8(10)-53	Escherichia coli	b1944	[pn:flit protein] [gn:flit]
CONTIG458	22007193_f2_57	2512	8174	426	142	633	5.0(10)-62	Escherichia coli	b1946	[pn:flagellar motor switch protein flin] [gn:flin]
CONTIG458	10015706_f3_74	2513	8175	1560	520	2350	5.5(10)-244	Escherichia coli	b1927	[pn:cytoplasmic alpha-amylase] [gn:amya]
CONTIG458	26694825_f3_78	2514	8176	729	243	608	7.0(10)-71	Escherichia coli	b1940	[pn:flagellar assembly protein flit] [gn:flit]
CONTIG458	2535457_f3_82	2515	8177	1068	356	1634	4.2(10)-168	Escherichia coli	b1945	[pn:cg site no. 774] [gn:flim]

CONTIG458	12397666_f3_84	2516	8178	804	268	730	2.6(10)-72	Escherichia coli	b1948	[pn:flagellar biosynthetic protein flip] [gn:flip]
CONTIG458	32306326_f3_87	2517	8179	258	86	272	9.0(10)-24	Escherichia coli	b1953	[pn:hypothetical protein]
CONTIG458	29939465_c1_93	2518	8180	348	116	306	2.2(10)-27	Escherichia coli	b1952	[pn:dsrb protein] [gn:dsrb]
CONTIG458	4332291_c1_105	2519	8181	309	103	131	7.7(10)-9	Bacillus subtilis	S14505	hypothetical protein 12 (flaa operon) - bacillus subtilis
CONTIG458	21910312_c1_112	2520	8182	336	112	406	5.7(10)-38	Escherichia coli	b1937	[pn:flagellar hook-basal body complex protein flie] [gn:flie]
CONTIG458	4006555_c1_113	2521	8183	489	163	486	1.8(10)-46	Escherichia coli	b1928	[pn:hypothetical 15.0 kd protein in amya-flie intergenic region] [gn:yedd]
CONTIG458	9800156_c1_123	2522	8184	810	270	1133	5.2(10)-115	Escherichia coli	b1922	[pn:ma polymerase sigma transcription factor for flagellar operon] [gn:flia]
CONTIG458	6928936_c1_124	2523	8185	567	189	718	4.9(10)-71	Escherichia coli	b1921	[pn:fliz protein] [gn:fliz]
CONTIG458	20837_c1_125	2524	8186	1059	353	1369	5.0(10)-140	Escherichia coli	b1919	[pn:hypothetical protein]
CONTIG458	33491312_c1_126	2525	8187	690	230	966	2.6(10)-97	Escherichia coli	b1918	[pn:yecg]
CONTIG458	33991554_c2_129	2526	8188	509	169	528	6.7(10)-51	Escherichia coli	b1956	[pn:hypothetical protein]
CONTIG458	24507182_c3_179	2527	8189	1728	576	2023	2.5(10)-209	Escherichia coli	U47614	or:escherichia coli pn:flagellin gn:flc le:1 re:1758 di:direct
CONTIG458	32692516_c3_181	2528	8190	855	285	1110	1.3(10)-112	Escherichia coli	b1920	[pn:fly protein precursor] [gn:fly]
CONTIG458	34414030_c3_184	2529	8191	759	253	1160	7.0(10)-118	Escherichia coli	b1917	[pn:hypothetical protein] [gn:yecg]
CONTIG459	3239041_f1_4	2530	8192	444	148	141	6.7(10)-10	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:59 re:580 di:direct nt:orf2
CONTIG459	26735180_f1_7	2531	8193	933	311	1015	1.7(10)-102	Haemophilus influenzae	HI0362	[pn:adhesin b precursor] [gn:fima]
CONTIG459	24422707_f1_10	2532	8194	1167	389	635	3.1(10)-62	Haemophilus influenzae	HI0359	[pn:gb]
CONTIG459	13163556_f1_12	2533	8195	1380	460	1480	8.8(10)-152	Escherichia coli	b0585	[pn:enterochelin esterase] [gn:fes]

CONTIG459	781658_f1_l3	2534	8196	3885	1295	4972	0	Escherichia coli	b0586	[pn:enterobactin synthetase component f] [gn:entf]
CONTIG459	21505208_f2_35	2535	8197	954	318	669	7.5(10)-66	Haemophilus influenzae	HI0360	[pn:hydrophobic membrane protein]
CONTIG459	33875656_f2_42	2536	8198	249	83	262	1.0(10)-22	Escherichia coli	B31958	hypothetical 8k protein (fes 3' region) - escherichia coli
CONTIG459	1425657_f3_61	2537	8199	1521	507	90	0.12	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:<1 re:588 di:direct
CONTIG459	5316943_f3_62	2538	8200	2043	681	3181	0	Escherichia coli	b0314	[pn:high-affinity choline transport protein] [gn:beti]
CONTIG459	20318790_f3_63	2539	8201	348	116	114	5.0(10)-7	Escherichia coli	b3555	[pn:hypothetical 11.0 kd protein in bise-cspa intergenic region]
CONTIG459	12370787_f3_66	2540	8202	948	316	819	9.6(10)-82	Haemophilus influenzae	HI0361	[pn:iii dicitrate transport atp-binding protein fece] [gn:fece]
CONTIG459	6539591_f3_81	2541	8203	354	118	91	0.00129	Mus musculus	P21952	octamer-binding transcription factor 6 (oct-6) (pou-domain transcription factor scip).
CONTIG459	679507_f3_87	2542	8204	1263	421	1338	9.8(10)-137	Escherichia coli	b0591	[pn:hypothetical membrane protein p43] [gn:ybda]
CONTIG459	10554188_c1_91	2543	8205	996	332	1279	1.7(10)-130	Escherichia coli	b0592	[pn:ferriterobactin-binding periplasmic protein precursor] [gn:fepb]
CONTIG459	31848911_c1_95	2544	8206	1320	440	1102	1.0(10)-111	Escherichia coli	b0589	[pn:ferric enterobactin transport protein fepg] [gn:fepg]
CONTIG459	19538413_c1_115	2545	8207	663	221	911	1.7(10)-91	Escherichia coli	b0313	[pn:regulatory protein beti] [gn:beti]
CONTIG459	5267181_c2_127	2546	8208	1059	353	1010	5.5(10)-102	Escherichia coli	b0590	[pn:ferric enterobactin transport protein fepd] [gn:fepd]
CONTIG459	29979075_c2_146	2547	8209	1593	531	2240	2.6(10)-232	Escherichia coli	b0312	[pn:betaine aldehyde dehydrogenase] [gn:betb]
CONTIG459	16970218_c3_156	2548	8210	720	240	275	4.2(10)-24	Escherichia coli	P21500	very hypothetical 18.0 kd protein in fepb 3'region.
CONTIG459	16270216_c3_161	2549	8211	978	326	1127	2.2(10)-114	Escherichia coli	b0588	[pn:ferric enterobactin transport atp-binding protein fepc] [gn:fepc]

CONTIG459	16219582_c3_162	2550	8212	327	109	92	0.0011	Nicotiana alata	U88587	[de:nicotiana alata 120 kda style glycoprotein (naprp5) mrna, completecds.] [pn:120 kda style glycoprotein] [gn:naprp5] [nt:style-specific protein possessing features of]
CONTIG459	24254052_c3_172	2551	8213	2265	755	2862	3.1(10)-298	Escherichia coli	b0584	[pn:ferrienterobactin receptor precursor] [gn:fepa]
CONTIG459	12134390_c3_173	2552	8214	762	254	465	3.2(10)-44	Escherichia coli	b0583	[pn:enterobactin synthetase component d] [gn:entd]
CONTIG459	33678892_c3_184	2553	8215	1722	574	2774	6.5(10)-289	Escherichia coli	b0311	[pn:choline dehydrogenase] [gn:beta]
CONTIG46	25397792_f2_2	2554	8216	435	145	462	6.5(10)-44	Escherichia coli	b0814	[pn:outer membrane protein x precursor] [gn:ompx]
CONTIG460	4392308_f1_11	2555	8217	531	177	624	4.5(10)-61	Escherichia coli	b3995	[pn:hypothetical protein] [gn:yjae]
CONTIG460	13759633_f1_12	2556	8218	1899	633	3132	0	Escherichia coli	b3994	[pn:thic protein] [gn:thic]
CONTIG460	32661541_f1_14	2557	8219	945	315	799	1.3(10)-79	Escherichia coli	b3992	[pn:thif]
CONTIG460	35267665_f1_19	2558	8220	357	119	189	5.5(10)-15	Bacillus subtilis	licA	[pn:phosphotransferase system] [gn:celc]
CONTIG460	16510407_f2_52	2559	8221	237	79	173	2.7(10)-13	Escherichia coli	S77700	[PN:thiG1 protein]
CONTIG460	13073963_f2_53	2560	8222	1800	600	1706	9.9(10)-176	Escherichia coli	b3990	[pn:thih protein] [gn:thih]
CONTIG460	11019027_f2_54	2561	8223	1656	552	245	3.8(10)-20	Escherichia coli	b1535	[pn:ydeh]
CONTIG460	24256925_f2_55	2562	8224	342	114	219	3.7(10)-18	Bacillus subtilis	ydhM	[pn:hypothetical protein]
CONTIG460	10196957_f3_65	2563	8225	1734	578	2535	1.3(10)-263	Escherichia coli	b4006	[pn:phosphoribosylaminoimidazolec arboxamide formyltransferase and imp cyclohydrolase]
CONTIG460	22869525_f3_66	2564	8226	1308	436	2030	4.5(10)-210	Escherichia coli	b4005	[pn:phosphoribosylglycineamide synthetase] [gn:purd]
CONTIG460	35339517_f3_74	2565	8227	717	239	899	3.2(10)-90	Escherichia coli	b3993	[pn:thie protein] [gn:thie]
CONTIG460	32245792_f3_77	2566	8228	885	295	1226	7.2(10)-125	Escherichia coli	b3991	[pn:thig protein] [gn:thig]

CONTIG460	24647902_c1_92	2567	8229	4167	1389	6740	0	Escherichia coli	b3987	[pn:dna-directed rna polymerase, beta-subunit] [gn:rpob]
CONTIG460	31291250_c1_112	2568	8230	612	204	886	7.7(10)-89	Escherichia coli	b3999	[pn:hypothetical 22.6 kd protein in heme-hupa intergenic region] [gn:yjag]
CONTIG460	24025251_c1_113	2569	8231	276	92	424	7.0(10)-40	Escherichia coli	b4000	[pn:histonelike dna-binding protein hu-alpha] [gn:hupa]
CONTIG460	16525791_c2_126	2570	8232	4266	1422	6482	0	Escherichia coli	b3988	[pn:dna-directed rna polymerase, beta'''-subunit] [gn:rpoc]
CONTIG460	16219716_c2_140	2571	8233	1092	364	1701	3.2(10)-175	Escherichia coli	b3997	[pn:uroporphyrinogen decarboxylase] [gn:heme]
CONTIG460	3962943_c2_141	2572	8234	681	227	1077	4.4(10)-109	Escherichia coli	b3998	[pn:hypothetical 24.9 kd protein in heme-hupa intergenic region] [gn:yjaf]
CONTIG460	13792203_c3_147	2573	8235	258	86	382	2.0(10)-35	Escherichia coli	b3985	[pn:50s ribosomal subunit protein 110] [gn:rplj]
CONTIG460	14094452_c3_148	2574	8236	366	122	372	2.2(10)-34	Escherichia coli	b3986	[pn:50s ribosomal subunit protein 17/112] [gn:rplj]
CONTIG460	33492292_c3_162	2575	8237	795	265	1160	7.0(10)-118	Escherichia coli	b3996	[pn:hypothetical 29.8 kd protein in thic-heme intergenic region] [gn:yjad]
CONTIG460	6508428_c3_168	2576	8238	708	236	906	5.9(10)-91	Escherichia coli	b4001	[pn:hypothetical 26.3 kd protein in hupa-hydh intergenic region] [gn:yjah]
CONTIG461	36428933_f1_16	2577	8239	897	299	944	5.5(10)-95	Escherichia coli	b0826	[pn:molybdopterin biosynthesis moeb protein] [gn:moeb]
CONTIG461	24120405_f1_19	2578	8240	915	305	1322	4.7(10)-135	Escherichia coli	b0824	[pn:hypothetical protein]
CONTIG461	26041332_f1_31	2579	8241	2403	801	560	2.7(10)-52	Saccharomyces cerevisiae	YBR229C	[pn:glucosidase ii, catalytic subunit] [gn:rot2]
CONTIG461	24103381_f2_39	2580	8242	1329	443	2083	1.1(10)-215	Escherichia coli	b0835	[pn:hypothetical protein] [gn:ylig]
CONTIG461	32229531_f2_40	2581	8243	555	185	101	0.00072	Streptococcus pneumoniae	L29323	or:streptococcus pneumoniae pn:methyl transferase le:508 re:1125 di:complement sr:transposon tn5252 dna; and streptococcus pneumoniae (strain sp1000 nt:member of the mtr gene cluster; putative

CONTIG461	19633331_f2_50	2582	8244	1173	391	1531	3.5(10)-157	Escherichia coli	b0823	[pn:hypothetical protein]
CONTIG461	11845627_f2_52	2583	8245	879	293	1168	1.0(10)-118	Escherichia coli	b0822	[pn:hypothetical protein]
CONTIG461	2540786_f2_57	2584	8246	1347	449	923	9.3(10)-93	Escherichia coli	b3093	[pn:hexuronate transporter] [gn:exut]
CONTIG461	24620452_f2_61	2585	8247	1236	412	161	7.7(10)-10	Pseudomonas aeruginosa	JQ0133	hypothetical 26.4k protein - pseudomonas aeruginosa
CONTIG461	287575_f3_64	2586	8248	690	230	837	1.2(10)-83	Escherichia coli	b0838	[pn:hypothetical protein]
CONTIG461	9770206_f3_77	2587	8249	1287	429	1701	3.2(10)-175	Escherichia coli	b0827	[pn:molybdopterin biosynthesis moea protein] [gn:moea]
CONTIG461	16101557_f3_81	2588	8250	1455	485	2307	2.0(10)-239	Escherichia coli	b0823	[pn:hypothetical protein]
CONTIG461	2734432_f3_82	2589	8251	1293	431	1983	4.4(10)-205	Escherichia coli	b0821	[pn:hypothetical protein]
CONTIG461	22949052_c1_113	2590	8252	1887	629	2443	7.9(10)-254	Escherichia coli	b0829	[pn:hypothetical protein]
CONTIG461	16832768_c1_118	2591	8253	414	138	358	6.9(10)-33	Escherichia coli	b0836	[pn:hypothetical protein]
CONTIG461	23573328_c2_137	2592	8254	690	230	955	3.7(10)-96	Escherichia coli	b0825	[pn:hypothetical protein]
CONTIG461	34414165_c2_142	2593	8255	945	315	1234	1.0(10)-125	Escherichia coli	b0828	[pn:hypothetical protein in moea-grxa intergenic region] [gn:ybik]
CONTIG461	23446041_c2_147	2594	8256	942	314	1292	7.2(10)-132	Escherichia coli	b0831	[pn:hypothetical protein]
CONTIG461	22146880_c2_151	2595	8257	1152	384	1539	4.9(10)-158	Escherichia coli	b0837	[pn:hypothetical protein]
CONTIG461	13704165_c2_153	2596	8258	345	115	248	5.9(10)-21	Escherichia coli	b0839	[pn:penicillin-binding protein 6 precursor] [gn:dacc]
CONTIG461	19583290_c3_154	2597	8259	1632	544	2615	4.7(10)-272	Escherichia coli	b0820	[pn:hypothetical protein]
CONTIG461	20179036_c3_164	2598	8260	1125	375	244	8.5(10)-20	Escherichia coli	b3934	[pn:transcriptional repressor cytr] [gn:cytr]
CONTIG461	31730393_c3_180	2599	8261	1623	541	2250	2.2(10)-233	Escherichia coli	b0830	[pn:hypothetical protein]
CONTIG461	30511340_c3_182	2600	8262	963	321	1247	4.2(10)-127	Escherichia coli	b0832	[pn:hypothetical protein]

CONTIG462	15132125_f1_2	2601	8263	828	276	907	4.5(10)-91	Escherichia coli	b1859	[pn:hypothetical 27.8 kd protein in msbb-ruvb intergenic region] [gn:yebi]
CONTIG462	24330036_f1_10	2602	8264	300	100	325	2.2(10)-29	Salmonella typhimurium	Q56031	virulence protein msga.
CONTIG462	14879752_f2_33	2603	8265	762	254	1110	1.3(10)-112	Escherichia coli	b1858	[pn:hypothetical protein] [gn:yebm]
CONTIG462	21677187_f3_66	2604	8266	594	198	794	4.2(10)-79	Escherichia coli	b1867	[pn:hypothetical 21.8 kd protein in asps 5'''region] [gn:yecd]
CONTIG462	35604677_f3_68	2605	8267	822	274	155	7.7(10)-10	Bacteriophage M1	P08231	tail fiber protein gp37 (fragment).
CONTIG462	36125431_f3_71	2606	8268	216	72	93	8.3(10)-5	Bacteriophage phi-80	P17651	adsorption-inhibiting cor protein.
CONTIG462	22163441_c1_92	2607	8269	396	132	156	1.8(10)-11	Bacteriophage lambda	P03737	minor tail protein m.
CONTIG462	6062568_c1_94	2608	8270	597	199	198	6.2(10)-16	Yersinia pestis	AF053947	[de:yersinia pestis plasmid pmt1, complete plasmid sequence.] [pn:unknown]
CONTIG462	34574067_c1_103	2609	8271	465	155	685	1.5(10)-67	Escherichia coli	b1865	[pn:datp pyrophosphohydrolase] [gn:ntpa]
CONTIG462	5208563_c1_104	2610	8272	768	256	1208	5.7(10)-123	Escherichia coli	b1864	[pn:hypothetical 26.4 kd protein in ruvc-asps intergenic region] [gn:yebc]
CONTIG462	24430416_c1_106	2611	8273	870	290	810	8.6(10)-81	Escherichia coli	b1861	[pn:holliday junction dna helicase ruva] [gn:ruva]
CONTIG462	116702_c2_110	2612	8274	3492	1164	228	8.5(10)-18	Salmonella typhimurium	AF007380	[PN:lambdaphage H tail component homolog] [DE:Salmonella typhimurium lambda phage K tail component homolog gene, partial cds, lambda phage L tail component homolog, copper-zinc superoxide dismutase (sodC), attachment and invasion prote
CONTIG462	1063557_c2_111	2613	8275	762	254	434	6.0(10)-41	Coxiella burnetii	Y15898	[de:coxiella burnetii plasmid qprs dna.] [pn:hypothetical protein] [gn:orf 248]

CONTIG462	26966_c2_113	2614	8276	3873	1291	1007	4.4(10)-101	Yersinia pestis	AF053947	[de:yersinia pestis plasmid pmt1, complete plasmid sequence.] [pn:phage lambda host specific protein.j]
CONTIG462	33828125_c2_120	2615	8277	606	202	736	6.0(10)-73	Escherichia coli	b1158	[pn:dna-invertase pin] [gn:pin]
CONTIG462	33752027_c2_125	2616	8278	543	181	750	2.0(10)-74	Escherichia coli	b1863	[pn:crossover junction endodeoxyribonuclease ruvc] [gn:ruvc]
CONTIG462	31875053_c2_129	2617	8279	477	159	442	8.6(10)-42	Escherichia coli	b1857	[pn:31.1 kd protein in msbb-ruvb intergenic region] [gn:yeb1]
CONTIG462	3393908_c3_137	2618	8280	855	285	319	9.4(10)-29	Yersinia pestis	AF053947	[de:yersinia pestis plasmid pmt1, complete plasmid sequence.] [pn:phage lambda minor tail protein l homolog]
CONTIG462	22541631_c3_148	2619	8281	1821	607	2868	7.2(10)-299	Escherichia coli	b1866	[pn:aspartyl-trna synthetase] [gn:asps]
CONTIG462	25431592_c3_149	2620	8282	1014	338	1623	6.2(10)-167	Escherichia coli	b1860	[pn:holliday junction dna helicase ruvb] [gn:ruvb]
CONTIG463	32605317_f1_1	2621	8283	591	197	428	2.6(10)-40	Shigella flexneri	P20102	mercuric resistance protein merd.
CONTIG463	4788505_f1_3	2622	8284	1323	441	1941	1.2(10)-200	Pseudomonas aeruginosa	U49101	or:pseudomonas aeruginosa pn:tnia gn:tnia lc:6735 re:8450
CONTIG463	13162887_f1_4	2623	8285	495	165	675	1.8(10)-66	Klebsiella pneumoniae	S32177	transposase - klebsiella pneumoniae
CONTIG463	12111286_f1_9	2624	8286	1563	521	2589	2.7(10)-269	Pseudomonas aeruginosa	Q57541	transposase for insertion sequences is1326/is1353.
CONTIG463	24644811_f1_15	2625	8287	1020	340	1583	1.1(10)-162	Corynebacterium glutamicum	Y14748	[de:corynebacterium glutamicum plasmid pcg4 integron (incg) sequence.] [gn:int]
CONTIG463	26614682_f2_25	2626	8288	510	170	810	8.6(10)-81	Escherichia coli	L29404	or:escherichia coli pn:mercuric reductase gn:mera lc:<1 re:663 di:direct sr:escherichia coli (strain dh1) dna nt:putative
CONTIG463	16141561_f2_27	2627	8289	927	309	1060	2.7(10)-107	Escherichia coli	L29404	or:escherichia coli lc:1273 re:1980 di:direct sr:escherichia coli (strain dh1) dna nt:urf2; putative



CONTIG463	11891882_f2_29	2628	8290	738	246	1268	2.6(10)-129	Salmonella ordonez	S34451	hypothetical protein (insertion sequence is261) - salmonellaordonez plasmid pip173
CONTIG463	24500431_f2_32	2629	8291	276	92	269	7.2(10)-23	Klebsiella pneumoniae	S32179	hypothetical protein 6 - klebsiella pneumoniae
CONTIG463	30494567_f2_37	2630	8292	204	68	111	5.7(10)-6	Ralstonia eutropha	D64144	or:ralstonia eutropha le:777 re:1793 di:direct sr:alcaligenes eutrophus (strain.nh9) plasmid:penh91 dna nt:orfa2
CONTIG463	4401693_f2_39	2631	8293	795	265	1327	1.3(10)-135	Insertion sequence IS1326	U38187	or:insertion sequence is1326 pn:istb le:1618 re:2403 di:direct nt:contains ntp binding domain motifs
CONTIG463	2838312_f2_44	2632	8294	348	116	258	2.7(10)-22	Citrobacter freundii	Z54241	or:citrobacter freundii pn:dna integrate gn:int le:<1 re:285 di:complement
CONTIG463	21667626_f2_49	2633	8295	3024	1008	4907	0	Escherichia coli	P13694	transposase for transposon tn3926.
CONTIG463	135818_f2_50	2634	8296	1032	344	1554	1.3(10)-159	Enterobacter aerogenes	U67194	or:enterobacter aerogenes pn:tnpa gn:tnpa le:10303 re:11307 di:complement
CONTIG463	25683166_f3_52	2635	8297	561	187	326	1.7(10)-29	Escherichia coli	L29404	or:escherichia coli le:1040 re:1276 di:direct sr:escherichia coli (strain dh1) dna nt:urf1, putative
CONTIG463	16462943_f3_57	2636	8298	873	291	1197	8.5(10)-122	Pseudomonas aeruginosa	U49101	or:pseudomonas aeruginosa pn:tnibdelta1 gn:tnibdelta1 le:5872 re:6732 di:complement nt:truncated version of tnib found in tn402, emb1
CONTIG463	9929075_f3_70	2637	8299	618	206	838	9.4(10)-84	Enterobacter cloacae	Y09025	resolvase family recombinase, tnpr,
CONTIG463	35629583_c1_84	2638	8300	435	145	708	5.5(10)-70	Escherichia coli	P10017	hypothetical 14.4 kd protein (orf 1).
CONTIG463	12398261_c1_85	2639	8301	447	149	667	1.2(10)-65	Escherichia coli	P10018	hypothetical 14.7 kd protein (orf 2).
CONTIG463	3416040_c1_87	2640	8302	930	310	1401	2.1(10)-143	Mycobacterium fortuitum	S10928	dihydropteroate synthase (ec 2.5.1.15) - mycobacteriumfortuitum transposon tn610
CONTIG463	10415891_c2_110	2641	8303	753	251	1335	2.0(10)-136	unidentified bacterium	X04555	or:unidentified bacterium sp.p10019 le:1080 re:1829 di:direct nt:aad(2) (aa 1-249)

CONTIG463	4723833_c2_111	2642	8304	387	129	581	1.6(10)-56	Plasmid pDGO100	L06418	or:plasmid pdgo100 pn:quaternary ammonium compound-resistance protein gn:suli le:1986 re:2333 di:direct sr:plasmid pdgo100 (clone: integron in7) dna nt:orf4; putative
CONTIG463	7050966_c2_113	2643	8305	603	201	889	3.7(10)-89	Pseudomonas aeruginosa	U12338	or:pseudomonas aeruginosa pn:unknown le:6192 re:6692 di:direct nt:orf5
CONTIG463	36520917_c2_123	2644	8306	411	137	151	5.9(10)-11	Xanthomonas sp.	S32799	hypothetical protein 1 - xanthomonas sp. transposon tn5053(fragment)
CONTIG463	33332337_c2_124	2645	8307	204	68	187	9.0(10)-15	Xanthomonas sp.	S32799	hypothetical protein 1 - xanthomonas sp. transposon tn5053(fragment)
CONTIG463	10582283_c2_125	2646	8308	210	70	94	0.00013	Xanthomonas sp.	S32799	hypothetical protein 1 - xanthomonas sp. transposon tn5053(fragment)
CONTIG463	26660405_c2_128	2647	8309	477	159	90	0.00519	Mycobacterium tuberculosis	AL009198	[de:mycobacterium tuberculosis sequence v004.] [pn:pgs-family protein] [gn:mtv004.01c] [nt:mtv004.01c, member of the m. tuberculosis pgfs]
CONTIG463	10970768_c3_130	2648	8310	411	137	374	1.3(10)-34	Escherichia coli	b2669	[pn:dna-binding protein stpa] [gn:stpa]
CONTIG463	9931525_c3_151	2649	8311	1548	516	1540	3.7(10)-158	Insertion sequence IS1353	U40482	or:insertion sequence is1353 le:671 re:1585 di:direct nt:orf6; possible alternate start site at nt 686
CONTIG463	21666540_c3_154	2650	8312	714	238	1128	1.8(10)-114	Transposon Tn1525	M12900	or:transposon tn1525 gn:p12 le:996 re:>1721 di:direct sr:transposon tn1525 dna nt:putative
CONTIG464	4333406_f1_3	2651	8313	1233	411	94	0.33	Gallus gallus	D88828	[de:gallus gallus mrna for chicken rabaptin-5, complete cds.] [pn:chicken rabaptin-5]
CONTIG464	4329843_f1_4	2652	8314	651	217	991	5.7(10)-100	Plasmid R478	U49054	or:plasmid r478 pn:terx gn:terx le:1800 re:2441 di:direct nt:shows similarity to r478 terz, terd, and tere
CONTIG464	3912918_f2_16	2653	8315	1152	384	873	1.8(10)-87	Serratia marcescens	U59239	or:serratia marcescens le:820 re:1413 di:complement nt:orf3
CONTIG464	568791_f2_29	2654	8316	186	62	279	1.6(10)-24	Escherichia coli	b3503	[pn:arsenate reductase] [gn:arsc]
CONTIG464	30157255_f2_31	2655	8317	279	93	99	0.00012	Escherichia coli	U95365	transposase, is5b,

CONTIG464	26770126_f3_35	2656	8318	1191	397	810	8.6(10)-81	Serratia marcescens	U59239	or:serratia marcescens le:<1 re:715 di:complement nt:orf4
CONTIG464	33402312_f3_36	2657	8319	1035	345	100	0.01499	Haemophilus influenzae	HI0023	[pn:citrate lyase beta chain] [gn:cite]
CONTIG464	12297782_f3_37	2658	8320	501	167	666	1.6(10)-65	Plasmid R478	U49054	or:plasmid r478 pn:terw gn:terw le:205 re:672 di:direct
CONTIG464	2140678_f3_38	2659	8321	747	249	944	5.5(10)-95	Plasmid R478	U49054	or:plasmid r478 pn:tery gn:tery le:1184 re:1777 di:direct
CONTIG464	4956283_f3_39	2660	8322	687	229	373	1.8(10)-34	Plasmid R478	U49054	or:plasmid r478 pn:tery gn:tery le:1184 re:1777 di:direct
CONTIG464	34194711_f3_40	2661	8323	1125	375	195	4.2(10)-15	Escherichia coli	b2073	[pn:hypothetical protein]
CONTIG464	672833_f3_43	2662	8324	1590	550	114	0.0014	Saccharomyces cerevisiae	YKL166C	[pn:camp-dependent protein kinase 3, catalytic chain] [gn:tpk3]
CONTIG464	4339135_f3_47	2663	8325	1026	342	1684	2.1(10)-173	Escherichia coli	b1370	[pn:insertion element is5 hypothetical protein] [gn:yi52_5]
CONTIG464	35212692_c1_54	2664	8326	1236	412	679	6.7(10)-67	Escherichia coli	b1650	[pn:hypothetical protein] [gn:nema]
CONTIG464	10563465_c1_59	2665	8327	420	140	102	9.3(10)-6	Escherichia coli	b2861	[pn:insertion element is2 hypothetical 13.4 kd protein]
CONTIG464	24783465_c2_79	2666	8328	786	262	315	2.5(10)-28	Bacillus subtilis	ykvO	[pn:hypothetical protein]
CONTIG464	21775383_c2_82	2667	8329	300	100	446	3.2(10)-42	Escherichia coli	D90774	or:escherichia coli gn:is5 le:13994 re:14362 di:direct sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:orf_id:o263#20; similar to [swissprot accession]
CONTIG464	22458580_c2_83	2668	8330	384	128	490	7.0(10)-47	Escherichia coli	b1371	[pn:hypothetical protein]
CONTIG464	3940943_c3_106	2669	8331	711	237	90	0.05299	Pseudomonas syringae	AF036929	[de:pseudomonas syringae disulfide oxidoreductase (dsba) gene, completecds.] [pn:disulfide oxidoreductase] [gn:dsba] [nt:dsba]
CONTIG464	10411407_c3_108	2670	8332	483	161	298	1.6(10)-26	Rhizobium sp.	S34667	hypothetical protein 140 - rhizobium sp.
CONTIG464	25572937_c3_109	2671	8333	1623	541	759	2.2(10)-75	Rhizobium sp.	P50360	hypothetical 29.3 kd protein in region 2 of sym plasmid (no1265).
CONTIG465	22837807_f1_36	2672	8334	828	276	1294	4.5(10)-132	Escherichia coli	b0166	[pn:2,3,4,5-tetrahydropyridine-2-carboxylate n-succinyltransferase]

CONTIG465	35971925_f1_37	2673	8335	423	141	640	9.0(10)-63	Escherichia coli	b0163	[pn:hypothetical 15.1 kd protein in htra-dapd intergenic region] [gn:yaeh]
CONTIG465	34236578_f2_78	2674	8336	2742	914	4041	0	Escherichia coli	b0167	[pn:uridylyltransferase] [gn:glnd]
CONTIG465	33407706_f2_87	2675	8337	747	249	1114	5.2(10)-113	Escherichia coli	b0159	[pn:pfs protein] [gn:pfs]
CONTIG465	13067707_f2_89	2676	8338	723	241	886	7.7(10)-89	Escherichia coli	b0157	[pn:hypothetical protein in heml-pfs intergenic region] [gn:yads]
CONTIG465	24470635_f3_124	2677	8339	831	277	1239	3.0(10)-126	Escherichia coli	b0168	[pn:methionine aminopeptidase] [gn:map]
CONTIG465	12698840_f3_131	2678	8340	840	280	1023	2.2(10)-103	Escherichia coli	b0158	[pn:hypothetical 29.4 kd protein in heml-pfs intergenic region] [gn:yadt]
CONTIG465	35625650_c1_134	2679	8341	360	120	403	1.2(10)-37	Escherichia coli	b0155	[pn:hypothetical protein in heml-pfs intergenic region] [gn:yadq]
CONTIG465	22773593_c1_141	2680	8342	1545	515	2268	2.7(10)-235	Escherichia coli	b0160	[pn:deoxyguanosinetriphosphate triphosphohydrolase] [gn:dgt]
CONTIG465	25476662_c1_142	2681	8343	1491	497	1973	5.0(10)-204	Escherichia coli	b0161	[pn:heat shock protein htra] [gn:htra]
CONTIG465	26735086_c1_151	2682	8344	648	216	651	6.2(10)-64	Escherichia coli	b0171	[pn:uridine 5'''-monophosphate kinase] [gn:pyrh]
CONTIG465	9776538_c1_152	2683	8345	582	194	769	1.8(10)-76	Escherichia coli	b0172	[pn:ribosome recycling factor] [gn:frr]
CONTIG465	31797711_c1_154	2684	8346	1428	476	1850	5.4(10)-191	Escherichia coli	b0176	[pn:hypothetical protein in cdsa 3'''region] [gn:yael]
CONTIG465	31445208_c1_155	2685	8347	585	195	689	5.7(10)-68	Escherichia coli	b0180	[pn:3r-hydroxymyristol acyl carrier protein dehydrase] [gn:fabz]
CONTIG465	25431660_c1_156	2686	8348	792	264	1258	2.8(10)-128	Escherichia coli	b0181	[pn:udp-n-acetylglucosamine o-acyltransferase] [gn:lpax]
CONTIG465	26822803_c1_159	2687	8349	3501	1167	5547	0	Escherichia coli	b0184	[pn:dna polymerase iii, alpha chain] [gn:dnae]
CONTIG465	13870910_c1_160	2688	8350	972	324	1391	2.3(10)-142	Escherichia coli	b0185	[pn:acetyl-coenzyme a carboxylase carboxyl transferase subunit alpha] [gn:acca]
CONTIG465	20885086_c2_163	2689	8351	372	124	480	8.0(10)-46	Escherichia coli	b0156	[pn:hypothetical 12.1 kd protein in heml-pfs intergenic region] [gn:yadr]
CONTIG465	32074135_c2_177	2690	8352	777	259	1210	3.6(10)-123	Escherichia coli	b0169	[pn:30s ribosomal protein s2] [gn:rpsb]

CONTIG465	22011677_c2_178	2691	8353	936	312	1253	9.9(10)-128	Escherichia coli	b0170	[pn:elongation factor ts] [gn:tsf]
CONTIG465	12985030_c2_179	2692	8354	330	110	438	2.2(10)-41	Escherichia coli	b0171	[pn:uridine 5'-monophosphate kinase] [gn:pyrh]
CONTIG465	3958312_c2_181	2693	8355	768	256	1206	9.5(10)-123	Escherichia coli	b0174	[pn:hypothetical protein] [gn:yaes]
CONTIG465	24634376_c2_182	2694	8356	861	287	1074	9.1(10)-109	Escherichia coli	b0175	[pn:phosphatidate cytidyltransferase] [gn:cdsa]
CONTIG465	4554813_c2_184	2695	8357	1449	483	2171	5.2(10)-225	Escherichia coli	b0177	[pn:hypothetical protein] [gn:yaet]
CONTIG465	25441525_c2_185	2696	8358	1002	334	1484	3.2(10)-152	Escherichia coli	b0177	[pn:hypothetical protein] [gn:yaet]
CONTIG465	2595412_c2_186	2697	8359	561	187	605	4.5(10)-59	Escherichia coli	b0178	[pn:histone-like protein hlp-1 precursor] [gn:hlpal]
CONTIG465	9869053_c2_187	2698	8360	1029	343	1513	2.7(10)-155	Escherichia coli	b0179	[pn:udp-3-o-3-hydroxymyristoyl glucosamine n-acyltransferase] [gn:lpdx]
CONTIG465	31927308_c2_189	2699	8361	630	210	927	3.5(10)-93	Escherichia coli	b0183	[pn:ribonuclease hii] [gn:rmhb]
CONTIG465	4494032_c2_193	2700	8362	2136	712	3424	0	Escherichia coli	b0186	[pn:lysine decarboxylase] [gn:ldcc]
CONTIG465	14664812_c2_194	2701	8363	402	134	538	5.7(10)-52	Escherichia coli	b0187	[pn:hypothetical protein]
CONTIG465	12114076_c3_202	2702	8364	1200	400	1846	1.3(10)-190	Escherichia coli	b0162	[pn:hypothetical 44.3 kd protein in htra-dapd intergenic region] [gn:yaeg]
CONTIG465	33603957_c3_210	2703	8365	1272	424	1767	3.3(10)-182	Escherichia coli	b0173	[pn:hypothetical protein in fir 3' region] [gn:yaem]
CONTIG465	12300082_c3_219	2704	8366	1176	392	1661	5.7(10)-171	Escherichia coli	b0182	[pn:lipid-a-disaccharide synthase] [gn:lpbx]
CONTIG465	14875750_c3_225	2705	8367	1077	359	1052	2.0(10)-106	Escherichia coli	b0188	[pn:cell cycle protein mes] [gn:mesj]
CONTIG466	4792812_fl_1	2706	8368	273	91	426	4.2(10)-40	Escherichia coli	b3704	[pn:mase p, protein component] [gn:mpa]
CONTIG466	25676928_fl_2	2707	8369	1137	379	1798	1.8(10)-185	Escherichia coli	b3705	[pn:60 kd protein] [gn:yide]
CONTIG466	14586582_fl_3	2708	8370	1377	459	2050	3.5(10)-212	Escherichia coli	b3706	[pn:50 kd protein] [gn:thdf]

CONTIG466	4697650_f1_4	2709	8371	1287	429	1260	1.8(10)-128	Escherichia coli	b3710	[pn:hypothetical 41.5 kd protein in tnab 3'''region] [gn:yidy]
CONTIG466	14954081_f1_5	2710	8372	777	259	802	6.2(10)-80	Escherichia coli	b3712	[pn:hypothetical 28.0 kd protein in tnab-bglb intergenic region]
CONTIG466	36503530_f1_6	2711	8373	603	201	867	8.0(10)-87	Escherichia coli	b3713	[pn:hypothetical protein] [gn:yief]
CONTIG466	14729535_f2_30	2712	8374	606	202	629	1.3(10)-61	Escherichia coli	b3705	[pn:60 kd protein] [gn:yidc]
CONTIG466	12601081_f2_33	2713	8375	306	102	178	4.7(10)-13	Escherichia coli	A38160	thdf protein - escherichia coli (fragment)
CONTIG466	16454200_f2_36	2714	8376	1029	343	1163	3.3(10)-118	Escherichia coli	b3711	[pn:hypothetical transcriptional regulator in tnab-bglb intergenic region] [gn:yidz]
CONTIG466	2036331_f3_68	2715	8377	288	96	236	5.7(10)-20	Haemophilus influenzae	H11000	[pn:hypothetical protein]
CONTIG466	20317500_f3_74	2716	8378	741	247	1004	2.3(10)-101	Escherichia coli	b3715	[pn:hypothetical protein] [gn:yieh]
CONTIG466	34587752_c1_100	2717	8379	594	198	198	6.2(10)-16	Escherichia coli	b0530	[pn:hypothetical protein] [gn:sfma]
CONTIG466	15759625_c1_107	2718	8380	963	321	1357	9.5(10)-139	Escherichia coli	b3727	[pn:phosphate transport system permease protein pstc] [gn:pstc]
CONTIG466	20573253_c1_109	2719	8381	831	277	1170	6.2(10)-119	Escherichia coli	b3724	[pn:peripheral membrane protein u] [gn:phou]
CONTIG466	28557318_c2_126	2720	8382	399	133	598	2.5(10)-58	Escherichia coli	b3738	[pn:atp synthase f0 subunit a] [gn:atpb]
CONTIG466	34647257_c2_127	2721	8383	555	185	782	8.0(10)-78	Escherichia coli	b3735	[pn:atp synthase f1 delta subunit] [gn:atph]
CONTIG466	34614218_c2_128	2722	8384	1554	518	2381	2.8(10)-247	Escherichia coli	b3734	[pn:atp synthase f1 alpha subunit] [gn:atpa]
CONTIG466	35831955_c2_129	2723	8385	915	305	1381	2.7(10)-141	Escherichia coli	b3733	[pn:atp synthase f1 gamma subunit] [gn:atpg]
CONTIG466	25682955_c2_130	2724	8386	432	144	661	5.4(10)-65	Escherichia coli	b3731	[pn:atp synthase f1 epsilon subunit] [gn:atpc]
CONTIG466	22843762_c2_131	2725	8387	1425	475	2082	1.3(10)-215	Escherichia coli	b3730	[pn:udp-n-acetylglucosamine pyrophosphorylase] [gn:glmu]
CONTIG466	4102305_c2_132	2726	8388	1845	615	2868	7.2(10)-299	Escherichia coli	b3729	[pn:glutamine amidotransferase]
CONTIG466	11142193_c2_139	2727	8389	786	262	1220	3.1(10)-124	Escherichia coli	b3725	[pn:phosphate transport atp-binding protein pstb] [gn:pstb]

CONTIG466	25989638_c2_141	2728	8390	1377	459	1938	2.6(10)-200	Escherichia coli	b3714	[pn:hypothetical 46.9 kd protein in tnab-bglb intergenic region]
CONTIG466	10659431_c2_147	2729	8391	480	160	90	0.11	Saccharomyces cerevisiae	YNR044W	[pn:a-agglutinin anchor subunit] [gn:agal]
CONTIG466	26585912_c3_150	2730	8392	258	86	130	1.0(10)-8	Haemophilus influenzae	HI0484	[pn:atp synthase c chain] [gn:atpc]
CONTIG466	13082037_c3_151	2731	8393	507	169	601	1.2(10)-58	Escherichia coli	b3736	[pn:atp synthase f0 subunit b] [gn:atpf]
CONTIG466	29964050_c3_155	2732	8394	1506	502	2132	7.0(10)-221	Escherichia coli	b3732	[pn:atp synthase f1 beta subunit] [gn:atpd]
CONTIG466	5313943_c3_158	2733	8395	696	232	348	7.9(10)-32	Escherichia coli	b3143	[pn:hypothetical 25.7 kd fimbrial chaperone in agai- mtr intergeni] [gn:yrai]
CONTIG466	35370308_c3_159	2734	8396	2538	846	1405	7.7(10)-144	Escherichia coli	b0532	[pn:hypothetical protein] [gn:sfmd]
CONTIG466	5267193_c3_160	2735	8397	1086	362	177	3.2(10)-11	Escherichia coli	b0941	[pn:hypothetical protein]
CONTIG466	19572182_c3_161	2736	8398	1074	358	1554	1.3(10)-159	Escherichia coli	b3728	[pn:periplasmic phosphate-binding protein] [gn:psts]
CONTIG466	36617687_c3_164	2737	8399	909	303	1153	3.8(10)-117	Escherichia coli	b3726	[pn:phosphate transport system permease protein psta] [gn:psta]
CONTIG467	1275005_f1_42	2738	8400	303	101	91	0.005	Saccharomyces cerevisiae	YIR019C	[pn:extracellular alpha-1,4-glucan glucosidase] [gn:sta]
CONTIG467	33461682_c1_117	2739	8401	2181	727	446	7.5(10)-42	Bacillus subtilis	yknV	[pn:hypothetical protein]
CONTIG467	6835407_c2_118	2740	8402	18048	6016	2534	3.1(10)-269	Synechocystis sp.	S76109	[PN:hypothetical protein] [OR:Synechocystis sp.] [SR:PCC 6803, PCC 6803] [SR:PCC 6803, ] cyad protein.
CONTIG467	35742257_c3_137	2741	8403	1197	399	360	4.2(10)-33	Bordetella pertussis	P11091	
CONTIG467	6054666_c3_139	2742	8404	783	261	143	1.8(10)-7	Klebsiella pneumoniae	L23111	or:klebsiella pneumoniae pn:fimbrial adhesin gn:fimk le:1139 re:2380 di:direct sr:klebsiella pneumoniae
CONTIG468	5133412_f1_2	2743	8405	330	110	391	2.2(10)-36	Escherichia coli	b0379	[pn:hypothetical protein]
CONTIG468	20397016_f1_8	2744	8406	1335	445	1652	5.2(10)-170	Escherichia coli	b0376	[pn:yaih]

CONTIG468	33212805_f1_13	2745	8407	1728	576	299	4.7(10)-26	Prevotella melaninogenica	U27587	or:prevotella melaninogenica pn:hemolysin a gn:hly le:188 re:1186 di:direct nt:zymogram analysis confirms the hemolytic glycoprotein x precursor.
CONTIG468	24477091_f1_21	2746	8408	876	292	95	0.14999	Equine herpesvirus 1	P28968	[pn:membrane protein]
CONTIG468	3986527_f1_23	2747	8409	1155	385	183	6.0(10)-12	Helicobacter pylori	HP0567	[pn:hypothetical 41.5 kd protein in psif-proc intergenic region] [gn:yaic]
CONTIG468	21878332_f1_24	2748	8410	1821	607	279	1.0(10)-23	Escherichia coli	b0385	[pn:hypothetical 27.4 kd protein in fec1-fimb intergenic region]
CONTIG468	16025932_f1_26	2749	8411	789	263	1178	8.8(10)-120	Escherichia coli	b4306	[pn:d-alanine] [gn:ddla]
CONTIG468	20432_f2_36	2750	8412	1221	407	1549	4.2(10)-159	Escherichia coli	b0381	[pn:delta-aminolevulinic acid dehydratase] [gn:hemb]
CONTIG468	4460882_f2_43	2751	8413	1095	365	1513	2.7(10)-155	Escherichia coli	b0369	[pn:hypothetical protein] [gn:ycir]
CONTIG468	21537927_f2_59	2752	8414	1959	653	644	3.3(10)-63	Escherichia coli	b1285	
CONTIG468	864080_f2_64	2753	8415	183	61	129	3.6(10)-8	Podospira anserina	X55026	or:mitochondrion podospira anserina le:44740 re:45549 di:complement sr:podospira anserina nt:orf16; no atg start codon; author-
CONTIG468	26830152_f2_65	2754	8416	444	148	446	3.2(10)-42	Escherichia coli	U70214	or:escherichia coli le:102767 re:103075 di:complement nt:hypothetical protein
CONTIG468	12915832_f3_107	2755	8417	423	141	676	1.3(10)-66	Escherichia coli	U70214	or:escherichia coli le:102415 re:102852 di:complement nt:hypothetical protein
CONTIG468	32227262_c1_109	2756	8418	522	174	703	1.8(10)-69	Escherichia coli	b0257	[pn:hypothetical protein]
CONTIG468	5078942_c1_111	2757	8419	1530	510	1823	3.8(10)-188	Escherichia coli	b0258	[pn:hypothetical 43.2 kd protein in per-argf intergenic region] [gn:ykfc]
CONTIG468	14880168_c1_112	2758	8420	210	70	100	1.5(10)-5	Escherichia coli	b4286	[pn:hypothetical protein]
CONTIG468	35267890_c1_131	2759	8421	957	319	1238	3.8(10)-126	Escherichia coli	b0368	[pn:hypothetical protein] [gn:taud]
CONTIG468	2931593_c1_138	2760	8422	432	144	549	4.0(10)-53	Escherichia coli	b0374	[pn:hypothetical protein] [gn:yaiu]



CONTIG468	36148311_c1_142	2761	8423	1239	413	1814	3.5(10)-187	Escherichia coli	b0377	[pn:sbma protein] [gn:sbma]
CONTIG468	15664812_c1_143	2762	8424	1110	370	1475	3.0(10)-151	Escherichia coli	b0378	[pn:hypothetical protein] [gn:yaiv]
CONTIG468	16532656_c2_149	2763	8425	396	132	404	9.1(10)-38	Escherichia coli	b0255	[pn:hypothetical protein]
CONTIG468	32066327_c2_160	2764	8426	1425	475	2146	2.2(10)-222	Escherichia coli	b0260	[pn:hypothetical 51.5 kd transport protein in per-argf intergenic region] [gn:ykfd]
CONTIG468	3191_c2_161	2765	8427	1038	346	1487	1.6(10)-152	Escherichia coli	b0365	[pn:hypothetical protein]
CONTIG468	36229058_c2_164	2766	8428	846	282	793	5.5(10)-79	Escherichia coli	b0367	[pn:hypothetical protein] [gn:tauc]
CONTIG468	585340_c2_170	2767	8429	2391	797	996	1.7(10)-100	Escherichia coli	b0374	[pn:hypothetical protein] [gn:yaiv]
CONTIG468	30136001_c2_171	2768	8430	639	213	510	5.4(10)-49	Escherichia coli	b0375	[pn:hypothetical protein] [gn:yaiv]
CONTIG468	16176655_c3_178	2769	8431	507	169	797	2.1(10)-79	Escherichia coli	Y07545	[de:e.coli plasmid p0157 dna, 5'''-region of the ehc-hemolysin operon.] [gn:orf5]
CONTIG468	31803342_c3_179	2770	8432	276	92	93	0.00063	Xenopus laevis	S31719	proline-rich protein - african clawed frog
CONTIG468	34645806_c3_189	2771	8433	951	317	1268	2.6(10)-129	Escherichia coli	b0261	[pn:hypothetical 33.4 kd protein in per-argf intergenic region]
CONTIG468	17089012_c3_191	2772	8434	783	261	1139	1.2(10)-115	Escherichia coli	b0366	[pn:hypothetical protein] [gn:taub]
CONTIG468	267338_c3_202	2773	8435	252	84	297	2.0(10)-26	Escherichia coli	b0380	[pn:hypothetical protein]
CONTIG468	25910005_c3_206	2774	8436	672	224	524	1.8(10)-50	Escherichia coli	b1033	[pn:hypothetical 43.9 kd protein in msyb-htrb intergenic region]
CONTIG469	25992268_fl_1	2775	8437	1167	389	1708	6.0(10)-176	Escherichia coli	b4375	[gn:ycee] [pn:peptide-chain-release factor 3]
CONTIG469	24632962_fl_10	2776	8438	732	244	1185	1.6(10)-120	Escherichia coli	b4384	[pn:purine-nucleoside phosphorylase] [gn:deod]
CONTIG469	24744157_fl_18	2777	8439	1389	463	2188	8.3(10)-227	Escherichia coli	b4389	[pn:sms protein] [gn:rada]
CONTIG469	1206557_fl_22	2778	8440	906	302	332	3.8(10)-30	Escherichia coli	b3243	[pn:hypothetical protein] [gn:yhcs]

CONTIG469	34478458_f1_27	2779	8441	414	138	423	8.9(10)-40	Escherichia coli	b4393	[pn:trpr] [gn:trpr]
CONTIG469	19729541_f1_28	2780	8442	432	144	192	2.7(10)-15	Escherichia coli	J01715	or:escherichia coli le:765 re:1178 di:direct sr:escherichia coli dna [1],[2] and mma [2]
CONTIG469	34179716_f1_29	2781	8443	571	191	775	4.5(10)-77	Escherichia coli	b4395	[pn:probable phosphoglycerate mutase 2] [gn:gpmh]
CONTIG469	22460465_f2_30	2782	8444	642	214	726	7.0(10)-72	Escherichia coli	b4376	[pn:periplasmic protein] [gn:osmy]
CONTIG469	35242941_f2_32	2783	8445	327	109	292	6.7(10)-26	Escherichia coli	P39408	hypothetical 28.9 kd protein in osmy-deoc intergenic region.
CONTIG469	1353888_f2_33	2784	8446	468	156	666	1.6(10)-65	Escherichia coli	b4378	[pn:yjiv]
CONTIG469	14730178_f2_39	2785	8447	1239	413	2087	4.2(10)-216	Escherichia coli	b4383	[pn:phosphotomutase] [gn:deob]
CONTIG469	3381262_f2_48	2786	8448	1071	357	1457	2.3(10)-149	Escherichia coli	b4388	[pn:phosphoserine phosphatase] [gn:serb]
CONTIG469	14588437_f2_51	2787	8449	1770	590	115	0.0004	Pseudomonas aeruginosa	JQ0133	hypothetical 26.4k protein - pseudomonas aeruginosa
CONTIG469	36428938_f3_59	2788	8450	1113	371	1490	7.5(10)-153	Escherichia coli	b4377	[pn:hypothetical 39.8 kd protein in osmy-deoc intergenic region] [gn:yjju]
CONTIG469	23476055_f3_65	2789	8451	948	316	1135	3.2(10)-115	Escherichia coli	b4381	[pn:deoxyribose-phosphate aldolase] [gn:deoc]
CONTIG469	12898503_f3_66	2790	8452	1422	474	2050	3.5(10)-212	Escherichia coli	b4382	[pn:thymidine phosphorylase] [gn:deoa]
CONTIG469	16303762_f3_77	2791	8453	258	86	95	0.00129	Nephila clavipes	AF027735	[de:nephila clavipes minor ampullate silk protein misp1 mma, partial cds.] [pn:minor ampullate silk protein misp1]
CONTIG469	10634625_f3_78	2792	8454	1236	412	1967	2.2(10)-203	Escherichia coli	b4390	[pn:transcriptional regulator nadr] [gn:nadr]
CONTIG469	15036641_f3_84	2793	8455	1983	661	3026	0	Escherichia coli	b4392	[pn:soluble lytic transglycosylase] [gn:slt]
CONTIG469	1011_c1_87	2794	8456	567	189	644	3.3(10)-63	Escherichia coli	b4394	[pn:hypothetical 18.6 kd protein in trpr-gpmh intergenic region]
CONTIG469	23849007_c1_97	2795	8457	1041	347	192	3.5(10)-13	Escherichia coli	b4051	[pn:quinone oxidoreductase] [gn:qor]

CONTIG469	5094427_c2_141	2796	8458	321	107	133	6.2(10)-9	Escherichia coli	b3215	[pn:hypothetical 25.3 kd fimbrial chaperone in gltf- nant intergenic region] [gn:yhca]
CONTIG469	32673426_c2_148	2797	8459	1629	543	2363	2.3(10)-245	Escherichia coli	b4380	[pn:hypothetical 58.0 kd protein in osmy-deoc intergenic region]
CONTIG469	32285036_c3_161	2798	8460	1764	588	2632	7.4(10)-274	Escherichia coli	b4391	[pn:abc transporter in nadr-slt intergenic region] [gn:yijk]
CONTIG469	33865627_c3_169	2799	8461	663	221	788	1.8(10)-78	Escherichia coli	b4387	[pn:smp protein precursor] [gn:smp]
CONTIG469	5339642_c3_170	2800	8462	1050	350	1653	4.0(10)-170	Escherichia coli	b4386	[pn:lipoate-protein ligase a] [gn:lpia]
CONTIG469	5193818_c3_171	2801	8463	696	232	223	1.3(10)-18	Escherichia coli	b3219	[pn:hypothetical protein] [gn:yhcf]
CONTIG469	4804068_c3_172	2802	8464	699	233	252	1.2(10)-21	Escherichia coli	b3219	[pn:hypothetical protein] [gn:yhcf]
CONTIG469	33790903_c3_181	2803	8465	864	288	1010	5.5(10)-102	Escherichia coli	b4379	[pn:hypothetical 31.5 kd protein in osmy-deoc intergenic region] [gn:yijw]
CONTIG47	25814591_f2_2	2804	8466	315	105	395	8.3(10)-37	Escherichia coli	b1183	[pn:umud protein] [gn:umud]
CONTIG470	19714505_f1_5	2805	8467	570	190	355	1.3(10)-32	Escherichia coli	b0892	[pn:hypothetical protein in sers 5''' region] [gn:yca]
CONTIG470	24640653_f1_8	2806	8468	2451	817	3753	0	Escherichia coli	b0894	[pn:anaerobic dimethyl sulfoxide reductase chain a precursor] [gn:dmsa]
CONTIG470	14181530_f1_10	2807	8469	1152	384	1451	1.0(10)-148	Escherichia coli	b0898	[pn:hypothetical protein in dmsc 3''' region] [gn:yca]
CONTIG470	10572341_f1_20	2808	8470	768	256	838	9.4(10)-84	Escherichia coli	b0906	[pn:hypothetical protein] [gn:yca]
CONTIG470	16836093_f1_21	2809	8471	1095	365	1718	5.2(10)-177	Escherichia coli	b0907	[pn:phosphoserine aminotransferase] [gn:serc]
CONTIG470	32422910_f2_33	2810	8472	468	156	708	5.5(10)-70	Escherichia coli	b0889	[pn:leucine-responsive regulatory protein] [gn:lrp]
CONTIG470	33882161_f2_34	2811	8473	3705	1235	2299	0	Escherichia coli	b0890	[pn:cell division protein ftsk] [gn:flsk]
CONTIG470	14572206_f2_41	2812	8474	621	207	1113	6.7(10)-113	Escherichia coli	b0895	[pn:anaerobic dimethyl sulfoxide reductase chain b] [gn:dmsb]
CONTIG470	9802290_f2_49	2813	8475	495	165	443	6.7(10)-42	Escherichia coli	b0910	[pn:cytidylate kinase] [gn:cmk]

CONTIG470	3297258_f2_50	2814	8476	327	109	469	1.2(10)-44	Escherichia coli	b0912	[pn:integration host factor beta-subunit] [gn:himd]
CONTIG470	31304137_f2_55	2815	8477	993	331	1183	2.6(10)-120	Escherichia coli	b0915	[pn:hypothetical 35.6 kd protein in msba-kdsb intergenic region] [gn:ycab]
CONTIG470	14072667_f2_57	2816	8478	780	260	1130	1.1(10)-114	Escherichia coli	b0918	[pn:3-deoxy-manno-octulosonate cytidyltransferase] [gn:kdsb]
CONTIG470	4803443_f3_63	2817	8479	624	208	923	9.3(10)-93	Escherichia coli	b0891	[pn:outer membrane lipoprotein carrier protein precursor] [gn:lola]
CONTIG470	2536578_f3_64	2818	8480	1182	394	1760	1.8(10)-181	Escherichia coli	b0892	[pn:hypothetical protein in sers 5'''region] [gn:ycal]
CONTIG470	22845400_f3_65	2819	8481	1314	438	2080	2.2(10)-215	Escherichia coli	b0893	[pn:seryl-trna synthetase] [gn:sers]
CONTIG470	14473425_f3_68	2820	8482	936	312	1206	9.5(10)-123	Escherichia coli	b0896	[pn:anaerobic dimethyl sulfoxide reductase chain c] [gn:dmse]
CONTIG470	275088_f3_76	2821	8483	1296	432	1752	1.3(10)-180	Escherichia coli	b0908	[pn:3-phosphoshikimate 1-carboxyvinyltransferase] [gn:aroa]
CONTIG470	36526905_f3_77	2822	8484	444	148	564	1.0(10)-54	Escherichia coli	b0910	[pn:cytidylate kinase] [gn:cmk]
CONTIG470	4490692_f3_78	2823	8485	1683	561	2321	6.7(10)-241	Escherichia coli	b0911	[pn:30s ribosomal protein s1] [gn:rpas]
CONTIG470	7244037_f3_80	2824	8486	2280	760	2325	2.5(10)-241	Escherichia coli	b0913	[pn:hypothetical protein in msba 5'''region] [gn:ycal]
CONTIG470	25970202_f3_81	2825	8487	1749	583	2574	1.0(10)-267	Escherichia coli	b0914	[pn:probable transport atp-binding protein msba] [gn:msba]
CONTIG470	24507201_f3_82	2826	8488	1245	415	1670	6.4(10)-172	Escherichia coli	b0916	[pn:hypothetical protein]
CONTIG470	6058562_f3_83	2827	8489	228	76	264	6.2(10)-23	Escherichia coli	b0917	[pn:hypothetical protein]
CONTIG470	26595130_f3_85	2828	8490	900	300	1423	9.5(10)-146	Escherichia coli	b0919	[pn:hypothetical protein]
CONTIG470	30363283_c1_105	2829	8491	777	259	1312	5.5(10)-134	Escherichia coli	b0902	[pn:pyruvate formate-lyase 1 activating enzyme] [gn:pfla]
CONTIG470	36621094_c2_127	2830	8492	302	100	403	1.2(10)-37	Escherichia coli	b0920	[pn:hypothetical protein in kdsb-kicb intergenic region] [gn:ycbc]
CONTIG470	24084592_c2_148	2831	8493	1968	656	2744	1.0(10)-285	Escherichia coli	b0905	[pn:hypothetical protein] [gn:ycal]
CONTIG470	22376412_c2_149	2832	8494	969	323	1304	3.8(10)-133	Escherichia coli	b0904	[pn:probable formate transporter] [gn:foca]

CONTIG470	30736688_c2_150	2833	8495	2298	766	3671	0	Escherichia coli	b0903	[pn:formate acetyltransferase I] [gn:pf1b]
CONTIG470	15100141_c2_165	2834	8496	318	106	94	0.00052	Caenorhabditis elegans	Z74033	or:caenorhabditis elegans pn:f38b7.3 le:join(22734 re:22880,22931 di:direct nt:cdna est yk117e9.5 comes from this gene; cdna est
CONTIG471	1992161_f1_6	2835	8497	1689	563	1476	2.2(10)-151	Escherichia coli	b3767	[pn:acetohydroxy acid synthase ii, large subunit] [gn:ilvg_1]
CONTIG471	4147812_f1_10	2836	8498	186	62	92	0.0016	Escherichia coli	M32253	or:escherichia coli gn:ilvd le:3652 re:5499 di:direct sr:escherichia coli (strain k-12) dna
CONTIG471	32145043_f2_26	2837	8499	222	74	235	7.5(10)-20	synthetic construct	M15619	or:artificial sequence le:29 re:>232 di:direct sr:e.coli (strain se5000) synthetic dna, clone pkb1 nt:orf16-lacZ fusion protein
CONTIG471	31662801_f2_30	2838	8500	390	130	551	2.3(10)-53	Escherichia coli	b3764	[pn:hypothetical 13.1 kd protein in pssr-ilvl intergenic region] [gn:yife]
CONTIG471	3417882_f2_38	2839	8501	1560	520	2459	1.6(10)-255	Escherichia coli	b3772	[pn:threonine dehydratase biosynthetic] [gn:ilva]
CONTIG471	22444682_f2_41	2840	8502	1503	501	2319	1.1(10)-240	Escherichia coli	b3774	[pn:ketol-acid reductoisomerase] [gn:ilvc]
CONTIG471	35986463_f3_48	2841	8503	432	144	113	6.2(10)-7	Escherichia coli	M87049	or:escherichia coli gn:o137 le:1223 re:1636 di:direct
CONTIG471	15819651_f3_55	2842	8504	288	96	381	2.5(10)-35	Escherichia coli	b3769	[pn:acetohydroxy acid synthase ii, small subunit] [gn:ilvm]
CONTIG471	24426711_f3_56	2843	8505	942	314	1571	2.0(10)-161	Escherichia coli	b3770	[pn:branched-chain amino-acid aminotransferase] [gn:ilve]
CONTIG471	23540917_f3_57	2844	8506	1854	618	2677	1.3(10)-278	Escherichia coli	b3771	[pn:dihydroxyacid dehydratase] [gn:ilvd]
CONTIG471	30352281_c1_66	2845	8507	426	142	437	2.8(10)-41	Escherichia coli	b3605	[pn:lctd] [gn:lidd]
CONTIG471	14845313_c1_67	2846	8508	1428	476	893	1.3(10)-89	Pseudomonas putida	U10895	or:pseudomonas putida pn:pcak gn:pcak le:261 re:1607 di:direct
CONTIG471	4567968_c1_68	2847	8509	483	161	787	2.3(10)-78	Escherichia coli	b3606	[pn:hypothetical 17.7 kd protein in lctd-cyse intergenic region] [gn:yibk]
CONTIG471	14959652_c1_81	2848	8510	855	285	776	3.5(10)-77	Escherichia coli	b3762	[pn:hypothetical 22.4 kd protein in trpt-pssr intergenic region] [gn:yifa]
CONTIG471	6932937_c2_95	2849	8511	909	303	1490	7.5(10)-153	Escherichia coli	b3773	[pn:transcriptional activator protein ilvy] [gn:ilvy]

CONTIG471	12773450_c3_117	2850	8512	228	76	99	0.00014	Escherichia coli	b3605	[pn:lctd] [gn:lldd]
CONTIG471	33386290_c3_138	2851	8513	1551	517	1973	5.0(10)-204	Escherichia coli	b3765	[pn:hypothetical 56.2 kd protein in pssr-1vl intergenic region] [gn:yifb]
CONTIG471	12781963_c3_148	2852	8514	189	63	101	1.2(10)-5	Homo sapiens	A44803	pgl protein - human (fragment)
CONTIG472	24350086_fl_14	2853	8515	1488	496	1897	5.7(10)-196	Escherichia coli	b1439	[pn:hypothetical protein]
CONTIG472	35428135_fl_21	2854	8516	267	89	272	9.0(10)-24	Escherichia coli	b1445	[pn:hypothetical protein]
CONTIG472	22113325_f2_45	2855	8517	834	278	252	1.2(10)-21	Escherichia coli	P52646	hypothetical 5.0 kd protein in hslj 3'region.
CONTIG472	22461461_f2_53	2856	8518	1461	487	2002	4.2(10)-207	Escherichia coli	b1444	[pn:hypothetical protein]
CONTIG472	4806693_f2_62	2857	8519	243	81	331	5.0(10)-30	Escherichia coli	b1446	[pn:hypothetical protein]
CONTIG472	4348510_f2_64	2858	8520	585	195	133	4.7(10)-9	Escherichia coli	b1434	[pn:hypothetical protein]
CONTIG472	26070290_f2_65	2859	8521	1101	367	1462	7.0(10)-150	Escherichia coli	b1449	[pn:hypothetical protein]
CONTIG472	24251918_f3_85	2860	8522	798	266	123	2.1(10)-5	Bacillus subtilis	ybfK	[pn:hypothetical protein]
CONTIG472	5098387_f3_96	2861	8523	1689	563	1524	1.8(10)-156	Escherichia coli	b3544	[pn:periplasmic dipeptide transport protein precursor] [gn:dppa]
CONTIG472	25519182_f3_101	2862	8524	714	238	469	1.2(10)-44	Escherichia coli	b1450	[pn:hypothetical protein]
CONTIG472	10944025_c1_106	2863	8525	693	231	760	1.7(10)-75	Escherichia coli	b1448	[pn:hypothetical protein]
CONTIG472	135836_c1_119	2864	8526	351	117	461	8.4(10)-44	Escherichia coli	b1797	[pn:hypothetical protein]
CONTIG472	23957058_c1_120	2865	8527	381	127	384	1.2(10)-35	Escherichia coli	b1797	[pn:hypothetical protein]
CONTIG472	23728427_c1_121	2866	8528	186	62	129	1.3(10)-8	Escherichia coli	b1796	[pn:hypothetical protein]
CONTIG472	6366068_c1_130	2867	8529	1023	341	1031	3.2(10)-104	Pseudomonas aeruginosa	Y10528	[PN:cyanide insensitive terminal oxidase] [GN:ciOB] [DE:P.aeruginosa cioA and cioB genes.] [LE:1746] [RE:2753] [DI:direct]

CONTIG472	6131633_c1_138	2868	8530	243	81	249	5.4(10)-21	Escherichia coli	b1345	[pn:hypothetical protein]
CONTIG472	511665_c2_139	2869	8531	387	129	482	6.9(10)-46	Escherichia coli	b1451	[pn:hypothetical protein]
CONTIG472	11753260_c2_159	2870	8532	1452	484	1827	1.5(10)-188	Pseudomonas aeruginosa	Y10528	[PN:cyanide insensitive terminal oxidase] [GN:cioA] [DE:P.aeruginosa cioA and cioB genes.] [LE:276] [RE:1742] [DI:direct]
CONTIG472	24415937_c2_161	2871	8533	441	147	363	2.0(10)-33	Escherichia coli	b1379	[pn:beta-lactamase precursor] [gn:hsli]
CONTIG472	10807708_c3_171	2872	8534	474	158	647	1.6(10)-63	Escherichia coli	b1447	[pn:hypothetical protein]
CONTIG472	11955093_c3_192	2873	8535	3537	1179	5614	0	Escherichia coli	b1378	[pn:hypothetical protein] [gn:ydbk]
CONTIG472	29876562_c3_193	2874	8536	1155	385	1487	1.6(10)-152	Escherichia coli	b1377	[pn:hypothetical protein]
CONTIG473	20586437_fl_1	2875	8537	567	189	712	2.1(10)-70	Escherichia coli	b3033	[pn:hypothetical 16.5 kd protein in icc-tolc intergenic region] [gn:yqib]
CONTIG473	176443_fl_2	2876	8538	837	279	1158	1.2(10)-117	Escherichia coli	b3032	[pn:icc protein] [gn:icc]
CONTIG473	22775277_fl_3	2877	8539	1896	632	2898	4.7(10)-302	Escherichia coli	b3030	[pn:topoisomerase iv subunit] [gn:pare]
CONTIG473	10761003_fl_13	2878	8540	2277	759	3581	0	Escherichia coli	b3019	[pn:topoisomerase iv subunit] [gn:parc]
CONTIG473	5991325_fl_14	2879	8541	768	256	988	1.2(10)-99	Escherichia coli	b3018	[pn:l-acyl-glycerol-3-phosphate acyltransferase] [gn:plsc]
CONTIG473	4142253_fl_26	2880	8542	235	79	163	3.2(10)-12	Escherichia coli	b3005	[pn:biopolymer transport exbd protein] [gn:exbd]
CONTIG473	32223181_fl_27	2881	8543	702	234	965	3.2(10)-97	Escherichia coli	b3034	[pn:hypothetical protein] [gn:yqie]
CONTIG473	35678406_fl_33	2882	8544	1359	453	720	3.0(10)-71	Enterobacter cloacae	AB000622	or:enterobacter cloacae pn:mely gn:mely le:481 re:1758 di:direct sr:enterobacter cloacae (strain:iid977) dna
CONTIG473	4703276_fl_37	2883	8545	303	101	264	6.2(10)-23	Escherichia coli	I80320	hypothetical protein I - escherichia coli
CONTIG473	16924207_fl_38	2884	8546	417	139	347	1.0(10)-31	Escherichia coli	I80320	hypothetical protein I - escherichia coli

CONTIG473	4563840_f2_47	2885	8547	1539	513	2084	8.6(10)-216	Escherichia coli	b3017	[pn:sufi protein precursor] [gn:sufi]
CONTIG473	31924155_f2_54	2886	8548	618	206	121	1.6(10)-5	Escherichia coli	b3010	[pn:hypothetical protein]
CONTIG473	24648451_f2_55	2887	8549	906	302	1230	2.7(10)-125	Escherichia coli	b3010	[pn:hypothetical protein]
CONTIG473	32444783_f3_61	2888	8550	726	242	927	3.5(10)-93	Escherichia coli	b3031	[pn:hypothetical 15.2 kd protein in icc 3 <sup>'''</sup> region] [gn:yqia]
CONTIG473	29787518_f3_78	2889	8551	453	151	387	5.7(10)-36	Escherichia coli	b3024	[pn:hypothetical protein] [gn:ygiw]
CONTIG473	13070218_f3_83	2890	8552	2265	755	1968	1.7(10)-203	Escherichia coli	b3015	[pn:hypothetical protein]
CONTIG473	29776000_f3_89	2891	8553	753	251	960	1.1(10)-96	Escherichia coli	b3006	[pn:biopolymer transport exbb protein] [gn:exbb]
CONTIG473	189540_c1_91	2892	8554	1347	449	1771	1.3(10)-182	Escherichia coli	b3008	[pn:cystathionine beta-lyase] [gn:metc]
CONTIG473	11047306_c1_119	2893	8555	840	280	184	4.5(10)-12	Haloferax sp.	P21562	hypothetical 80.2 kd protein in the S' region of gyra and gyrb (orf 4).
CONTIG473	23489836_c1_122	2894	8556	378	126	191	5.0(10)-14	Haloferax sp.	P21562	hypothetical 80.2 kd protein in the S' region of gyra and gyrb (orf 4).
CONTIG473	9771052_c2_125	2895	8557	669	223	140	8.6(10)-10	Escherichia coli	M16489	or:escherichia coli le:627 re:1199 di:complement sr:escherichia coli dna nt:orf 4; putative
CONTIG473	29720016_c2_128	2896	8558	690	230	857	9.0(10)-86	Escherichia coli	b3009	[pn:hypothetical 24.1 kd protein in metc-sufi intergenic region] [gn:yghb]
CONTIG473	2931541_c2_145	2897	8559	1353	451	1330	6.9(10)-136	Escherichia coli	b3026	[pn:hypothetical protein] [gn:ygiy]
CONTIG473	35370336_c2_148	2898	8560	453	151	498	1.0(10)-47	Escherichia coli	b2665	[pn:hypothetical protein] [gn:yga]
CONTIG473	24220313_c2_154	2899	8561	900	300	175	1.7(10)-11	Bacillus subtilis	ydeC	[pn:hypothetical protein]
CONTIG473	16118916_c3_166	2900	8562	1242	414	1761	1.5(10)-181	Escherichia coli	b3011	[pn:hypothetical protein]
CONTIG473	36141433_c3_167	2901	8563	846	282	1175	1.8(10)-119	Escherichia coli	b3012	[pn:hypothetical protein]
CONTIG473	36110841_c3_178	2902	8564	678	226	918	3.1(10)-92	Escherichia coli	b3025	[pn:hypothetical protein] [gn:ygi]



CONTIG473	35288206_c3_180	2903	8565	699	233	959	1.3(10)-96	Escherichia coli	b3028	[pn:modulator of drug activity b] [gn:mdab]
CONTIG473	4698587_c3_181	2904	8566	327	109	459	1.3(10)-43	Escherichia coli	b3029	[pn:hypothetical 11.5 kd protein in mdab 3'''region] [gn:ygin]
CONTIG474	14957706_fl_7	2905	8567	1239	413	314	3.2(10)-28	Bacillus subtilis	ydeR	[pn:hypothetical protein]
CONTIG474	14179715_fl_22	2906	8568	714	238	644	3.3(10)-63	Escherichia coli	b1072	[pn:flagellar basal body p-ring protein flga precursor] [gn:flga]
CONTIG474	22792015_fl_23	2907	8569	336	112	358	6.9(10)-33	Escherichia coli	b1071	[pn:anti-sigma factor] [gn:flgm]
CONTIG474	25488888_fl_31	2908	8570	621	207	784	5.0(10)-78	Escherichia coli	b1063	[pn:hypothetical 20.5 kd protein in pyrc 3'''region] [gn:yceb]
CONTIG474	16495826_f2_33	2909	8571	360	120	189	5.5(10)-15	Haemophilus influenzae	U20229	ori:haemophilus influenzae pn:unknown le:4561 re:>4926 di:complement nt:orf121
CONTIG474	24650878_f2_38	2910	8572	3123	1041	2554	0	Escherichia coli	b1084	[pn:ribonuclease e] [gn:rne]
CONTIG474	22777150_f2_56	2911	8573	465	155	485	2.3(10)-46	Escherichia coli	b1070	[pn:flagella synthesis protein flgn] [gn:flgn]
CONTIG474	15729168_f2_65	2912	8574	1632	544	1518	8.1(10)-156	Escherichia coli	b1065	[pn:hypothetical protein]
CONTIG474	2923912_f2_66	2913	8575	357	119	320	7.2(10)-29	Escherichia coli	b1061	[pn:dna-damage-inducible protein ij] [gn:dini]
CONTIG474	22896012_f3_68	2914	8576	624	208	788	1.8(10)-78	Escherichia coli	b1087	[pn:hypothetical 23.2 kd protein in me-rpmf intergenic region] [gn:ycef]
CONTIG474	35634750_f3_99	2915	8577	1104	368	1539	4.9(10)-158	Escherichia coli	b1062	[pn:dihydroorotase] [gn:pyrc]
CONTIG474	2477266_c1_106	2916	8578	1068	356	1299	1.3(10)-132	Escherichia coli	b1068	[pn:virulence factor mvim homolog] [gn:mvim]
CONTIG474	31901507_c1_107	2917	8579	1620	540	2017	1.1(10)-208	Escherichia coli	b1069	[pn:virulence factor mvim homolog] [gn:mvim]
CONTIG474	26377035_c1_110	2918	8580	768	256	772	9.3(10)-77	Escherichia coli	b1075	[pn:flagellar hook formation protein flgd] [gn:flgd]
CONTIG474	35235306_c1_112	2919	8581	816	272	1108	2.2(10)-112	Escherichia coli	b1078	[pn:flagellar basal-body rod protein flgg] [gn:flgg]
CONTIG474	31772268_c1_113	2920	8582	756	252	1075	7.2(10)-109	Escherichia coli	b1079	[pn:flagellar l-ring protein precursor] [gn:flgh]
CONTIG474	23603381_c1_114	2921	8583	1101	367	1484	3.2(10)-152	Escherichia coli	b1080	[pn:flagellar p-ring protein precursor] [gn:flgi]

CONTIG474	25900202_c1_115	2922	8584	954	318	1165	2.1(10)-118	Escherichia coli	b1081	[pn:flagellar protein flg] [gn:flg]
CONTIG474	36022506_c1_116	2923	8585	1683	561	2059	3.8(10)-213	Escherichia coli	b1082	[pn:flagellar hook-associated protein 1] [gn:flgk]
CONTIG474	13862837_c1_127	2924	8586	195	65	135	2.8(10)-9	Escherichia coli	b1085	[pn:hypothetical protein]
CONTIG474	12369760_c1_131	2925	8587	534	178	757	3.6(10)-75	Escherichia coli	b1088	[pn:hypothetical 19.3 kd protein in me-rpmf intergenic region]
CONTIG474	22866711_c2_135	2926	8588	621	207	996	1.7(10)-100	Escherichia coli	b1066	[pn:ribosomal-protein-alanine acetyltransferase] [gn:rimj]
CONTIG474	5273568_c2_136	2927	8589	660	220	786	3.1(10)-78	Escherichia coli	b1067	[pn:hypothetical protein in rimj 3'" region] [gn:yceh]
CONTIG474	4312512_c2_142	2928	8590	441	147	526	1.1(10)-50	Escherichia coli	b1073	[pn:putative flagellar basal-body rod protein flgb] [gn:flgb]
CONTIG474	3129202_c2_143	2929	8591	405	135	643	4.2(10)-63	Escherichia coli	b1074	[pn:putative flagellar basal-body rod protein flgc] [gn:flgc]
CONTIG474	16289216_c2_147	2930	8592	768	256	1134	4.0(10)-115	Escherichia coli	b1077	[pn:putative flagellar basal-body rod protein flgf] [gn:flgf]
CONTIG474	4165937_c2_153	2931	8593	957	319	1123	5.9(10)-114	Escherichia coli	b1083	[pn:flagellar hook-associated protein 3] [gn:flg]
CONTIG474	35369016_c2_156	2932	8594	957	319	329	8.1(10)-30	Escherichia coli	b3060	[pn:hypothetical protein] [gn:ygip]
CONTIG474	11209818_c2_162	2933	8595	186	62	284	4.7(10)-25	Escherichia coli	b1089	[pn:50s ribosomal protein l32] [gn:rpmf]
CONTIG474	33645805_c2_163	2934	8596	321	107	218	7.0(10)-18	Escherichia coli	b1090	[pn:plsx protein] [gn:plsx]
CONTIG474	525278_c3_173	2935	8597	1212	404	1547	7.0(10)-159	Escherichia coli	b1076	[pn:flagellar hook protein flge] [gn:flge]
CONTIG474	34557338_c3_191	2936	8598	270	90	197	7.9(10)-16	Escherichia coli	b1085	[pn:hypothetical protein]
CONTIG474	26173166_c3_192	2937	8599	969	323	1383	1.7(10)-141	Escherichia coli	b1086	[pn:hypothetical 36.0 kd protein in me-rpmf intergenic region] [gn:yceec]
CONTIG475	21501261_fl_10	2938	8600	1218	406	114	0.00051	Escherichia coli	b1053	[pn:hypothetical 43.9 kd protein in msyb-htrb intergenic region] [gn:yceec]
CONTIG475	12538181_fl_32	2939	8601	1206	402	328	1.0(10)-29	Escherichia coli	b2214	[pn:hypothetical 38.5 kd protein in ada-omp intergenic region]
CONTIG475	5206911_fl_38	2940	8602	1014	338	450	1.2(10)-42	Escherichia coli	b2428	[pn:hypothetical protein] [gn:yfeu]

CONTIG475	6328406_f1_43	2941	8603	1506	502	1915	7.0(10)-198	Escherichia coli	b1961	[pn:dna-cytosine methyltransferase] [gn:dcn]
CONTIG475	21884378_f1_47	2942	8604	963	321	1088	3.0(10)-110	Escherichia coli	b1958	[pn:hypothetical protein in vsr 5' region] [gn:yedj]
CONTIG475	163125_f2_70	2943	8605	1008	336	164	5.5(10)-10	Escherichia coli	b0338	[pn:cyn operon transcriptional activator] [gn:cynr]
CONTIG475	29926551_f2_90	2944	8606	552	184	681	4.0(10)-67	Escherichia coli	b1960	[pn:patch repair protein] [gn:vst]
CONTIG475	19540756_f2_93	2945	8607	1182	394	959	1.3(10)-96	Escherichia coli	b1956	[pn:hypothetical protein]
CONTIG475	19610027_f3_110	2946	8608	942	314	492	4.4(10)-47	Escherichia coli	b0768	[pn:hypothetical transcriptional regulator in mode-bioa intergenic region] [gn:ybhd]
CONTIG475	1365625_f3_123	2947	8609	2805	935	706	9.1(10)-70	Bacillus subtilis	yqIG	[pn:hypothetical protein]
CONTIG475	782525_f3_128	2948	8610	771	257	712	2.1(10)-70	Escherichia coli	b1962	[pn:hypothetical protein in seru-dcm intergenic region] [gn:yedj]
CONTIG475	14064562_f3_135	2949	8611	213	71	205	1.1(10)-16	Escherichia coli	b1957	[pn:hypothetical protein]
CONTIG475	16207680_f3_136	2950	8612	258	86	193	1.8(10)-14	Escherichia coli	b1956	[pn:hypothetical protein]
CONTIG475	11197708_c1_146	2951	8613	519	173	267	3.0(10)-23	Escherichia coli	P09183	very hypothetical 20.3 kd protein in dcm 3' region (orf 3).
CONTIG475	6453252_c1_147	2952	8614	648	216	132	6.0(10)-9	Escherichia coli	P09183	very hypothetical 20.3 kd protein in dcm 3' region (orf 3).
CONTIG475	4070467_c1_148	2953	8615	1179	393	1096	4.2(10)-111	Escherichia coli	b1377	[pn:hypothetical protein]
CONTIG475	15656_c1_151	2954	8616	1644	548	1602	1.0(10)-164	Escherichia coli	b4125	[pn:hypothetical 60.6 kd protein in dcub-lysu intergenic region]
CONTIG475	25437753_c1_166	2955	8617	1455	485	1821	6.4(10)-188	Klebsiella pneumoniae	P16482	citrate-proton symport (citrate transporter) (citrate carrier protein).
CONTIG475	3159530_c1_167	2956	8618	795	265	107	0.0011	Bacillus subtilis	yvgL	[pn:hypothetical protein] [gn:yvsd]
CONTIG475	175006_c1_169	2957	8619	837	279	1212	2.2(10)-123	Escherichia coli	b1976	[pn:hypothetical protein]
CONTIG475	4022768_c1_173	2958	8620	1752	584	1683	2.7(10)-173	Escherichia coli	S44018	iuca protein - escherichia coli

CONTIG475	9766281_c1_176	2959	8621	705	235	730	2.6(10)-72	Escherichia coli	AF016586	[de:escherichia coli plasmid pcolv-k311 lysine n6-hydroxylase (aera)gene, complete cds.] [pn:lysine n6-hydroxylase] [gn:aera] [nt:monooxygenase]
CONTIG475	36525756_c1_177	2960	8622	555	185	462	6.5(10)-44	Escherichia coli	AF016587	[de:escherichia coli plasmid pcolv-k311 lysine n6-hydroxylase mutant(aera) gene, complete cds.] [pn:lysine n6-hydroxylase mutant] [gn:aera] [nt:monooxygenase; p14g site-directed mutant]
CONTIG475	32454093_c2_184	2961	8623	921	307	1095	5.5(10)-111	Escherichia coli	b1959	[pn:hypothetical 32.2 kd protein in vsr 5''' region] [gn:yeda]
CONTIG475	4939042_c2_197	2962	8624	1569	523	928	2.7(10)-93	Escherichia coli	b3063	[pn:hypothetical 52.9 kd protein in ttdb-rpsu intergenic region] [gn:ygie]
CONTIG475	2398507_c2_201	2963	8625	1329	443	363	2.0(10)-33	Plasmid pNAD2	D10678	or:plasmid pna2 pn:6-aminohexanoate-dimer hydrolase gn:nylb ec:3.5.1.46 le:611 re:1801 di:direct sr:plasmid pna2 dna
CONTIG475	35729157_c2_208	2964	8626	1908	636	1971	8.1(10)-204	Escherichia coli	S50883	iucc protein - escherichia coli
CONTIG475	16828512_c3_216	2965	8627	282	94	94	7.7(10)-5	Drosophila melanogaster	P13238	vitelline membrane protein vm26ab precursor (protien tu-4) (protein sv23).
CONTIG475	20526691_c3_219	2966	8628	735	245	882	2.0(10)-88	Escherichia coli	b4124	[pn:hypothetical 27.4 kd protein in dcub-lysu intergenic region]
CONTIG475	4510206_c3_223	2967	8629	1695	565	2482	5.7(10)-258	Escherichia coli	b4122	[pn:fumarate hydratase class i, anaerobic] [gn:fumb]
CONTIG475	6539193_c3_237	2968	8630	960	320	1100	1.6(10)-111	Escherichia coli	S44019	iucb protein - escherichia coli
CONTIG475	12314388_c3_240	2969	8631	459	153	91	0.023	Escherichia coli	AF016587	[de:escherichia coli plasmid pcolv-k311 lysine n6-hydroxylase mutant(aera) gene, complete cds.] [pn:lysine n6-hydroxylase mutant] [gn:aera] [nt:monooxygenase; p14g site-directed mutant]
CONTIG475	23625277_c3_241	2970	8632	1221	407	1254	7.7(10)-128	Escherichia coli	S01042	cloacin receptor precursor - escherichia coli plasmidpcolv-k30

CONTIG476	3400177_f1_1	2971	8633	714	238	168	9.4(10)-13	Escherichia coli	b0375	[pn:hypothetical protein] [gn:yaiv]
CONTIG476	3912575_f1_18	2972	8634	207	69	140	8.6(10)-10	Escherichia coli	b1705	[pn:hypothetical 7.1 kd protein in aroh-nlpc intergenic region]
CONTIG476	16538217_f1_20	2973	8635	1197	399	1487	1.6(10)-152	Shigella dysenteriae	U64516	[de:shigella dysenteriae shuv, shuu, shuy, shux, shuw pseudogene, shut, outer membrane heme receptor shua (shua), and shus genes, completecds.] [pn:unknown] [gn:shus]
CONTIG476	11110830_f2_37	2974	8636	684	228	182	3.1(10)-14	Escherichia coli	b0375	[pn:hypothetical protein] [gn:yaiv]
CONTIG476	12689051_f2_60	2975	8637	2067	689	2705	1.3(10)-281	Shigella dysenteriae	U64516	or:shigella dysenteriae pn:outer membrane heme receptor shua gn:shua le:491 re:2473 di:direct
CONTIG476	15724136_f2_64	2976	8638	1071	357	722	1.8(10)-71	Yersinia enterocolitica	S54438	hemin permease - yersinia enterocolitica
CONTIG476	7157942_f3_91	2977	8639	1176	392	1187	9.8(10)-121	Escherichia coli	b1688	[pn:hypothetical protein]
CONTIG476	581252_f3_99	2978	8640	1059	353	1279	1.7(10)-130	Escherichia coli	b1703	[pn:hypothetical 19.8 kd protein in ppsa-aroh intergenic region] [gn:ydia]
CONTIG476	976542_f3_100	2979	8641	1071	357	1553	1.6(10)-159	Escherichia coli	b1704	[pn:3-deoxy-d-arabinoheptulosonate 7-phosphate synthase]
CONTIG476	16219541_f3_108	2980	8642	846	282	741	1.8(10)-73	Yersinia enterocolitica	X77867	or:yersinia enterocolitica pn:hemin binding protein gn:hemt le:1175 re:2014 di:direct
CONTIG476	35397915_f3_110	2981	8643	972	324	370	3.7(10)-34	Escherichia coli	b0151	[pn:ferrichrome transport ap-binding protein fhuc] [gn:fhuc]
CONTIG476	30331908_c1_121	2982	8644	1071	357	1601	1.3(10)-164	Escherichia coli	b1714	[pn:phenylalanyl-trna synthetase alpha chain] [gn:phe5]
CONTIG476	13164051_c1_123	2983	8645	1074	358	1188	7.7(10)-121	Escherichia coli	b1711	[pn:vitamin b12 transport system permease protein btuc] [gn:btuc]
CONTIG476	4032843_c1_124	2984	8646	561	187	738	3.7(10)-73	Escherichia coli	b1710	[pn:vitamin b12 transport periplasmic protein btue] [gn:btue]
CONTIG476	21752052_c1_138	2985	8647	2433	811	3905	0	Escherichia coli	b1702	[pn:phosphoenolpyruvate synthase] [gn:ppsa]
CONTIG476	2618785_c1_139	2986	8648	1485	495	1219	4.0(10)-124	Escherichia coli	b0270	[pn:hypothetical 50.6 kd protein in per-arg intergenic region]

CONTIG476	15744032_c1_140	2987	8649	2379	793	992	4.5(10)-100	Escherichia coli	b2132	[pn:periplasmic beta-glucosidase precursor] [gn:bglx]
CONTIG476	35234831_c2_151	2988	8650	2280	760	3516	0	Escherichia coli	b1713	[pn:phenylalanyl-trna synthetase beta chain] [gn:phet]
CONTIG476	34105290_c2_155	2989	8651	729	243	608	2.2(10)-59	Escherichia coli	b1707	[pn:hypothetical protein]
CONTIG476	12986466_c2_182	2990	8652	453	151	532	2.5(10)-51	Escherichia coli	b1686	[pn:hypothetical protein]
CONTIG476	12703468_c3_188	2991	8653	312	104	491	5.5(10)-47	Escherichia coli	b1712	[pn:integration host factor alpha-subunit] [gn:hima]
CONTIG476	35676636_c3_190	2992	8654	813	271	812	5.4(10)-81	Escherichia coli	b1709	[pn:vitamin b12 transport atp-binding protein btud] [gn:btud]
CONTIG476	34191502_c3_191	2993	8655	504	168	653	3.7(10)-64	Escherichia coli	b1708	[pn:probable lipoprotein nlpc precursor] [gn:nlpc]
CONTIG476	38136_c3_193	2994	8656	1584	528	1970	1.0(10)-203	Escherichia coli	b1706	[pn:hypothetical protein]
CONTIG476	6054688_c3_213	2995	8657	3129	1043	4774	0	Escherichia coli	b1687	[pn:hypothetical protein] [gn:ydi]
CONTIG476	21766510_c3_215	2996	8658	231	77	238	3.6(10)-20	Escherichia coli	b1685	[pn:hypothetical protein]
CONTIG476	36439017_c3_216	2997	8659	546	182	243	1.1(10)-20	Escherichia coli	b4314	[pn:type 1 fimbrial subunit] [gn:fima]
CONTIG477	35187530_fl_3	2998	8660	1884	628	2268	2.7(10)-235	Escherichia coli	b1222	[pn:nitrate/nitrite sensor protein narx] [gn:narx]
CONTIG477	15817592_fl_23	2999	8661	678	226	537	7.4(10)-52	Escherichia coli	b1219	[pn:hypothetical 12.7 kd protein in chac-nar1 intergenic region] [gn:ychn]
CONTIG477	12632675_fl_27	3000	8662	1131	377	1379	4.4(10)-141	Escherichia coli	b1216	[pn:putative calcium/proton antiporter] [gn:chaa]
CONTIG477	21994077_fl_33	3001	8663	870	290	1241	1.8(10)-126	Escherichia coli	b1208	[pn:hypothetical 30.9 kd protein in hemm-prs intergenic region] [gn:yehb]
CONTIG477	24656552_fl_34	3002	8664	1686	562	2131	9.0(10)-221	Escherichia coli	b1206	[pn:hypothetical protein in pth-prs intergenic region] [gn:yehm]
CONTIG477	26753326_f2_41	3003	8665	669	223	865	1.3(10)-86	Escherichia coli	b1221	[pn:nitrate/nitrite response regulator protein narl] [gn:narl]
CONTIG477	11988405_f2_47	3004	8666	246	82	102	0.00012	Murine herpesvirus 68	U97553	[de:murine herpesvirus 68 strain wums, complete genome.] [pn:unknown] [gn:gammahv.m6]

CONTIG477	23625001_f2_62	3005	8667	1899	633	565	8.0(10)-55	Escherichia coli	b4355	[pn:methyl-accepting chemotaxis protein i] [gn:tsr]
CONTIG477	12634832_f2_78	3006	8668	522	174	638	1.5(10)-62	Escherichia coli	b1209	[pn:hemm protein] [gn:hemm]
CONTIG477	5197277_f2_81	3007	8669	588	196	818	1.2(10)-81	Escherichia coli	b1204	[pn:peptidyl-tRNA hydrolase] [gn:pth]
CONTIG477	26173518_f2_82	3008	8670	1107	369	1584	8.4(10)-163	Escherichia coli	b1203	[pn:hypothetical gtp-binding protein in pth 3''' region] [gn:ychf]
CONTIG477	33228141_f3_131	3009	8671	1017	339	1563	1.3(10)-160	Escherichia coli	b1207	[pn:ribose-phosphate pyrophosphokinase] [gn:prsa]
CONTIG477	32228386_c1_148	3010	8672	279	93	433	7.7(10)-41	Escherichia coli	b1205	[pn:hypothetical 10.5 kd protein in pth-prs intergenic region] [gn:ychh]
CONTIG477	23694406_c1_149	3011	8673	1116	372	507	1.1(10)-48	Klebsiella oxytoca	P10488	albicidin resistance protein.
CONTIG477	15876283_c1_159	3012	8674	888	296	972	5.9(10)-98	Escherichia coli	b1212	[pn:possibly protoporphyrinogen oxidase] [gn:hemk]
CONTIG477	32690636_c1_163	3013	8675	843	281	1020	4.9(10)-103	Escherichia coli	b1218	[pn:cation transport protein chac] [gn:chac]
CONTIG477	22062836_c1_170	3014	8676	1365	455	1689	6.2(10)-174	Klebsiella pneumoniae	L27431	[de:klebsiella pneumoniae nitrate transporter component (nasf), nitratetransporter component (nase), nitrate transporter atpase component(nasd), and"] [pn:nitrate transporter component] [gn:nasf]
CONTIG477	6047667_c1_186	3015	8677	1440	480	1753	1.0(10)-180	Escherichia coli	b1223	[pn:nitrite extrusion protein] [gn:nark]
CONTIG477	24032082_c2_203	3016	8678	1164	388	1698	6.9(10)-175	Escherichia coli	b1211	[pn:peptide chain release factor 1] [gn:prfa]
CONTIG477	34585931_c2_209	3017	8679	249	83	251	1.5(10)-21	Escherichia coli	b1217	[pn:cation transport regulator chab] [gn:chab]
CONTIG477	24724187_c2_212	3018	8680	1227	409	900	2.5(10)-90	Klebsiella pneumoniae	A55859	regulatory protein nasr - klebsiella pneumoniae
CONTIG477	25672942_c2_226	3019	8681	2505	835	732	1.6(10)-72	Methanobacterium thermoautotrophicum	MTH1552	[pn:formate dehydrogenase, alpha subunit homolog]
CONTIG477	1291575_c2_227	3020	8682	1419	473	1532	2.7(10)-157	Escherichia coli	b1220	[pn:hypothetical protein in narl 5''' region] [gn:ychp]

CONTIG477	3144165_c3_243	3021	8683	1302	434	1913	1.1(10)-197	Escherichia coli	b1210	[pn:glutamyl-tRNA reductase] [gn:hema]
CONTIG477	16219457_c3_248	3022	8684	417	139	395	8.3(10)-37	Escherichia coli	b1213	[pn:hypothetical protein]
CONTIG477	11212778_c3_249	3023	8685	813	271	1139	1.2(10)-115	Escherichia coli	b1214	[pn:hypothetical protein in hemk-kdsa] [gn:ycha]
CONTIG477	4001668_c3_250	3024	8686	864	288	1362	2.7(10)-139	Escherichia coli	b1215	[pn:2-dehydro-3-deoxyphosphooctonate aldolase] [gn:kdsa]
CONTIG477	34396092_c3_265	3025	8687	891	297	895	8.5(10)-90	Klebsiella pneumoniae	L27431	[de:klebsiella pneumoniae nitrate transporter component (nasf), nitratetransporter component (nase), nitrate transporter atpase component(nasd), and nitrate reductase small"] [pn:nitrate transporter component] [gn:nase]
CONTIG477	5275443_c3_266	3026	8688	798	266	1270	1.6(10)-129	Klebsiella pneumoniae	L27431	[de:klebsiella pneumoniae nitrate transporter component (nasf), nitratetransporter component (nase), nitrate transporter atpase component(nasd)] [pn:nitrate transporter atpase component] [gn:nasd]
CONTIG477	5192968_c3_267	3027	8689	3996	1332	2499	9.1(10)-260	Escherichia coli	b3365	[pn:nadh-nitrate oxidoreductase apoprotein] [gn:nirb]
CONTIG478	32453438_fl_1	3028	8690	582	194	111	7.2(10)-5	Bacillus subtilis	pksA	[pn:transcriptional regulator]
CONTIG478	12548567_fl_15	3029	8691	918	306	878	5.4(10)-88	Escherichia coli	b1466	[pn:respiratory nitrate reductase 2 delta chain] [gn:narw]
CONTIG478	12213952_fl_18	3030	8692	765	255	854	1.8(10)-85	Escherichia coli	b1462	[pn:hypothetical protein]
CONTIG478	6735393_fl_19	3031	8693	285	95	446	3.2(10)-42	Escherichia coli	b4294	[pn:insertion element islf protein insa] [gn:insa_7]
CONTIG478	2242625_f2_52	3032	8694	219	73	251	1.5(10)-21	Escherichia coli	b2215	[pn:outer membrane protein c precursor] [gn:ompc]
CONTIG478	4725761_f2_57	3033	8695	1473	491	1795	3.7(10)-185	Escherichia coli	b1469	[pn:nitrite extrusion protein] [gn:naru]
CONTIG478	32313576_f2_62	3034	8696	1590	530	2635	3.5(10)-274	Escherichia coli	b1467	[pn:respiratory nitrate reductase 2 beta chain] [gn:nary]



CONTIG478	2911290_f2_67	3035	8697	234	78	296	2.6(10)-26	Escherichia coli	S40546	hypothetical protein - escherichia coli
CONTIG478	25437756_f2_71	3036	8698	345	115	237	4.5(10)-20	Bacillus subtilis	yezG	[pn:hypothetical protein]
CONTIG478	1207283_f2_89	3037	8699	897	299	693	2.2(10)-68	Escherichia coli	b1328	[pn:hypothetical protein] [gn:ycjz]
CONTIG478	22929826_f2_90	3038	8700	1599	533	2129	1.5(10)-220	Escherichia coli	b1453	[pn:hypothetical protein] [gn:ansp]
CONTIG478	10312956_f2_93	3039	8701	996	332	1314	3.3(10)-134	Escherichia coli	b1451	[pn:hypothetical protein]
CONTIG478	2456415_f3_100	3040	8702	3783	1261	6042	0	Escherichia coli	b1468	[pn:respiratory nitrate reductase 2 alpha chain] [gn:narz]
CONTIG478	31508556_f3_105	3041	8703	729	243	885	9.9(10)-89	Escherichia coli	b1465	[pn:respiratory nitrate reductase 2 gamma chain] [gn:narv]
CONTIG478	25567885_f3_108	3042	8704	525	175	898	4.0(10)-90	Escherichia coli	b0264	[pn:insertion element isl protein insb] [gn:insb_2]
CONTIG478	21500925_f3_110	3043	8705	537	179	237	3.5(10)-19	Eikenella corrodens	P35649	hypothetical 66.3 kd protein in hag2 5'region.
CONTIG478	16987716_f3_111	3044	8706	270	90	185	1.5(10)-13	Eikenella corrodens	P35649	hypothetical 66.3 kd protein in hag2 5'region.
CONTIG478	21970417_f3_112	3045	8707	954	318	479	1.0(10)-45	Eikenella corrodens	P35649	hypothetical 66.3 kd protein in hag2 5'region.
CONTIG478	4884652_c1_148	3046	8708	402	134	102	9.3(10)-6	Enterobacter aerogenes	U67194	or:enterobacter aerogenes pn:pep l gn:tnpa le:11649 re:12071 di:complement nt:orf1
CONTIG478	26429653_c1_151	3047	8709	1020	340	539	4.5(10)-52	Bacillus subtilis	ymnO	[pn:hypothetical protein]
CONTIG478	24667250_c1_154	3048	8710	1329	443	1352	3.2(10)-138	Bacillus subtilis	ymnJ	[pn:hypothetical protein]
CONTIG478	35664657_c1_158	3049	8711	837	279	559	3.5(10)-54	Escherichia coli	b1917	[pn:hypothetical protein] [gn:yecc]
CONTIG478	5260443_c1_159	3050	8712	897	299	165	1.2(10)-10	Escherichia coli	b1920	[pn:fly protein precursor] [gn:fly]
CONTIG478	6439130_c1_160	3051	8713	273	91	244	8.3(10)-21	Escherichia coli	b1461	[pn:hypothetical 8.7 kd protein in rhse-narv intergenic region] [gn:ydee]
CONTIG478	189212_c1_161	3052	8714	1017	339	1053	1.6(10)-106	Escherichia coli	b1054	[pn:membrane protein affecting cell division, growth and high temperature survival] [gn:htrb]

CONTIG478	33787750_c1_169	3053	8715	882	294	1153	3.8(10)-117	Escherichia coli	b1463	[pn:hypothetical protein]
CONTIG478	4329693_c2_205	3054	8716	2112	704	2500	7.2(10)-260	Escherichia coli	b3340	[pn:fusa] [gn:fusa]
CONTIG478	5899187_c3_232	3055	8717	1074	358	1425	5.9(10)-146	Escherichia coli	b1452	[pn:hypothetical protein]
CONTIG478	1267275_c3_237	3056	8718	576	192	97	0.00259	Bacillus subtilis	yqaC	[pn:hypothetical protein]
CONTIG478	19821086_c3_239	3057	8719	1137	379	879	4.2(10)-88	Bacillus subtilis	yxep	[pn:hypothetical protein] [gn:lp9h]
CONTIG478	15908556_c3_241	3058	8720	1062	354	206	1.7(10)-16	Escherichia coli	b1920	[pn:fly protein precursor] [gn:fly]
CONTIG478	4036693_c3_242	3059	8721	519	173	134	3.7(10)-9	Methanobacterium thermoautotrophicum	MTH676	[pn:unknown]
CONTIG478	5197943_c3_243	3060	8722	936	312	215	9.8(10)-18	Escherichia coli	b1918	[pn:yecc]
CONTIG478	4489843_c3_244	3061	8723	627	209	851	3.8(10)-85	Escherichia coli	b1454	[pn:hypothetical protein]
CONTIG478	14885165_c3_249	3062	8724	297	99	390	2.7(10)-36	Escherichia coli	D93826	hypothetical 11k protein (insertion sequence is 1) -escherichia coli this protein is coded by the insertion sequence is 1.
CONTIG478	1290917_c3_265	3063	8725	1395	465	1230	2.7(10)-125	Salmonella typhimurium	P37594	methyl viologen resistance protein smva.
CONTIG478	26175336_c3_266	3064	8726	216	72	137	1.8(10)-9	Bacteriophage PA2	E25647	hypothetical lc protein - phage pa2
CONTIG479	34552086_fl_3	3065	8727	786	262	1064	1.1(10)-107	Escherichia coli	b1868	[pn:hypothetical protein in asps 5' region] [gn:yecc]
CONTIG479	4423942_fl_4	3066	8728	429	143	581	1.6(10)-56	Escherichia coli	b1869	[pn:hypothetical protein]
CONTIG479	10167501_fl_5	3067	8729	1011	337	1481	6.9(10)-152	Escherichia coli	b1871	[pn:hypothetical protein]
CONTIG479	7206518_fl_8	3068	8730	1743	581	2582	1.5(10)-268	Escherichia coli	b1876	[pn:arginyl-tRNA synthetase] [gn:args]
CONTIG479	16048255_f2_45	3069	8731	1347	449	1223	1.5(10)-124	Escherichia coli	b1579	[pn:hypothetical protein]

CONTIG479	3364465_f2_47	3070	8732	840	280	1190	4.7(10)-121	Escherichia coli	b1870	[pn:hypothetical protein]
CONTIG479	39052_f2_76	3071	8733	1317	439	646	2.1(10)-63	Escherichia coli	b4077	[pn:glutamate- aspartate carrier] [gn:gltp]
CONTIG479	6104837_f3_140	3072	8734	441	147	516	1.2(10)-49	Escherichia coli	b1895	[pn:hypothetical 17.1 kd protein in flhD-otsA intergenic region]
CONTIG479	24645818_c1_144	3073	8735	936	312	1293	5.7(10)-132	Escherichia coli	b1889	[pn:chemotaxis motB protein] [gn:motb]
CONTIG479	35330006_c1_149	3074	8736	792	264	103	0.0023	Escherichia coli	b0944	[pn:hypothetical fimbrial chaperone in pepN-pyrd intergenic region] [gn:ycbf]
CONTIG479	13080431_c1_161	3075	8737	465	155	371	2.8(10)-34	Escherichia coli	b1882	[pn:chemotaxis protein cheY] [gn:cheY]
CONTIG479	35188562_c2_181	3076	8738	1431	477	1957	2.5(10)-202	Escherichia coli	b1896	[pn:alpha trehalose phosphate synthase] [gn:otsA]
CONTIG479	26661516_c2_183	3077	8739	606	202	934	6.2(10)-94	Escherichia coli	b1891	[pn:flagellar transcriptional activator] [gn:flhC]
CONTIG479	25604716_c2_184	3078	8740	903	301	1159	9.0(10)-118	Escherichia coli	b1890	[pn:chemotaxis motA protein] [gn:motA]
CONTIG479	36349037_c2_186	3079	8741	2076	692	2572	1.7(10)-267	Escherichia coli	b1888	[pn:chemotaxis protein cheA] [gn:cheA]
CONTIG479	24492202_c2_187	3080	8742	573	191	102	0.0004	Myxococcus xanthus	P27755	protein u precursor.
CONTIG479	15760407_c2_197	3081	8743	1635	545	1629	1.3(10)-167	Escherichia coli	b1885	[pn:methyl-accepting chemotaxis protein iv] [gn:tap]
CONTIG479	4570318_c2_199	3082	8744	1059	353	1653	4.0(10)-170	Escherichia coli	b1883	[pn:protein-glutamate methyltransferase] [gn:cheB]
CONTIG479	12579542_c2_200	3083	8745	720	240	795	3.3(10)-79	Escherichia coli	b1881	[pn:chemotaxis protein cheZ] [gn:cheZ]
CONTIG479	3407752_c2_201	3084	8746	1173	391	1419	2.6(10)-145	Escherichia coli	b1880	[pn:hypothetical protein] [gn:flhB]
CONTIG479	9929581_c2_205	3085	8747	432	144	472	5.7(10)-45	Escherichia coli	b1878	[pn:hypothetical protein] [gn:flhE]
CONTIG479	14855425_c2_206	3086	8748	1599	533	197	1.2(10)-12	Escherichia coli	b3657	[pn:hypothetical 51.0 kd protein in glts-selC intergenic region] [gn:yicJ]
CONTIG479	24744006_c2_212	3087	8749	564	188	860	4.4(10)-86	Escherichia coli	b1875	[pn:hypothetical protein] [gn:yecM]
CONTIG479	23445328_c2_213	3088	8750	765	255	478	1.3(10)-45	Escherichia coli	b1874	[pn:copper homeostasis protein] [gn:cutC]

CONTIG479	30101377_c3_221	3089	8751	462	154	503	3.0(10)-48	Escherichia coli	b1892	[pn:flagellar transcriptional activator flhD] [gn:flhD]
CONTIG479	23726553_c3_224	3090	8752	519	173	693	2.2(10)-68	Escherichia coli	b1887	[pn:purine binding chemotaxis protein] [gn:chew]
CONTIG479	31806341_c3_227	3091	8753	609	203	1119	3.1(10)-6	Myxococcus xanthus	P27755	protein u precursor.
CONTIG479	20604540_c3_229	3092	8754	2400	800	169	7.2(10)-9	Escherichia coli	b3144	[pn:hypothetical outer membrane usher protein in agal-mtr intergenic region] [gn:yraj]
CONTIG479	11056950_c3_230	3093	8755	1773	591	1501	5.2(10)-154	Escherichia coli	b1886	[pn:methyl-accepting chemotaxis protein ii] [gn:tar]
CONTIG479	4036688_c3_232	3094	8756	876	292	1309	1.2(10)-133	Escherichia coli	b1884	[pn:chemotaxis protein methyltransferase] [gn:cher]
CONTIG479	15117832_c3_237	3095	8757	2094	698	2605	5.4(10)-271	Escherichia coli	b1879	[pn:flagellar biosynthesis protein flhA] [gn:flhA]
CONTIG479	13852002_c3_239	3096	8758	1167	389	422	1.1(10)-39	Bacillus subtilis	yesR	[pn:hypothetical protein]
CONTIG48	6023461_c1_7	3097	8759	333	111	110	1.3(10)-6	Bacteriophage p1	L01408	or:bacteriophage p1 pn:partition protein gn:parb le:<1 re:456 di:direct sr:bacteriophage p1 (individual isolate p1kc) dna
CONTIG480	26740760_f1_13	3098	8760	621	207	631	8.0(10)-62	Helicobacter pylori	HP1563	[pn:alkyl hydroperoxide reductase] [gn:tsaa]
CONTIG480	24477316_f1_35	3099	8761	3147	1049	2369	5.5(10)-246	Escherichia coli	b0397	[pn:exonuclease sbcc] [gn:sbcc]
CONTIG480	24035875_f1_39	3100	8762	942	314	1343	2.8(10)-137	Escherichia coli	b0393	[pn:hypothetical 34.0 kd protein in arom-araj intergenic region]
CONTIG480	6461591_f1_45	3101	8763	822	274	1206	9.5(10)-123	Escherichia coli	b0386	[pn:pyrroline-5-carboxylate reductase] [gn:proc]
CONTIG480	4580282_f2_49	3102	8764	900	300	1472	6.2(10)-151	Escherichia coli	b0411	[pn:nucleoside-specific channel-forming protein tsx precursor] [gn:tsx]
CONTIG480	25547211_f2_80	3103	8765	1323	441	1708	6.0(10)-176	Escherichia coli	b0398	[pn:exonuclease sbcd] [gn:sbcd]
CONTIG480	21611505_f3_104	3104	8766	639	213	798	1.6(10)-79	Escherichia coli	b0404	[pn:hypothetical 23.0 kd protein in malz-quea intergenic region] [gn:yajb]
CONTIG480	14329693_c1_138	3105	8767	273	91	306	2.2(10)-27	Escherichia coli	b0382	[pn:hypothetical protein fragment in ddla-phoa intergenic region]

CONTIG480	14556508_c1_141	3106	8768	195	65	284	4.7(10)-25	Escherichia coli	b0389	[pn:hypothetical 7.3 kd protein in aro-arom intergenic region]
CONTIG480	21875076_c1_145	3107	8769	1107	369	1330	6.9(10)-136	Escherichia coli	b0394	[pn:hypothetical protein in aro-arom intergenic region] [gn:yajf]
CONTIG480	25945251_c1_157	3108	8770	717	239	1153	3.8(10)-117	Escherichia coli	b0399	[pn:phosphate regulon transcriptional regulatory protein]
CONTIG480	26587916_c1_158	3109	8771	1299	433	1920	2.1(10)-198	Escherichia coli	b0400	[pn:phosphate regulon sensor protein phor] [gn:phor]
CONTIG480	14875018_c1_159	3110	8772	1326	442	1898	4.4(10)-196	Escherichia coli	b0401	[pn:branched chain amino acid transport system ii carrier protein] [gn:brnq]
CONTIG480	14588430_c1_160	3111	8773	1455	485	1858	7.7(10)-192	Escherichia coli	b0402	[pn:hypothetical protein] [gn:proy]
CONTIG480	21492016_c1_161	3112	8774	1845	615	2711	3.1(10)-282	Escherichia coli	b0403	[pn:maltodextrin glucosidase] [gn:malz]
CONTIG480	6536091_c1_165	3113	8775	474	158	97	0.0008	Nephila clavipes	AF027735	[de:nephila clavipes minor ampullate silk protein misp1 mrna, partial cds.] [pn:minor ampullate silk protein misp1]
CONTIG480	12994007_c1_172	3114	8776	1107	369	1748	3.5(10)-180	Escherichia coli	b0405	[pn:trna ribosyltransferase-isomerase] [gn:quea]
CONTIG480	5327_c1_175	3115	8777	333	111	531	3.2(10)-51	Escherichia coli	b0407	[pn:hypothetical 11.9 kd protein in tgt-secd intergenic region] [gn:yajc]
CONTIG480	23280_c1_176	3116	8778	1875	625	2610	1.6(10)-271	Escherichia coli	b0408	[pn:protein-export membrane protein secd] [gn:secd]
CONTIG480	36073311_c1_178	3117	8779	384	128	130	2.3(10)-8	Mycobacterium tuberculosis	Z95210	unknown,,mtcy21c12.05. len
CONTIG480	15055465_c2_187	3118	8780	471	157	613	6.5(10)-60	Escherichia coli	b0387	[pn:yaii]
CONTIG480	35629031_c2_188	3119	8781	669	223	569	3.0(10)-55	Escherichia coli	b0388	[pn:shikimate kinase ii] [gn:aro]
CONTIG480	1953465_c2_189	3120	8782	321	107	435	4.7(10)-41	Escherichia coli	b0391	[pn:hypothetical 10.2 kd protein in aro-araj intergenic region]
CONTIG480	13911282_c2_215	3121	8783	1026	342	1414	8.5(10)-145	Escherichia coli	b0409	[pn:protein-export membrane protein secf] [gn:secf]
CONTIG480	12501906_c2_217	3122	8784	744	248	218	4.7(10)-18	Bacillus subtilis	yobV	[pn:hypothetical protein]

CONTIG480	35251317_c3_219	3123	8785	591	197	549	4.0(10)-53	Escherichia coli	b1053	[pn:hypothetical 43.9 kd protein in msyb-htrb intergenic region] [gn:ycee]
CONTIG480	3004066_c3_220	3124	8786	363	121	447	2.6(10)-42	Escherichia coli	b0384	[pn:phosphate starvation-inducible protein psif] [gn:psif]
CONTIG480	21759841_c3_221	3125	8787	1200	400	996	1.7(10)-100	Escherichia coli	b0385	[pn:hypothetical 41.5 kd protein in psif-proc intergenic region] [gn:yaic]
CONTIG480	5214091_c3_225	3126	8788	693	231	799	1.3(10)-79	Escherichia coli	b0390	[pn:arom protein] [gn:arom]
CONTIG480	36580083_c3_251	3127	8789	1197	399	1968	1.7(10)-203	Escherichia coli	b0406	[pn:queueine trna-ribosyltransferase] [gn:tgt]
CONTIG481	34042063_fl_6	3128	8790	1419	473	2247	4.5(10)-233	Escherichia coli	b3962	[pn:unknown dehydrogenase a] [gn:udha]
CONTIG481	14541018_fl_13	3129	8791	1164	388	1789	1.6(10)-184	Escherichia coli	b3957	[pn:acetylornithine deacetylase] [gn:arge]
CONTIG481	9806318_fl_26	3130	8792	2523	841	2841	5.2(10)-296	Escherichia coli	b3947	[pn:phosphoenolpyruvate-protein phosphotransferase ptsa] [gn:ptsa]
CONTIG481	29892965_f2_59	3131	8793	2670	890	4102	0	Escherichia coli	b3956	[pn:phosphoenolpyruvate carboxylase] [gn:ppc]
CONTIG481	27137_f2_60	3132	8794	912	304	1168	1.0(10)-118	Escherichia coli	b3954	[pn:hypothetical transcriptional regulator in glda- ppc intergenic] [gn:yjio]
CONTIG481	4428176_f3_85	3133	8795	1179	393	1788	2.0(10)-184	Escherichia coli	b3965	[pn:uracil-5-methyltransferase] [gn:trma]
CONTIG481	33620166_f3_116	3134	8796	732	244	900	2.5(10)-90	Escherichia coli	b3946	[pn:transaldolase-like protein] [gn:talc]
CONTIG481	5977218_f3_117	3135	8797	1116	372	1734	1.1(10)-178	Escherichia coli	b3945	[pn:glycerol dehydrogenase] [gn:glga]
CONTIG481	4103427_c1_149	3136	8798	921	307	1508	9.5(10)-155	Escherichia coli	b3961	[pn:hydrogen peroxide-inducible genes activator] [gn:oxyr]
CONTIG481	10004056_c1_158	3137	8799	885	295	1324	3.0(10)-135	Escherichia coli	b3967	[pn:glutamate racemase] [gn:muri]
CONTIG481	29484442_c2_159	3138	8800	2487	829	3802	0	Escherichia coli	b3940	[pn:aspartokinase ii/homoserine dehydrogenase ii] [gn:metl]
CONTIG481	26765951_c2_174	3139	8801	339	113	436	3.7(10)-41	Escherichia coli	b3950	[pn:phosphotransfera] [gn:frwb]
CONTIG481	16110041_c2_175	3140	8802	2349	783	3547	0	Escherichia coli	b3951	[pn:formate acetyltransferase 2] [gn:pflD]

CONTIG481	26050393_c2_177	3141	8803	420	140	437	2.8(10)-41	Escherichia coli	b3953	[pn:phosphotransferase] [gn:frwd]
CONTIG481	5176592_c2_187	3142	8804	1014	338	1561	2.2(10)-160	Escherichia coli	b3958	[pn:n-acetyl-gamma-glutamyl-phosphate reductase] [gn:argc]
CONTIG481	5119143_c2_188	3143	8805	783	261	1143	4.5(10)-116	Escherichia coli	b3959	[pn:acetylglutamate kinase] [gn:argb]
CONTIG481	22135833_c2_189	3144	8806	1419	473	2100	1.7(10)-217	Escherichia coli	b3960	[pn:argininosuccinate lyase] [gn:argh]
CONTIG481	24788962_c2_192	3145	8807	807	269	942	9.0(10)-95	Escherichia coli	b3963	[pn:hypothetical 26.6 kd protein in udha-trma intergenic region]
CONTIG481	22462803_c2_193	3146	8808	378	126	463	5.2(10)-44	Escherichia coli	b3964	[pn:hypothetical 13.0 kd protein in udha-trma intergenic region]
CONTIG481	477291_c3_199	3147	8809	624	208	888	4.7(10)-89	Escherichia coli	b3939	[pn:cystathionine gamma-synthase] [gn:metb]
CONTIG481	665893_c3_206	3148	8810	894	298	1444	5.7(10)-148	Escherichia coli	b3941	[pn:5,10 methylenetetrahydrofolate reductase] [gn:metf]
CONTIG481	36126911_c3_207	3149	8811	2202	734	3234	0	Escherichia coli	b3942	[pn:catalase hydroperoxidase i] [gn:katg]
CONTIG481	5891450_c3_214	3150	8812	1131	377	1281	1.1(10)-130	Escherichia coli	b3949	[pn:phosphotransferase] [gn:frwc]
CONTIG481	22464582_c3_219	3151	8813	987	329	1123	5.9(10)-114	Escherichia coli	b3952	[pn:probable pyruvate formate-lyase 2 activating enzyme] [gn:pfic]
CONTIG481	22915793_c3_239	3152	8814	1890	630	2190	5.0(10)-227	Escherichia coli	b3966	[pn:vitamin b12 receptor precursor] [gn:btub]
CONTIG482	820216_fl_1	3153	8815	984	328	300	9.6(10)-27	Haemophilus influenzae	H11248	[pn:hypothetical protein]
CONTIG482	5115718_fl_6	3154	8816	1002	334	961	8.6(10)-97	Escherichia coli	b3909	[pn:2-keto-3-deoxygluconate permease] [gn:kdgt]
CONTIG482	24423878_fl_12	3155	8817	531	177	608	2.2(10)-59	Escherichia coli	b4107	[pn:protein] [gn:phnb]
CONTIG482	26602316_fl_13	3156	8818	1029	343	1594	7.2(10)-164	Escherichia coli	b4105	[pn:carbon phosphorus lyase] [gn:phnd]
CONTIG482	14317706_fl_18	3157	8819	669	223	837	1.2(10)-83	Escherichia coli	b4100	[pn:phnh protein] [gn:phnh]
CONTIG482	24066662_fl_21	3158	8820	870	290	1440	1.5(10)-147	Escherichia coli	b4098	[pn:phnj protein] [gn:phnj]
CONTIG482	34239388_fl_25	3159	8821	648	216	641	7.0(10)-63	Escherichia coli	b4094	[pn:phosphonates transport atp-binding protein phnn] [gn:phnn]

CONTIG482	17054132_f1_37	3160	8822	1080	360	499	7.9(10)-48	Bacillus subtilis	rbsC	[pn:ribose abc transporter]
CONTIG482	884702_f1_39	3161	8823	864	288	566	6.2(10)-55	Bacillus subtilis	fbaA	[pn:fructose-1,6-bisphosphate aldolase] [gn:tsr]
CONTIG482	24644438_f1_45	3162	8824	492	164	412	1.3(10)-38	Escherichia coli	b4078	[pn:hypothetical 25.1 kd protein in glp-fdhf intergenic region] [gn:yjco]
CONTIG482	32612590_f1_46	3163	8825	189	63	263	8.0(10)-23	Escherichia coli	b4078	[pn:hypothetical 25.1 kd protein in glp-fdhf intergenic region] [gn:yjco]
CONTIG482	10987586_f2_54	3164	8826	492	164	568	3.7(10)-55	Escherichia coli	b4108	[pn:phna protein] [gn:phna]
CONTIG482	2125183_f2_61	3165	8827	759	253	1098	2.6(10)-111	Escherichia coli	b4102	[pn:phnf protein] [gn:phnf]
CONTIG482	25785827_f2_62	3166	8828	453	151	522	2.8(10)-50	Escherichia coli	b4101	[pn:phng protein] [gn:phng]
CONTIG482	33854562_f2_69	3167	8829	1266	422	1630	1.1(10)-167	Escherichia coli	b4095	[pn:phnm protein] [gn:phnm]
CONTIG482	12927258_f2_70	3168	8830	774	258	598	2.5(10)-58	Escherichia coli	b4093	[pn:phno protein] [gn:phno]
CONTIG482	3958592_f2_71	3169	8831	768	256	1084	8.0(10)-110	Escherichia coli	b4092	[pn:phnp protein] [gn:phnp]
CONTIG482	4116642_f2_75	3170	8832	1548	516	1154	3.1(10)-117	Bacillus subtilis	rbsA	[pn:ribose abc transporter]
CONTIG482	13866580_f2_77	3171	8833	1008	336	332	3.8(10)-30	Bacillus subtilis	rbsB	[pn:ribose abc transporter]
CONTIG482	19567043_f3_99	3172	8834	816	272	1144	3.5(10)-116	Escherichia coli	b4106	[pn:phosphonates transport atp-binding protein] [gn:phnc]
CONTIG482	6767917_f3_102	3173	8835	846	282	607	2.7(10)-59	Escherichia coli	b4104	[pn:phne]
CONTIG482	16661008_f3_106	3174	8836	1170	390	1563	1.3(10)-160	Escherichia coli	b4099	[pn:phni protein] [gn:phni]
CONTIG482	35283517_f3_108	3175	8837	840	280	1119	1.6(10)-113	Escherichia coli	b4097	[pn:phosphonates transport atp-binding protein phnk] [gn:phnk]
CONTIG482	4488588_f3_109	3176	8838	690	230	1003	3.1(10)-101	Escherichia coli	b4096	[pn:phosphonates transport atp-binding protein phnl] [gn:phnl]
CONTIG482	19708567_f3_123	3177	8839	1083	361	152	2.0(10)-8	Bacillus subtilis	yjE	[pn:hypothetical protein]



CONTIG482	23447063_f3_127	3178	8840	444	148	701	3.1(10)-69	Escherichia coli	b4079	[pn:formate dehydrogenase, formate-hydrogen-lyase-linked, selenocysteine-containing polypeptide] [gn:fdhf]
CONTIG482	10646036_f3_128	3179	8841	1728	576	2902	1.8(10)-302	Escherichia coli	b4079	[pn:formate dehydrogenase, formate-hydrogen-lyase-linked, selenocysteine-containing polypeptide] [gn:fdhf]
CONTIG482	2756663_c2_176	3180	8842	189	63	175	1.0(10)-12	Escherichia coli	b4077	[pn:glutamate- aspartate carrier] [gn:gltp]
CONTIG482	14629593_c2_194	3181	8843	2310	770	593	8.5(10)-58	Rhizobium leguminosarum	Z70305	or:rhizobium leguminosarum pn:fixl gn:fixl   le:1546 re:3471 di:direct nt:putative heme-binding, oxygen sensing protein
CONTIG482	10337827_c2_205	3182	8844	237	79	276	3.3(10)-24	Escherichia coli	B35720	hypothetical 28.6k protein - escherichia coli
CONTIG482	4375276_c2_215	3183	8845	417	139	94	6.5(10)-5	Escherichia coli	C35720	hypothetical 12.4k protein - escherichia coli
CONTIG482	2538135_c2_221	3184	8846	909	303	788	1.8(10)-78	Escherichia coli	b4110	[pn:hypothetical 31.8 kd protein in phna-prop intergenic region] [gn:yjcz]
CONTIG482	22736068_c2_225	3185	8847	1587	529	2169	8.5(10)-225	Escherichia coli	b4111	[pn:proline/betaine transporter] [gn:prop]
CONTIG482	15807841_c3_231	3186	8848	744	248	369	4.7(10)-34	Escherichia coli	b3405	[pn:ompr] [gn:ompr]
CONTIG482	14112961_c3_242	3187	8849	240	80	173	2.7(10)-13	Escherichia coli	D35720	hypothetical 13.7k protein - escherichia coli
CONTIG482	25833337_c3_256	3188	8850	1023	341	172	3.5(10)-13	Escherichia coli	A35720	hypothetical 16.1k protein (phnq 3 region) - escherichiacoli
CONTIG482	135968_c3_261	3189	8851	2370	790	1679	9.0(10)-228	Escherichia coli	b4109	[pn:hypothetical 84.2 kd protein in phna-prop intergenic region] [gn:yjda]
CONTIG483	22834691_fl_6	3190	8852	891	297	1234	1.0(10)-125	Escherichia coli	b3455	[pn:high-affinity branched-chain amino acid transport atp-binding protein livg] [gn:livg]
CONTIG483	10677010_fl_8	3191	8853	1350	450	2061	2.3(10)-213	Escherichia coli	b3453	[pn:glycerol-3-phosphate-binding periplasmic protein precursor] [gn:ugpb]

CONTIG483	10634631_fl_9	3192	8854	891	297	1261	1.3(10)-128	Escherichia coli	b3452	[pn:sn-glycerol-3-phosphate transport system permease protein]
CONTIG483	16286306_fl_10	3193	8855	1119	373	1701	3.2(10)-175	Escherichia coli	b3450	[pn:sn-glycerol-3-phosphate transport atp-binding protein]
CONTIG483	2534500_fl_18	3194	8856	1083	361	1456	3.1(10)-149	Escherichia coli	b3438	[pn:gtmukr operon regulator]
CONTIG483	25509436_fl_19	3195	8857	549	183	687	9.4(10)-68	Escherichia coli	b3437	[pn:thermoresistant glucokinase]
CONTIG483	13946878_fl_20	3196	8858	1344	448	1223	1.5(10)-124	Escherichia coli	b3436	[pn:gntk]
CONTIG483	13132087_fl_25	3197	8859	2049	683	2777	3.2(10)-289	Escherichia coli	b3431	[pn:gntu_1]
CONTIG483	34010260_fl_35	3198	8860	927	309	1431	1.3(10)-146	Escherichia coli	b3458	[pn:glycogen operon protein glgx]
CONTIG483	6142567_fl_39	3199	8861	1362	454	1638	1.6(10)-168	Escherichia coli	b3456	[pn:leucine-specific binding protein precursor] [gn:livk]
CONTIG483	13854155_fl_40	3200	8862	741	247	1145	2.7(10)-116	Escherichia coli	b3454	[pn:high-affinity branched-chain amino acid transport permease protein livm] [gn:livm]
CONTIG483	5348437_fl_60	3201	8863	2205	735	3757	0	Escherichia coli	b3432	[pn:high-affinity branched-chain amino acid transport atp-binding] [gn:livf]
CONTIG483	3555135_fl_66	3202	8864	1647	549	2222	2.1(10)-230	Escherichia coli	b3429	[pn:1,4-alpha-glucan branching enzyme] [gn:glgb]
CONTIG483	6849191_fl_67	3203	8865	2466	822	3732	0	Escherichia coli	b3428	[pn:glycogen synthase] [gn:glga]
CONTIG483	22658562_fl_72	3204	8866	939	313	1163	3.3(10)-118	Escherichia coli	b3457	[pn:alpha-glucan phosphorylase] [gn:glgp]
CONTIG483	12526018_fl_79	3205	8867	858	286	1098	2.6(10)-111	Escherichia coli	b3451	[pn:high-affinity branched-chain amino acid transport permease protein livh] [gn:livh]
CONTIG483	14574037_fl_84	3206	8868	756	252	1016	1.3(10)-102	Escherichia coli	b3449	[pn:sn-glycerol-3-phosphate transport system permease protein]
CONTIG483	4105458_fl_85	3207	8869	1782	594	2482	5.7(10)-258	Escherichia coli	b3447	[pn:glycerophosphoryl diester phosphodiesterase] [gn:ugpd]
CONTIG483	2853408_fl_88	3208	8870	1041	347	1700	4.2(10)-175	Escherichia coli	b3440	[pn:gamma-glutamyltranspeptidase] [gn:ggd]
CONTIG483	16914058_fl_89	3209	8871	819	273	1145	2.7(10)-116	Escherichia coli	b3439	[pn:hypothetical 38.8 kd protein in gnt-r-ggt intergenic region] [gn:yhhx]
CONTIG483										[pn:hypothetical 26.3 kd protein in gnt-r-ggt intergenic region]

CONTIG483	20445390_f3_95	3210	8872	1173	391	1793	6.0(10)-185	Escherichia coli	b3433	[pn:aspartate semialdehyde dehydrogenase] [gn:asd]
CONTIG483	19767905_f3_104	3211	8873	1374	458	1994	3.0(10)-206	Escherichia coli	b3430	[pn:glucose-1-phosphate adenyllyltransferase] [gn:glgc]
CONTIG483	12605291_f3_113	3212	8874	258	86	94	0.00077	Saccharomyces cerevisiae	YNL138W	[pn:adenylate cyclase-associated protein, 70kda] [gn:sv2]
CONTIG483	20411717_c1_144	3213	8875	477	159	159	4.9(10)-17	Escherichia coli	b3448	[pn:hypothetical 16.6 kd protein in gg-ugpq intergenic region precursor] [gn:yhha]
CONTIG483	13019705_c2_186	3214	8876	318	106	104	5.7(10)-6	Escherichia coli	D90866	or:escherichia coli pn:d-serine dehydratase transcriptional activator gn:dsdc le:10864 re:11133 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [pir accession number a28674];
CONTIG483	17000951_c2_191	3215	8877	507	169	724	1.1(10)-71	Escherichia coli	b3441	[pn:hypothetical protein] [gn:yhhy]
CONTIG483	24725008_c3_207	3216	8878	1536	512	2144	3.7(10)-222	Escherichia coli	b3426	[pn:aerobic glycerol-3-phosphate dehydrogenase] [gn:glpd]
CONTIG483	34417125_c3_229	3217	8879	600	200	763	8.3(10)-76	Escherichia coli	b3434	[pn:hypothetical protein] [gn:yhgn]
CONTIG484	22737668_f1_1	3218	8880	639	213	782	8.0(10)-78	Escherichia coli	b0687	[pn:seqa protein] [gn:seqa]
CONTIG484	21665656_f1_2	3219	8881	1665	555	2446	3.7(10)-254	Escherichia coli	b0688	[pn:phosphoglucomutase] [gn:pgm]
CONTIG484	975428_f1_21	3220	8882	303	101	315	2.5(10)-28	Escherichia coli	b0699	[pn:hypothetical 8.3 kd protein in rhsc 5'''region] [gn:ybfa]
CONTIG484	25494783_f1_25	3221	8883	1125	375	1214	1.3(10)-123	Escherichia coli	b0712	[pn:hypothetical protein] [gn:ybgk]
CONTIG484	16618761_f1_31	3222	8884	501	167	604	5.9(10)-59	Escherichia coli	b0721	[pn:succinate dehydrogenase cytochrome b-556 subunit] [gn:sdhc]
CONTIG484	15663465_f2_67	3223	8885	1434	478	1878	5.9(10)-194	Escherichia coli	b0708	[pn:deoxyribodipyrimidine photolyase] [gn:phrb]
CONTIG484	16667078_f2_68	3224	8886	657	219	989	9.4(10)-100	Escherichia coli	b0711	[pn:hypothetical protein] [gn:ybgj]
CONTIG484	33878962_f2_69	3225	8887	792	264	887	6.0(10)-89	Escherichia coli	b0713	[pn:hypothetical protein] [gn:ybgj]

CONTIG484	23629837_f2_70	3226	8888	804	268	1138	1.5(10)-115	Escherichia coli	b0714	[pn:endonuclease viii, dna n-glycosylase with an ap lyase activity] [gn:nei]
CONTIG484	26063532_f2_72	3227	8889	1860	620	2955	0	Escherichia coli	b0723	[pn:succinate dehydrogenase flavoprotein subunit] [gn:sdha]
CONTIG484	33805415_f2_73	3228	8890	732	244	1234	1.0(10)-125	Escherichia coli	b0724	[pn:succinate dehydrogenase iron-sulfur protein] [gn:sdhb]
CONTIG484	32235416_f2_75	3229	8891	1260	420	1765	5.5(10)-182	Escherichia coli	b0727	[pn:e2] [gn:subb]
CONTIG484	25495836_f2_76	3230	8892	948	316	1117	2.6(10)-113	Escherichia coli	b0729	[pn:succinyl-coa synthetase alpha chain] [gn:sucd]
CONTIG484	24726711_f2_80	3231	8893	1686	562	2440	1.6(10)-253	Escherichia coli	b0732	[pn:ybgg] [gn:ybgg]
CONTIG484	6347132_f3_100	3232	8894	834	278	1208	5.7(10)-123	Escherichia coli	b0710	[pn:hypothetical protein] [gn:ybgi]
CONTIG484	29331261_f3_109	3233	8895	399	133	470	9.3(10)-45	Escherichia coli	b0722	[pn:succinate dehydrogenase 13 kd hydrophobic protein] [gn:sdhd]
CONTIG484	5190968_f3_114	3234	8896	2817	939	4678	0	Escherichia coli	b0726	[pn:2-oxoglutarate dehydrogenase e1 component] [gn:suca]
CONTIG484	25490691_f3_116	3235	8897	1194	398	1622	7.9(10)-167	Escherichia coli	b0728	[pn:succinyl-coa synthetase beta chain] [gn:succ]
CONTIG484	433468_f3_119	3236	8898	1929	643	2228	4.7(10)-231	Escherichia coli	b0731	[pn:heat-responsive regulatory protein] [gn:hrrsa]
CONTIG484	4586665_c1_147	3237	8899	1695	565	2180	5.7(10)-226	Escherichia coli	b0698	[pn:potassium-transporting atpase, a chain] [gn:kdpa]
CONTIG484	26458125_c1_148	3238	8900	2067	689	2744	1.0(10)-285	Escherichia coli	b0697	[pn:potassium-transporting atpase, b chain] [gn:kdpb]
CONTIG484	11931712_c1_149	3239	8901	588	196	632	6.4(10)-62	Escherichia coli	b0696	[pn:potassium-transporting atpase, c chain] [gn:kdpc]
CONTIG484	25988452_c1_150	3240	8902	2688	896	3570	0	Escherichia coli	b0695	[pn:sensor protein kdpd] [gn:kdpd]
CONTIG484	1423512_c1_151	3241	8903	195	65	90	0.00017	Haemophilus influenzae	HI0592	[pn:hypothetical protein]
CONTIG484	5289012_c1_152	3242	8904	2214	738	3359	0	Escherichia coli	b0693	[pn:ornithine decarboxylase, inducible] [gn:spet]
CONTIG484	24422650_c2_163	3243	8905	750	250	904	9.5(10)-91	Escherichia coli	b0730	[pn:fatty acyl responsive regulator] [gn:farr]
CONTIG484	10413587_c2_177	3244	8906	1221	407	1940	1.6(10)-200	Escherichia coli	b0720	[pn:gltA] [gn:gltA]

CONTIG484	36207127_c3_243	3245	8907	693	231	871	3.0(10)-87	Escherichia coli	b0694	[pn:kdp operon transcriptional regulatory protein kdpE] [gn:kdpe]
CONTIG484	14635091_c3_246	3246	8908	1389	463	1945	4.7(10)-201	Escherichia coli	b0692	[pn:putrescine-ornithine antiporter] [gn:pote]
CONTIG485	1992963_fl_1	3247	8909	531	177	215	9.8(10)-18	Saccharomyces cerevisiae	S62019	hypothetical protein ydr540c - yeast (saccharomycetaceae)
CONTIG485	26595012_fl_2	3248	8910	600	200	293	5.2(10)-26	Thiobacillus ferrooxidans	U66426	or:thiobacillus ferrooxidans
CONTIG485	25681417_fl_3	3249	8911	315	105	117	1.8(10)-6	Ralstonia eutropha	A47041	pn:transposase le:73 re:1284
CONTIG485	5985443_fl_60	3250	8912	807	269	1184	2.0(10)-120	Escherichia coli	P15026	transposase homolog (insertion element isae1) - alcaligeneseutrophus
CONTIG485	14238143_fl_100	3251	8913	1200	400	1910	2.3(10)-197	Escherichia coli	P15025	istb protein (insertion sequence is21).
CONTIG485	14657086_fl_101	3252	8914	333	111	132	4.2(10)-8	Bordetella parapertussis	Q06126	ista protein (insertion sequence is21).
CONTIG485	25491665_cl_154	3253	8915	597	199	587	3.7(10)-57	Escherichia coli	b4059	transposase for insertion sequence element is1001.
CONTIG485	15027217_cl_162	3254	8916	2112	704	107	0.01099	Oenococcus oeni	S42039	[pn:single-strand dna-binding protein] [gn:ssb]
CONTIG485	22933541_c2_190	3255	8917	2025	675	1098	2.6(10)-111	Haemophilus influenzae	H10444	hypothetical protein 2 - leuconostoc oenos
CONTIG485	34412908_c2_198	3256	8918	579	193	345	1.6(10)-31	Salmonella typhimurium	AF000360	[pn:topoisomerase iii] [gn:topb]
CONTIG485	34415760_c2_216	3257	8919	1482	494	99	0.17999	Escherichia coli	b1372	[PN:hypothetical protein] [DE:Salmonella typhimurium IncN plasmid pKM101 IS26 (IS46) element TnpA(tpnA) gene, complete cds.] [LE:1033] [RE:1674] [DI:complement]
CONTIG485	1258591_c3_230	3258	8920	294	98	100	1.5(10)-5	Synechocystis sp.	S77531	[pn:hypothetical protein]
CONTIG485	13167787_c3_248	3259	8921	1518	506	97	0.42999	Saccharomyces cerevisiae	YHR023 W	[OR:Synechocystis sp.] [SR:PCC 6803, PCC 6803] [SR:PCC 6803, ] [pn:myo ii myosin heavy chain] [gn:myo1]
CONTIG485	32714125_c3_266	3260	8922	327	109	208	5.4(10)-17	Escherichia coli	b1183	[pn:umud protein] [gn:umud]

CONTIG486	5112518_f1_1	3261	8923	1428	476	1982	5.5(10)-205	Escherichia coli	b4240	[pn:phosphotransferase system trehalose permease] [gn:treb]
CONTIG486	9781900_f1_2	3262	8924	1695	565	2480	9.4(10)-258	Escherichia coli	b4239	[pn:trehalose-6-phosphate hydrolase] [gn:trec]
CONTIG486	7204593_f1_4	3263	8925	483	161	778	2.1(10)-77	Escherichia coli	b4237	[pn:anaerobic ribonucleoside-triphosphate reductase activating pr] [gn:nrdg]
CONTIG486	897326_f1_24	3264	8926	588	196	785	3.8(10)-78	Escherichia coli	b4234	[pn:x] [gn:yjga]
CONTIG486	4947281_f1_36	3265	8927	468	156	318	1.2(10)-28	Escherichia coli	b4223	[pn:ile repressor protein] [gn:yjfa]
CONTIG486	4410943_f1_51	3266	8928	567	189	681	4.0(10)-67	Escherichia coli	b4216	[pn:f18] [gn:ytfj]
CONTIG486	3219057_f1_54	3267	8929	2073	691	2954	0	Escherichia coli	b4213	[pn:2'''',3'''-cyclic-nucleotide 2'''-phosphodiesterase] [gn:cpdb]
CONTIG486	5133512_f2_59	3268	8930	2151	717	3486	0	Escherichia coli	b4238	[pn:anaerobic ribonucleoside-triphosphate reductase] [gn:nrdd]
CONTIG486	15080013_f2_78	3269	8931	1050	350	1627	2.2(10)-167	Escherichia coli	b4232	[pn:fructose-1,6-bisphosphatase] [gn:fbp]
CONTIG486	36536467_f2_87	3270	8932	630	210	899	3.2(10)-90	Escherichia coli	b4226	[pn:pyrophosphate phospho] [gn:ppa]
CONTIG486	32210956_f2_102	3271	8933	672	224	910	2.2(10)-91	Escherichia coli	b4219	[pn:peptide methionine sulfoxide reductase] [gn:msra]
CONTIG486	24228382_f3_156	3272	8934	1341	447	1948	2.2(10)-201	Escherichia coli	b4218	[pn:hypothetical 49.8 kd protein in cysq-msra intergenic region]
CONTIG486	26460400_c1_163	3273	8935	225	75	335	1.8(10)-30	Escherichia coli	b4217	[pn:hypothetical 9.6 kd protein in cysq-msra intergenic region] [gn:ytfk]
CONTIG486	275252_c1_167	3274	8936	2940	980	4140	0	Escherichia coli	b4221	[pn:hypothetical 136.8 kd protein in msra-chpb intergenic region] [gn:ytfh]
CONTIG486	25439541_c1_168	3275	8937	834	278	1142	5.7(10)-116	Escherichia coli	b4221	[pn:hypothetical 136.8 kd protein in msra-chpb intergenic region] [gn:ytfh]
CONTIG486	32507077_c1_174	3276	8938	1053	351	1069	3.1(10)-108	Escherichia coli	b4230	[pn:hypothetical 35.7 kd protein in ppa-fbp intergenic region] [gn:ytfh]
CONTIG486	2050705_c1_192	3277	8939	1209	403	332	3.8(10)-30	Bacillus subtilis	ykgB	[pn:hypothetical protein]

CONTIG486	14165878_c2_209	3278	8940	774	258	1110	1.3(10)-112	Escherichia coli	b4214	[pn:csq protein] [gn:csq]
CONTIG486	35162813_c2_216	3279	8941	1860	620	2678	9.8(10)-279	Escherichia coli	b4220	[pn:hypothetical 64.8 kd protein in msra-chpbi intergenic region] [gn:ytfm]
CONTIG486	11027152_c2_222	3280	8942	1089	363	1366	1.1(10)-139	Escherichia coli	b4231	[pn:hypothetical 34.0 kd protein in ppa-fbp intergenic region] [gn:yfff]
CONTIG486	2775841_c2_223	3281	8943	1623	541	863	2.1(10)-86	Escherichia coli	b1886	[pn:methyl-accepting chemotaxis protein ii] [gn:tar]
CONTIG486	4556516_c2_226	3282	8944	1377	459	2270	1.7(10)-235	Escherichia coli	b4233	[pn:hypothetical 48.5 kd protein in fbp-pmba intergenic region]
CONTIG486	17057035_c2_235	3283	8945	1176	392	102	0.019	Escherichia coli	b0512	[pn:hypothetical protein] [gn:ybbx]
CONTIG486	25973588_c2_239	3284	8946	1923	641	567	4.9(10)-55	Bacillus subtilis	licR	[pn:transcriptional regulator] [gn:celr]
CONTIG486	4818753_c3_259	3285	8947	354	118	554	1.2(10)-53	Escherichia coli	b4222	[pn:hypothetical 12.9 kd protein in msra-chpbi intergenic region] [gn:ytfp]
CONTIG486	26054762_c3_261	3286	8948	1104	368	1511	4.5(10)-155	Escherichia coli	b4227	[pn:o318] [gn:ytfq]
CONTIG486	31719406_c3_262	3287	8949	1614	538	1694	1.8(10)-174	Escherichia coli	b4228	[pn:hypothetical abc transporter in ppa-fbp intergenic region] [gn:yftr]
CONTIG486	3401562_c3_268	3288	8950	1434	478	2089	2.6(10)-216	Escherichia coli	b4235	[pn:pmba protein] [gn:pmba]
CONTIG486	23697183_c3_269	3289	8951	390	130	376	8.5(10)-35	Escherichia coli	b4236	[pn:cytochrome b562] [gn:cybc]
CONTIG486	36520331_c3_274	3290	8952	1140	380	140	1.3(10)-6	Haemophilus influenzae	H10708	[pn:l-seryl-trna] [gn:sela]
CONTIG487	4689187_fl_1	3291	8953	789	263	1065	8.3(10)-108	Escherichia coli	b3364	[pn:hypothetical 43.2 kd protein in ppia-nirb intergenic region]
CONTIG487	35626332_fl_4	3292	8954	360	120	481	6.4(10)-46	Escherichia coli	b3366	[pn:nadh small subunit] [gn:nird]
CONTIG487	6439205_fl_31	3293	8955	756	252	1041	2.8(10)-105	Escherichia coli	b3399	[pn:hypothetical 27.1 kd protein in mreca-pcka intergenic region] [gn:yrfg]
CONTIG487	26612631_fl_44	3294	8956	249	83	335	1.8(10)-30	Escherichia coli	b3408	[pn:ferrous iron transport protein a] [gn:feoa]
CONTIG487	35816686_fl_45	3295	8957	2349	783	3358	0	Escherichia coli	b3409	[pn:ferrous iron transport protein b] [gn:feob]

CONTIG487	4039033_f1_47	3296	8958	585	195	962	6.7(10)-97	Escherichia coli	b3414	[pn:hypothetical 21.0 kd protein in bioh-gntt intergenic region]
CONTIG487	30331875_f2_49	3297	8959	747	249	647	1.6(10)-63	Escherichia coli	b3364	[pn:hypothetical 43.2 kd protein in ppia-nirb intergenic region]
CONTIG487	25517283_f2_50	3298	8960	2469	823	3923	0	Escherichia coli	b3365	[pn:nadh-nitrate oxidoreductase apoprotein] [gn:nirb]
CONTIG487	22696061_f2_53	3299	8961	183	61	155	2.2(10)-11	Escherichia coli	b3369	[pn:hypothetical 5.8 kd protein in cysg-trps intergenic region] [gn:yhrf]
CONTIG487	15652152_f2_70	3300	8962	2625	875	3985	0	Escherichia coli	b3396	[pn:mrca]
CONTIG487	29926075_f2_88	3301	8963	351	117	359	5.4(10)-33	Escherichia coli	b3410	[pn:hypothetical 8.7 kd protein in feob-bioh intergenic region] [gn:yhgg]
CONTIG487	6301518_f3_102	3302	8964	1383	461	2088	3.2(10)-216	Escherichia coli	b3368	[pn:siroheme synthase] [gn:cysg]
CONTIG487	34199138_f3_122	3303	8965	2151	717	2807	2.1(10)-292	Escherichia coli	b3398	[pn:hypothetical 79.5 kd protein in mrca-pcka intergenic region] [gn:yrfi]
CONTIG487	10573437_f3_123	3304	8966	477	159	545	1.1(10)-52	Escherichia coli	b3400	[pn:hypothetical 15.5 kd protein in mrca-pcka intergenic region] [gn:yrfh]
CONTIG487	15725656_f3_124	3305	8967	903	301	1313	4.4(10)-134	Escherichia coli	b3401	[pn:hypothetical 32.8 kd protein in mrca-pcka intergenic region] [gn:yrfi]
CONTIG487	24785136_f3_129	3306	8968	1677	559	2513	3.0(10)-261	Escherichia coli	b3403	[pn:phosphoenolpyruvate carboxykinase] [gn:pcka]
CONTIG487	14491316_f3_132	3307	8969	801	267	154	3.1(10)-11	Rhizobium sp.	S28675	hypothetical protein 5 - rhizobium sp. (strain ic3342)
CONTIG487	32158437_f3_133	3308	8970	654	218	776	3.5(10)-77	Escherichia coli	b3406	[pn:transcription elongation factor greb] [gn:greb]
CONTIG487	16292280_f3_134	3309	8971	2418	806	3440	0	Escherichia coli	b3407	[pn:hypothetical 81.4 kd protein in greb-feoa intergenic region] [gn:yhgf]
CONTIG487	7207652_f3_140	3310	8972	855	285	770	1.5(10)-76	Escherichia coli	b3413	[pn:hypothetical 27.7 kd protein in bioh-gntt intergenic region]
CONTIG487	6454662_f3_143	3311	8973	1200	400	1635	3.2(10)-168	Escherichia coli	b3415	[pn:high-affinity gluconate transporter] [gn:gntt]
CONTIG487	32697167_c1_153	3312	8974	1350	450	1961	9.4(10)-203	Escherichia coli	b3404	[pn:osmolarity sensor protein envz] [gn:envz]



CONTIG487	10978206_c1_170	3313	8975	486	162	189	2.6(10)-14	Escherichia coli	b3391	[pn:protein transport protein hofq precursor] [gn:hofq]
CONTIG487	9767138_c1_172	3314	8976	642	214	940	1.5(10)-94	Escherichia coli	b3390	[pn:shikimic acid kinase i] [gn:arok]
CONTIG487	31492893_c1_173	3315	8977	1104	368	1535	1.3(10)-157	Escherichia coli	b3389	[pn:3-dehydroquinate synthase] [gn:arob]
CONTIG487	30366667_c1_176	3316	8978	873	291	1187	9.8(10)-121	Escherichia coli	b3387	[pn:dna adenine methylase] [gn:dam]
CONTIG487	23829557_c1_177	3317	8979	699	233	1124	4.5(10)-114	Escherichia coli	b3386	[pn:dod protein] [gn:rpe]
CONTIG487	31886465_c1_180	3318	8980	252	84	90	0.00119	Plasmid R751	L13688	or:plasmid r751 gn:kfra le:2408 re:3466 di:direct sr:plasmid r751 dna
CONTIG487	15040951_c2_196	3319	8981	786	262	1084	8.0(10)-110	Escherichia coli	b3412	[pn:bioh] [gn:bioh]
CONTIG487	5864427_c2_205	3320	8982	834	278	1212	2.2(10)-123	Escherichia coli	b3405	[pn:ompr] [gn:ompr]
CONTIG487	14073292_c2_219	3321	8983	591	197	885	9.9(10)-89	Escherichia coli	b3397	[pn:hypothetical 21.2 kd protein in mra-pcka intergenic region] [gn:yrfc]
CONTIG487	2207675_c2_223	3322	8984	534	178	172	3.5(10)-13	Escherichia coli	b3394	[pn:hypothetical 20.8 kd protein in hofq-mrca intergenic region] [gn:yrfc]
CONTIG487	4197331_c2_224	3323	8985	456	152	197	7.9(10)-16	Escherichia coli	b3393	[pn:hypothetical 16.8 kd protein in hofq-mrca intergenic region] [gn:yrfb]
CONTIG487	32557191_c2_226	3324	8986	315	105	94	0.00022	Escherichia coli	b3390	[pn:shikimic acid kinase i] [gn:arok]
CONTIG487	23650278_c2_230	3325	8987	765	255	1045	1.1(10)-105	Escherichia coli	b3385	[pn:hypothetical 27.4 kd protein in trps-dod intergenic region] [gn:gph]
CONTIG487	24736537_c3_243	3326	8988	501	167	118	1.8(10)-7	Escherichia coli	D90866	or:escherichia coli pn:d-serine dehydratase transcriptional activator gn:dsdc le:10864 re:11133 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [pir accession number a28674];
CONTIG487	24431563_c3_258	3327	8989	1761	587	950	1.3(10)-95	Escherichia coli	b3402	[pn:hypothetical 64.6 kd protein in mra-pcka intergenic region] [gn:yhge]

CONTIG487	134667_c3_269	3328	8990	777	259	535	1.2(10)-51	Escherichia coli	b3395	[pn:hypothetical 30.0 kd protein in hofq-mrca intergenic region] [gn:yrfd]
CONTIG487	11720828_c3_270	3329	8991	468	156	295	3.2(10)-26	Escherichia coli	b3392	[pn:hypothetical 16.9 kd protein in hofq-mrca intergenic region] [gn:yrfa]
CONTIG487	35441390_c3_271	3330	8992	1149	383	1180	5.4(10)-120	Escherichia coli	b3391	[pn:protein transport protein hofq precursor] [gn:hofq]
CONTIG487	986077_c3_272	3331	8993	1290	430	902	1.1(10)-105	Escherichia coli	b3388	[pn:damx protein] [gn:damx]
CONTIG487	23556552_c3_273	3332	8994	1008	336	1646	2.2(10)-169	Escherichia coli	b3384	[pn:tryptophanyl-tRNA synthetase] [gn:trps]
CONTIG488	26458418_f1_32	3333	8995	1638	546	171	7.2(10)-10	Escherichia coli	b0591	[pn:hypothetical membrane protein p43] [gn:ybda]
CONTIG488	5960056_f1_33	3334	8996	846	282	963	5.2(10)-97	Pseudomonas fluorescens	U12537	or:pseudomonas fluorescens pn:esterase le:248 re:1066 di:direct
CONTIG488	25448950_f1_48	3335	8997	1017	339	119	0.00014	Neisseria gonorrhoeae	AF001598	[de:neisseria gonorrhoeae strain wr302 tex gene, partial cds, 5-methylcytosine methyltransferase (dcmh), n6-methyladeninemethyltransferase (damh) and restriction endonuclease (dc") [pn:restriction endonuclease] [gn:dcrh]
CONTIG488	31343942_f2_70	3336	8998	858	286	92	0.094	Pseudomonas alcaligenes	U84154	or:pseudomonas alcaligenes pn:putative transposase subunit gn:orf2 le:1104 re:2426 di:direct
CONTIG488	26457290_f2_74	3337	8999	1179	393	168	5.0(10)-10	Escherichia coli	b4345	[pn:mrcr protein] [gn:mrcr]
CONTIG488	16023540_f2_78	3338	9000	1035	345	334	2.3(10)-30	Bacillus subtilis	degA	[pn:transcriptional regulator]
CONTIG488	4476702_f3_93	3339	9001	618	206	456	2.7(10)-43	Pseudomonas sp.	AJ000330	[de:pseudomonas sp. dna for styrene catabolism genes.] [pn:styrene response regulator] [gn:styr]
CONTIG488	2086630_f3_108	3340	9002	1884	628	393	3.2(10)-36	Bacillus subtilis	yticJ	[pn:hypothetical protein]
CONTIG488	21596925_f3_115	3341	9003	2028	676	549	4.0(10)-53	Escherichia coli	b4346	[pn:5-methylcytosine-specific restriction enzyme b] [gn:mrcb]

CONTIG488	3909818_f3_122	3342	9004	1014	338	825	2.2(10)-82	Clostridium longisporum	Q46127	tryptophanyl-trna synthetase (ec 6.1.1.2) (tryptophan--trna ligase) (trprs).
CONTIG488	35214849_c1_125	3343	9005	370	123	498	1.0(10)-47	Enterobacter agglomerans	B38965	hypothetical protein b (insertion sequence is l222) -enterobacter agglomerans
CONTIG488	23930317_c1_134	3344	9006	414	138	101	0.00019	Pseudomonas aeruginosa	Z54213	or:pseudomonas aeruginosa pn:alginate lyase gn:algy le:1820 re:3874 di:direct
CONTIG488	1192705_c1_160	3345	9007	1155	385	1353	2.5(10)-138	Escherichia coli	b1128	[pn:hypothetical protein in pept-phoq intergenic region] [gn:ycfd]
CONTIG488	34647887_c2_169	3346	9008	516	172	186	1.2(10)-14	Vibrio cholerae	S81006	or:vibrio cholerae pn:hcp gn:hcp le:690 re:l208 di:direct sr:vibrio cholerae o17 nt:28 kda secreted hydrophilic protein; this sequence unknown, mtcy l64.07, mtcy l64.07. unknown, len
CONTIG488	6725052_c2_174	3347	9009	1113	371	602	9.5(10)-59	Mycobacterium tuberculosis	Z95150	[pn:hypothetical protein]
CONTIG488	16300816_c2_175	3348	9010	1434	478	425	5.5(10)-40	Bacillus subtilis	yjmB	
CONTIG488	25892702_c2_189	3349	9011	468	156	151	5.9(10)-11	Haemophilus influenzae	H10884	[pn:aerobic respiration control protein arca] [gn:arca]
CONTIG488	5338193_c2_190	3350	9012	5595	1865	235	1.0(10)-15	Bacillus subtilis	yloP	[pn:hypothetical protein]
CONTIG488	16586438_c2_192	3351	9013	630	210	406	5.7(10)-38	Bacillus subtilis	clpP	[pn:class iii heat-shock protein] [gn:yvdr]
CONTIG488	884575_c2_194	3352	9014	1017	339	307	1.7(10)-27	Bordetella pertussis	AF018255	[de:bordetella pertussis alcaligin siderophore system regulator (alcr)gene, complete cds.] [pn:alcr ] [gn:alcr] [nt:member of arac family; regulator of alcaligin]
CONTIG488	5908451_c3_195	3353	9015	279	93	327	1.3(10)-29	Escherichia coli	b4308	[pn:hypothetical 38.0 kd protein in fecI-fimb intergenic region] [gn:yjhr]
CONTIG488	4071963_c3_197	3354	9016	477	159	95	0.00013	Escherichia coli	b4347	[pn:hypothetical 14.6 kd protein in merb-hsdS intergenic region] [gn:yjiw]

CONTIG488	15814391_c3_203	3355	9017	612	204	734	9.9(10)-73	Escherichia coli	A60635	glutathione transferase (ec 2.5.1.18), fosfomycin-modifying -escherichia coli plasmid psu961 transposon tn2921 this enzyme inactivates the antibiotic phosphomycin by opening the epoxide ring and creating an adduct with glutathione.
CONTIG488	10271881_c3_220	3356	9018	711	237	259	2.1(10)-22	Escherichia coli	b0897	[pn:hypothetical 23.1 kd protein in dmsc 3'''region] [gn:ycac]
CONTIG488	2037813_c3_233	3357	9019	2112	704	1451	1.0(10)-148	Erwinia amylovora	AJ223062	[de:erwinia amylovora foxr gene.]
CONTIG489	7120253_fl_1	3358	9020	1143	381	1780	1.3(10)-183	Escherichia coli	b2097	[pn:ferrioxamine receptor] [gn:foxr]
CONTIG489	17065668_fl_38	3359	9021	618	206	936	3.8(10)-94	Escherichia coli	b2065	[pn:hypothetical protein]
CONTIG489	22299168_fl_45	3360	9022	2235	745	2883	1.8(10)-300	Escherichia coli	b2060	[pn:deoxycytidine triphosphate deaminase] [gn:ded]
CONTIG489	36407341_fl_47	3361	9023	510	170	780	1.3(10)-77	Escherichia coli	b2058	[pn:hypothetical protein]
CONTIG489	14163325_fl_49	3362	9024	1116	372	1176	1.3(10)-119	Escherichia coli	b2056	[pn:putative acetyl transferase] [gn:wcab]
CONTIG489	24650305_f2_74	3363	9025	1140	380	1771	1.3(10)-182	Escherichia coli	b2062	[pn:hypothetical protein] [gn:wza]
CONTIG489	1213533_f2_75	3364	9026	450	150	606	3.6(10)-59	Escherichia coli	b2061	[pn:hypothetical protein] [gn:wzb]
CONTIG489	4101718_f2_79	3365	9027	852	284	1378	5.5(10)-141	Escherichia coli	b2059	[pn:hypothetical protein] [gn:wcaa]
CONTIG489	4782830_f3_108	3366	9028	1020	340	1061	2.2(10)-107	Escherichia coli	b2068	[pn:dna-3-methyladenine glycosidase ii] [gn:alka]
CONTIG489	21900836_f3_113	3367	9029	534	178	95	0.01799	Methanobacterium thermoautotrophicum	MTH954	[pn:atp synthase, subunit b]
CONTIG489	6535041_f3_116	3368	9030	693	231	1071	1.8(10)-108	Escherichia coli	b2066	[pn:uridine kinase] [gn:udk]
CONTIG489	29728317_f3_117	3369	9031	1881	627	2079	2.8(10)-215	Escherichia coli	b2064	[pn:yega] [gn:asma]
CONTIG489	29504766_f3_127	3370	9032	1308	436	1789	1.6(10)-184	Escherichia coli	b2057	[pn:hypothetical protein] [gn:wcaac]

CONTIG489	24395411_c1_168	3371	9033	1449	483	1938	2.6(10)-200	Escherichia coli	b2078	[pn:sensor protein baes] [gn:baes]
CONTIG489	13906637_c2_191	3372	9034	3417	1139	4069	0	Escherichia coli	b2067	[pn:hypothetical 123.9 kd protein in udk-alka intergenic region] [gn:yege]
CONTIG489	31844831_c2_194	3373	9035	3186	1062	3946	0	Escherichia coli	b2075	[pn:hypothetical protein]
CONTIG489	10953_c2_195	3374	9036	3078	1026	4028	0	Escherichia coli	b2076	[pn:hypothetical protein]
CONTIG489	2550952_c2_196	3375	9037	1416	472	1529	5.5(10)-157	Escherichia coli	b2077	[pn:hypothetical protein] [gn:yege]
CONTIG489	650700_c2_197	3376	9038	357	119	91	0.00013	Daucus carota	U47097	or:daucus carota pn:glycine-rich protein le:<1 re:336 di:direct sr:carrot strain=danver half-long
CONTIG489	33628506_c2_202	3377	9039	1386	462	2084	8.6(10)-216	Escherichia coli	b2081	[pn:hypothetical protein] [gn:yege]
CONTIG489	25635955_c3_216	3378	9040	1629	543	2101	1.3(10)-217	Escherichia coli	b2063	[pn:hypothetical protein]
CONTIG489	10820407_c3_228	3379	9041	1452	484	1977	1.8(10)-204	Escherichia coli	b2069	[pn:yege]
CONTIG489	14486438_c3_230	3380	9042	1215	405	1533	2.1(10)-157	Escherichia coli	b2074	[pn:hypothetical protein]
CONTIG489	33473136_c3_236	3381	9043	768	256	1094	7.0(10)-111	Escherichia coli	b2079	[pn:transcriptional regulatory protein baer] [gn:baer]
CONTIG489	36523426_c3_241	3382	9044	921	307	1081	1.7(10)-109	Escherichia coli	b2086	[pn:hypothetical protein] [gn:vegs]
CONTIG49	31437686_c1_1	3383	9045	285	95	91	0.00083	Escherichia coli	D90807	or:escherichia coli pn:fusaric acid resistance protein fuse. le:18869 re:19858 di:direct sr:escherichia coli (strain:k12) dna_clone_lib:kohara lambda minise nt:orf_id:o316#23; similar to [swissprot accession]
CONTIG490	32464583_f1_1	3384	9046	1392	464	1788	2.0(10)-184	Escherichia coli	b3035	[pn:tolc] [gn:tolc]
CONTIG490	35644442_f1_2	3385	9047	204	68	160	6.5(10)-12	Escherichia coli	b3036	[pn:hypothetical 9.9 kd protein in tole-ribb/htrp intergenic region]
CONTIG490	3379790_f1_3	3386	9048	1200	400	1939	2.0(10)-200	Escherichia coli	b3038	[pn:hypothetical 45.0 kd protein in tole-ribb/htrp intergenic region] [gn:ygie]

CONTIG490	23632140_f1_6	3387	9049	2532	844	2377	7.7(10)-247	Escherichia coli	b0718	[pn:hypothetical protein]
CONTIG490	4331563_f1_7	3388	9050	771	257	698	6.4(10)-69	Escherichia coli	b0717	[pn:hypothetical protein]
CONTIG490	10672833_f1_31	3389	9051	600	200	350	4.9(10)-32	Bacillus subtilis	ureA	[pn:urease]
CONTIG490	22917040_f1_40	3390	9052	255	85	357	8.8(10)-33	Escherichia coli	b3065	[pn:30s ribosomal subunit protein s21] [gn:rpsu]
CONTIG490	4103953_f1_42	3391	9053	1980	660	2670	6.9(10)-278	Escherichia coli	b3067	[pn:rna polymerase sigma-70 factor] [gn:rpoD]
CONTIG490	554651_f1_44	3392	9054	1203	401	93	0.34999	Escherichia coli	b2592	[pn:clpb protein] [gn:clpb]
CONTIG490	24707030_f2_54	3393	9055	783	261	1044	1.3(10)-105	Escherichia coli	b3037	[pn:hypothetical 24.9 kd protein in tole-ribb/htrp intergenic region] [gn:ygiB]
CONTIG490	14932812_f2_64	3394	9056	375	125	290	1.1(10)-25	Escherichia coli	b0716	[pn:hypothetical protein]
CONTIG490	32304812_f2_79	3395	9057	1578	526	207	7.4(10)-14	Bacillus subtilis	yfiG	[pn:hypothetical protein]
CONTIG490	35430415_f2_80	3396	9058	846	282	931	1.3(10)-93	Klebsiella aerogenes	Q09063	urease operon ured protein.
CONTIG490	823905_f2_81	3397	9059	399	133	317	1.5(10)-28	Haemophilus influenzae	H10540	[pn:urease beta subunit] [gn:ureb]
CONTIG490	30718912_f2_87	3398	9060	726	242	817	1.6(10)-81	Klebsiella aerogenes	P18318	urease accessory protein uref.
CONTIG490	4097318_f2_88	3399	9061	627	209	691	3.6(10)-68	Helicobacter pylori	HP0068	[pn:urease accessory protein] [gn:ureg]
CONTIG490	9892668_f2_91	3400	9062	1812	604	2662	4.9(10)-277	Escherichia coli	b3066	[pn:dna primase] [gn:dnag]
CONTIG490	4492818_f2_94	3401	9063	504	168	526	1.1(10)-50	Escherichia coli	b3071	[pn:hypothetical protein]
CONTIG490	4539193_f3_102	3402	9064	783	261	1124	4.5(10)-114	Escherichia coli	b3040	[pn:hypothetical 26.5 kd protein in tole-ribb/htrp intergenic region] [gn:ygie]
CONTIG490	24235787_f3_103	3403	9065	681	227	376	8.5(10)-35	Escherichia coli	b0719	[pn:hypothetical fimbrial-like protein in glta 3''' region] [gn:ybgd]
CONTIG490	36051013_f3_104	3404	9066	741	247	739	2.8(10)-73	Escherichia coli	b0716	[pn:hypothetical protein]

CONTIG490	4797911_f3_107	3405	9067	342	114	424	7.0(10)-40	Escherichia coli	b3042	[pn:hypothetical protein]
CONTIG490	5086400_f3_121	3406	9068	702	234	852	3.1(10)-85	Escherichia coli	b3055	[pn:hypothetical protein in glne-cca intergenic region] [gn:ygim]
CONTIG490	10159776_f3_122	3407	9069	1302	434	1840	6.2(10)-190	Escherichia coli	b3056	[pn:trna nucleotidyltransferase] [gn:cca]
CONTIG490	11882677_f3_124	3408	9070	735	245	921	1.5(10)-92	Escherichia coli	b3059	[pn:hypothetical 22.2 kd protein in bacA-ttdA intergenic region]
CONTIG490	25525287_f3_129	3409	9071	1707	569	1859	6.0(10)-192	Bacillus subtilis	ureC	[pn:urease]
CONTIG490	5119193_f3_130	3410	9072	474	158	405	7.2(10)-38	Klebsiella aerogenes	P18317	urease accessory protein uree.
CONTIG490	23471885_f3_144	3411	9073	888	296	103	0.012	Rhodobacter sphaeroides	AJ000977	[de:rhodobacter sphaeroides dna for second chemotaxis operon andflanking genes.] [pn:transducer-like protein, tlpC] [gn:tlpC] [nt:tlpC shows weak homology to tlpA (formerly orf1) of]
CONTIG490	21692187_c1_148	3412	9074	1644	548	525	1.3(10)-50	Escherichia coli	b4355	[pn:methyl-accepting chemotaxis protein i] [gn:tsr]
CONTIG490	10553811_c1_150	3413	9075	825	275	991	5.7(10)-100	Escherichia coli	b3070	[pn:hypothetical protein]
CONTIG490	33867181_c1_172	3414	9076	390	130	555	9.1(10)-54	Escherichia coli	b3058	[pn:hypothetical 13.6 kd protein in bacA-ttdA intergenic region]
CONTIG490	4114702_c1_173	3415	9077	822	274	1144	3.5(10)-116	Escherichia coli	b3057	[pn:bacitracin resistance protein] [gn:baca]
CONTIG490	10969692_c2_196	3416	9078	1512	504	1862	2.8(10)-192	Escherichia coli	b3072	[pn:aerotaxis receptor protein] [gn:aer]
CONTIG490	26227307_c2_199	3417	9079	519	173	678	8.5(10)-67	Escherichia coli	b3068	[pn:hypothetical protein] [gn:ygif]
CONTIG490	24400916_c2_223	3418	9080	1323	441	1573	1.2(10)-161	Escherichia coli	b3054	[pn:hypothetical 48.4 kd protein in glne-cca intergenic region] [gn:ygif]
CONTIG490	36120650_c2_224	3419	9081	2877	959	3854	0	Escherichia coli	b3053	[pn:adenyl-transferase] [gn:gln]
CONTIG490	24276662_c3_255	3420	9082	1032	344	1552	2.1(10)-159	Escherichia coli	b3064	[pn:hypothetical 36.0 kd protein in ttdB-rpsu intergenic region]
CONTIG490	26578152_c3_282	3421	9083	1449	483	2299	1.3(10)-238	Escherichia coli	b3052	[pn:hypothetical protein]

CONTIG490	13671893_c3_284	3422	9084	666	222	995	2.2(10)-100	Escherichia coli	b3041	[pn:3,4-dihydroxy-2-butanone 4-phosphate synthase] [gn:ribb]
CONTIG490	4470033_c3_291	3423	9085	825	275	1230	2.7(10)-125	Escherichia coli	b3039	[pn:hypothetical 29.9 kd protein in tole-ribb intergenic region] [gn:ygld]
CONTIG491	2223507_fl_1	3424	9086	906	302	170	5.7(10)-11	Bacillus subtilis	ydeE	[pn:hypothetical protein]
CONTIG491	2056402_fl_9	3425	9087	1386	462	951	1.0(10)-95	Synechocystis sp.	S76228	[PN:hypothetical protein] [OR:Synechocystis sp.] [SR:PCC 6803,, PCC 6803] [SR:PCC 6803,,]
CONTIG491	25682967_fl_13	3426	9088	1092	364	322	1.8(10)-58	Synechocystis sp.	S77535	[PN:hypothetical protein] [OR:Synechocystis sp.] [SR:PCC 6803,, PCC 6803] [SR:PCC 6803,,]
CONTIG491	30728318_fl_22	3427	9089	681	227	729	3.2(10)-72	Escherichia coli	b1529	[pn:hypothetical protein in marr 5''' region] [gn:ydeb]
CONTIG491	1432191_fl_23	3428	9090	849	283	292	6.7(10)-26	Synechococcus PCC7942	U59236	or:synechococcus pcc7942 pn:unknown le:4337 re:>4953 di:direct nt:orf205
CONTIG491	13158461_fl_25	3429	9091	1428	476	1645	2.8(10)-169	Escherichia coli	b1525	[pn:hypothetical protein]
CONTIG491	24120381_fl_26	3430	9092	948	316	1338	9.8(10)-137	Escherichia coli	b1524	[pn:hypothetical protein] [gn:yneh]
CONTIG491	11069415_fl_27	3431	9093	987	329	225	1.7(10)-18	Escherichia coli	b0385	[pn:hypothetical 41.5 kd protein in psif-proc intergenic region] [gn:yaic]
CONTIG491	4379092_fl_28	3432	9094	1464	488	2225	9.9(10)-231	Escherichia coli	b1521	[pn:altronate oxidoreductase] [gn:uxab]
CONTIG491	32117968_fl_29	3433	9095	963	321	1268	2.6(10)-129	Escherichia coli	b1520	[pn:hypothetical protein]
CONTIG491	30494702_fl_32	3434	9096	1296	432	236	1.0(10)-17	Escherichia coli	b3911	[pn:cpxa] [gn:cpxa]
CONTIG491	29816950_fl_33	3435	9097	1326	442	210	1.1(10)-16	Bacillus subtilis	ykuV	[pn:hypothetical protein]
CONTIG491	4180191_f2_41	3436	9098	1974	658	3043	0	Escherichia coli	b1435	[pn:hypothetical protein]
CONTIG491	4305318_f2_44	3437	9099	420	140	106	7.7(10)-6	Mycobacterium tuberculosis	AL022002	[de:mycobacterium tuberculosis sequence v047.] [pn:hypothetical protein mtv047.09c] [gn:mtv047.09c] [nt:mtv047.09c,
CONTIG491	15023916_f2_45	3438	9100	1128	376	926	4.5(10)-93	Bacillus subtilis	yxjG	[pn:hypothetical protein] [gn:n15nr]



CONTIG491	16510187_f2_48	3439	9101	1614	538	412	1.3(10)-38	Methylophilus methylophilus	Y14964	[de:methylophilus methylophilus fmdd, fmde genes and partial fmfdgene.] [pn:putative transport protein] [gn:fmde]
CONTIG491	12145260_f2_52	3440	9102	681	227	723	1.3(10)-71	Escherichia coli	b1973	[pn:hypothetical protein]
CONTIG491	14554838_f2_60	3441	9103	825	275	155	3.7(10)-9	Escherichia coli	b1464	[pn:hypothetical 32.3 kd protein in rhse-narv intergenic region] [gn:ydde]
CONTIG491	21721965_f2_64	3442	9104	1653	551	485	2.3(10)-46	Escherichia coli	b1886	[pn:methyl-accepting chemotaxis protein ii] [gn:tar]
CONTIG491	4817693_f2_68	3443	9105	537	179	195	1.3(10)-15	Pseudomonas syringae	P16966	acetyltransferase (ec 2.3.1.-) (tabtoxin resistance protein).
CONTIG491	22479765_f2_72	3444	9106	453	151	262	1.0(10)-22	Acinetobacter calcoaceticus	AF009672	[PN:unknown] [DE:Acinetobacter calcoaceticus ADP1 vanillate demethylase region, vanillate demethylase (vanB) and vanillate demethylase (vanA) genes, complete cds.] [NT:putative acetyl transferase; ORF2] [LE:535] [RE:1002] [DI:complement]
CONTIG491	30118883_f2_75	3445	9107	792	264	503	3.0(10)-48	Escherichia coli	b3405	[pn:ompr] [gn:ompr]
CONTIG491	4429568_f2_80	3446	9108	1302	434	232	4.5(10)-17	Erwinia carotovora	JC1219	polygalacturonase (ec 3.2.1.15) precursor - erwinia carotovora this enzyme plays an important role in plant tissue maceration.
CONTIG491	5972143_f3_84	3447	9109	585	195	600	1.6(10)-58	Escherichia coli	b1434	[pn:hypothetical protein]
CONTIG491	34198965_f3_90	3448	9110	459	153	94	0.035	Caenorhabditis elegans	Z81518	[de:caenorhabditis elegans cosmid f28d9, complete sequence.] [pn:f28d9.a] [nt:protein predicted using genefinder; preliminary]
CONTIG491	992843_f3_96	3449	9111	810	270	319	9.4(10)-29	Escherichia coli	b3455	[pn:high-affinity branched-chain amino acid transport atp-binding protein livg] [gn:livg]
CONTIG491	5370443_f3_97	3450	9112	594	198	299	1.2(10)-26	Escherichia coli	b3454	[pn:high-affinity branched-chain amino acid transport atp-binding] [gn:livf]

CONTIG491	23984767_β_102	3451	9113	936	312	1014	2.1(10)-102	Escherichia coli	b1533	[pn:hypothetical 28.7 kd protein in marb-dcp intergenic region] [gn:yded]
CONTIG491	29558277_β_111	3452	9114	513	171	506	1.3(10)-48	Escherichia coli	b1523	[pn:hypothetical protein]
CONTIG491	31677037_β_115	3453	9115	1074	358	153	1.1(10)-8	Bacillus subtilis	yfhM	[pn:hypothetical protein]
CONTIG491	13697152_β_121	3454	9116	762	254	291	8.6(10)-26	Bacillus subtilis	fruR	[pn:transcriptional regulator]
CONTIG491	25484625_c1_125	3455	9117	669	223	496	1.6(10)-47	Escherichia coli	b2467	[pn:hypothetical 21.7 kd protein in tkb-marq intergenic region] [gn:yfh]
CONTIG491	22438307_c1_126	3456	9118	783	261	953	6.0(10)-96	Escherichia coli	b1519	[pn:hypothetical protein]
CONTIG491	24737775_c1_148	3457	9119	615	205	96	0.035	Microbacterium ammoniaphilum	X79027	or: microbacterium ammoniaphilum pn: unknown le: 3382 re: >4972 di: complement
CONTIG491	12600660_c1_150	3458	9120	912	304	1155	2.3(10)-117	Escherichia coli	b1526	[pn:hypothetical protein] [gn:yncj]
CONTIG491	21593738_c1_156	3459	9121	429	143	633	5.0(10)-62	Escherichia coli	b1531	[pn: multiple antibiotic resistance protein] [gn: mara]
CONTIG491	12986452_c1_173	3460	9122	1011	337	290	1.1(10)-25	Azorhizobium caulinodans	S52856	arac-like protein - azorhizobium caulinodans
CONTIG491	33603452_c1_176	3461	9123	237	79	292	6.7(10)-26	Escherichia coli	b1436	[pn:hypothetical protein]
CONTIG491	12140785_c1_180	3462	9124	1206	402	1035	1.3(10)-104	Escherichia coli	b1433	[pn:hypothetical protein]
CONTIG491	31442539_c2_184	3463	9125	476	158	264	6.2(10)-23	Escherichia coli	b3558	[pn: insertion element is150 hypothetical 33.3 kd protein] [gn: y15b]
CONTIG491	24785687_c2_197	3464	9126	600	200	113	2.5(10)-5	Bacillus subtilis	ydaF	[pn:hypothetical protein]
CONTIG491	10584683_c2_216	3465	9127	480	160	592	1.1(10)-57	Escherichia coli	b1530	[pn: multiple antibiotic resistance protein] [gn: marr]
CONTIG491	23626905_c2_219	3466	9128	1266	422	917	4.0(10)-92	Escherichia coli	b1534	[pn:hypothetical protein in marb-dcp intergenic region] [gn: ydef]
CONTIG491	35679180_c3_250	3467	9129	801	267	154	2.8(10)-11	Rhizobium sp.	S28675	hypothetical protein 5 - rhizobium sp. (strain ic3342)

CONTIG491	5265668_c3_274	3468	9130	1248	416	1474	3.7(10)-151	Escherichia coli	b1528	[pn:hypothetical protein in marr 5'' region] [gn:ydea]
CONTIG491	25897288_c3_276	3469	9131	234	78	142	5.2(10)-10	Escherichia coli	b1532	[pn:multiple antibiotic resistance protein marb] [gn:marb]
CONTIG491	6364437_c3_301	3470	9132	489	163	148	1.0(10)-9	Escherichia coli	b1433	[pn:hypothetical protein]
CONTIG492	16601011_fl_9	3471	9133	369	123	357	8.8(10)-33	Escherichia coli	b1273	[pn:hypothetical 9.4 kd protein in sobh-topa intergenic region]
CONTIG492	25554807_fl_16	3472	9134	1584	528	2099	2.2(10)-217	Escherichia coli	b1264	[pn:anthranilate synthase component i] [gn:trpe]
CONTIG492	31832017_fl_17	3473	9135	1272	424	1623	6.2(10)-167	Haemophilus influenzae	HI1431	[pn:tryptophan synthase, beta chain] [gn:trpb]
CONTIG492	29687656_fl_23	3474	9136	1050	350	926	4.5(10)-93	Escherichia coli	b1255	[pn:hypothetical protein in tonb-tpa intergenic region] [gn:ycic]
CONTIG492	2032531_fl_25	3475	9137	465	155	485	2.3(10)-46	Haemophilus influenzae	HI0827	[pn:p14 protein]
CONTIG492	46905_fl_43	3476	9138	2745	915	3902	0	Escherichia coli	b1241	[pn:alcohol/acetaldehyde dehydrogenase] [gn:adhe]
CONTIG492	25589537_f2_62	3477	9139	642	214	696	1.1(10)-68	Escherichia coli	b1270	[pn:i alamin adenosyltransferase] [gn:btur]
CONTIG492	26460037_f2_72	3478	9140	252	84	396	6.5(10)-37	Escherichia coli	b1930	[pn:hypothetical 8.6 kd protein in amya-flie intergenic region]
CONTIG492	23706650_f2_74	3479	9141	564	188	386	7.4(10)-36	Haemophilus influenzae	HI0826	[pn:sp]
CONTIG492	22833333_f2_76	3480	9142	2052	684	1755	6.2(10)-181	Escherichia coli	b0584	[pn:ferrienterobactin receptor precursor] [gn:fepa]
CONTIG492	21720266_f2_80	3481	9143	333	111	449	1.6(10)-42	Escherichia coli	b1251	[pn:hypothetical protein in kch-tonb intergenic region] [gn:ycii]
CONTIG492	32462652_f2_84	3482	9144	1512	504	2392	2.0(10)-248	Escherichia coli	U24201	or:escherichia coli pn:cardiolipin synthase gn:cls le:5896 re:7356 di:direct
CONTIG492	4350461_f2_85	3483	9145	342	114	555	9.1(10)-54	Escherichia coli	b1248	[pn:hypothetical protein]
CONTIG492	26282780_f3_101	3484	9146	654	218	960	1.1(10)-96	Escherichia coli	b1277	[pn:gtp cyclohydrolase ii] [gn:riba]
CONTIG492	6363568_f3_113	3485	9147	771	257	1039	4.7(10)-105	Escherichia coli	b1271	[pn:hypothetical oxidoreductase in btur-sobh intergenic region]
CONTIG492	35807292_f3_119	3486	9148	1653	551	2248	3.6(10)-233	Escherichia coli	b1263	[pn:anthranilate synthase component ii] [gn:trpd]

CONTIG492	15025302_f3_120	3487	9149	1362	454	1894	1.2(10)-195	Escherichia coli	b1262	[pn:indole-3-glycerol phosphate synthase] [gn:trpc]
CONTIG492	2117811_f3_121	3488	9150	861	287	693	2.2(10)-68	Haemophilus influenzae	H11432	[pn:tryptophan synthase alpha subunit] [gn:trpa]
CONTIG492	1448253_f3_122	3489	9151	1227	409	1588	3.1(10)-163	Escherichia coli	b1929	[pn:hypothetical 44.4 kd protein in amyala-flie intergenic region] [gn:yede]
CONTIG492	1298500_f3_124	3490	9152	435	145	435	4.7(10)-41	Escherichia coli	b0288	[pn:hypothetical protein]
CONTIG492	9819786_c1_153	3491	9153	351	117	330	6.4(10)-30	Escherichia coli	b1242	[pn:hypothetical protein in adhe-oppa intergenic region] [gn:yche]
CONTIG492	36022511_c1_156	3492	9154	954	318	1349	6.7(10)-138	Escherichia coli	b1244	[pn:oligopeptide transport system permease protein oppb] [gn:oppb]
CONTIG492	16198576_c1_157	3493	9155	921	307	1170	6.2(10)-119	Escherichia coli	b1245	[pn:oligopeptide transport system permease protein] [gn:oppc]
CONTIG492	6720317_c1_200	3494	9156	813	271	924	7.2(10)-93	Escherichia coli	b1278	[pn:phosphatidylglycerophosphatase b] [gn:pgpb]
CONTIG492	3909503_c2_206	3495	9157	471	157	562	1.7(10)-54	Escherichia coli	b1242	[pn:hypothetical protein in adhe-oppa intergenic region] [gn:yche]
CONTIG492	36219783_c2_208	3496	9158	1023	341	1588	3.1(10)-163	Escherichia coli	b1247	[pn:oligopeptide transport atp-binding protein oppf] [gn:oppf]
CONTIG492	495140_c2_209	3497	9159	864	288	175	5.5(10)-12	Bacillus subtilis	yugO	[pn:hypothetical protein]
CONTIG492	880092_c2_212	3498	9160	648	216	104	1.1(10)-5	Plasmid ColEI	J01566	or:plasmid col1 le:3943 re:4362 di:complement sr:plasmid col1 (clone: pew2762 and pmm1.) dna nt:entry exclusion protein 2 (exc2)
CONTIG492	23679576_c2_215	3499	9161	876	292	391	4.4(10)-63	Escherichia coli	b1252	[pn:tonb protein] [gn:tonb]
CONTIG492	25973287_c2_221	3500	9162	654	218	860	4.4(10)-86	Escherichia coli	b1256	[pn:hypothetical 22.9 kd protein in tonb-trpa intergenic region]
CONTIG492	34276692_c2_240	3501	9163	633	211	1025	1.3(10)-103	Escherichia coli	b1267	[pn:hypothetical 24.5 kd protein in trp1-btur intergenic region] [gn:ycio]
CONTIG492	26620277_c2_244	3502	9164	1062	354	1170	6.2(10)-119	Escherichia coli	b1272	[pn:possible protease] [gn:sohb]
CONTIG492	12694152_c2_247	3503	9165	2841	947	4175	0	Escherichia coli	b1276	[pn:aconitate hydratase 1] [gn:acna]
CONTIG492	3911432_c2_250	3504	9166	348	116	342	3.3(10)-31	Escherichia coli	b1279	[pn:hypothetical protein]

CONTIG492	3157830_c2_251	3505	9167	1089	363	1473	4.7(10)-151	Escherichia coli	b1280	[pn:hypothetical protein in pyrF 5' region] [gn:ycim]
CONTIG492	24273561_c3_254	3506	9168	1749	583	2422	1.3(10)-251	Escherichia coli	b1243	[pn:periplasmic oligopeptide-binding protein precursor] [gn:oppa]
CONTIG492	12750790_c3_257	3507	9169	1050	350	1607	3.1(10)-165	Escherichia coli	b1246	[pn:oligopeptide transport atp-binding protein oppd] [gn:oppd]
CONTIG492	15634788_c3_263	3508	9170	639	213	750	2.0(10)-74	Escherichia coli	b1798	[pn:hypothetical protein]
CONTIG492	167000_c3_286	3509	9171	951	317	1262	1.1(10)-128	Escherichia coli	b1266	[pn:hypothetical protein]
CONTIG492	4770943_c3_287	3510	9172	885	295	1355	1.5(10)-138	Escherichia coli	b1269	[pn:hypothetical 32.7 kd protein in trpL-bttr intergenic region] [gn:ycil]
CONTIG492	24667178_c3_291	3511	9173	2610	870	4117	0	Escherichia coli	b1274	[pn:dna topoisomerase i, omega protein i] [gn:topa]
CONTIG492	11180402_c3_292	3512	9174	990	330	1557	6.0(10)-160	Escherichia coli	b1275	[pn:cys regulon transcriptional activator] [gn:cysb]
CONTIG493	26306510_fl_4	3513	9175	192	64	213	1.6(10)-17	Escherichia coli	b1259	[pn:hypothetical protein in tonb-trpa intergenic region] [gn:ycig]
CONTIG493	26426302_fl_5	3514	9176	567	189	444	5.2(10)-42	Escherichia coli	U25418	or:escherichia coli gn:ycif le:4030 re:4530 di:direct
CONTIG493	1445785_fl_11	3515	9177	1653	551	870	3.7(10)-87	Streptomyces coelicolor	AJ001206	[de:streptomyces coelicolor a3(2), glycogen metabolism clusterii.] [pn:putative trehalose synthase]
CONTIG493	554561_fl_16	3516	9178	435	145	415	6.2(10)-39	Escherichia coli	b1536	[pn:hypothetical 14.1 kd protein in marb-dcp intergenic region]
CONTIG493	22844032_fl_22	3517	9179	573	191	629	1.3(10)-61	Escherichia coli	b2889	[pn:hypothetical protein]
CONTIG493	16174013_fl_32	3518	9180	534	178	848	8.1(10)-85	Escherichia coli	b2895	[pn:flavodoxin ii] [gn:fldb]
CONTIG493	11957058_fl_42	3519	9181	633	211	551	2.3(10)-53	Salmonella typhimurium	U75949	[de:salmonella typhimurium curved dna-binding protein (cbpa) gene andagp (agp) gene, partial cds; operon 1 containing suppressor for copper-sensitivity a (scsa) gene, "] [pn:suppressor for copper-sensitivity c] [gn:scsc]
CONTIG493	4882917_f2_53	3520	9182	903	301	345	1.6(10)-31	Bacillus subtilis	yjqC	[pn:hypothetical protein]

CONTIG493	22940910_f2_55	3521	9183	1077	359	910	2.2(10)-91	Pseudomonas aeruginosa	Y10528	[PN:cyanide insensitive terminal oxidase] [GN:cioB] [DE:P.aeruginosa cioA and cioB genes.] [LE:1746] [RE:2753] [DI:direct]
CONTIG493	20111291_f2_62	3522	9184	930	310	784	5.0(10)-78	Bacillus subtilis	yhxD	[pn:hypothetical protein]
CONTIG493	4797186_f2_93	3523	9185	1527	509	2399	3.6(10)-249	Escherichia coli	b2901	[pn:6-phospho-beta-glucosidase bglA] [gn:bglA]
CONTIG493	7300052_f2_97	3524	9186	606	202	322	4.5(10)-29	Haemophilus influenzae	H11115	[pn:thioredoxin]
CONTIG493	16854160_f3_101	3525	9187	921	307	1189	6.0(10)-121	Escherichia coli	b2845	[pn:hypothetical protein]
CONTIG493	23626681_f3_103	3526	9188	534	178	769	1.8(10)-76	Escherichia coli	U25423	or:escherichia coli gn:yoic le:4576 re:5082 di:direct
CONTIG493	16814418_f3_105	3527	9189	1452	484	1752	1.3(10)-180	Pseudomonas aeruginosa	Y10528	[PN:cyanide insensitive terminal oxidase] [GN:cioA] [DE:P.aeruginosa cioA and cioB genes.] [LE:276] [RE:1742] [DI:direct]
CONTIG493	26734683_f3_126	3528	9190	999	333	1451	1.0(10)-148	Escherichia coli	b2898	[pn:hypothetical protein] [gn:ygfZ]
CONTIG493	16525765_f3_129	3529	9191	762	254	181	7.5(10)-14	Bacillus subtilis	yfiA	[pn:hypothetical protein]
CONTIG493	17000052_f3_131	3530	9192	375	125	243	1.1(10)-20	Salmonella typhimurium	U75949	[de:salmonella typhimurium curved dna-binding protein (cbpa) gene andagp (agp) gene, partial cds; operon 1 containing suppressor forcopper-sensitivity a (scsa) gene;"] [pn:suppressor for copper-sensitivity a] [gn:scsa]
CONTIG493	13681500_f3_132	3531	9193	2013	671	1246	6.9(10)-173	Salmonella typhimurium	U75949	[de:salmonella typhimurium curved dna-binding protein (cbpa) gene andagp (agp) gene, partial cds; operon 1 containing suppressor forcopper-sensitivity a (scsa) gene;"] [pn:suppressor for copper-sensitivity b] [gn:scsb]

CONTIG493	16148542_c1_140	3532	9194	783	261	1016	1.3(10)-102	Escherichia coli	b2902	[pn:hypothetical oxidoreductase] [gn:ygff]
CONTIG493	5860055_c1_148	3533	9195	687	229	885	9.9(10)-89	Escherichia coli	b2899	[pn:hypothetical protein]
CONTIG493	3260282_c1_154	3534	9196	873	291	1007	1.2(10)-101	Escherichia coli	b2893	[pn:disulfide interchange protein, precursor] [gn:dsbc]
CONTIG493	910312_c1_156	3535	9197	1077	359	1581	1.7(10)-162	Escherichia coli	b2891	[pn:peptide chain release factor 2] [gn:prfb]
CONTIG493	4488443_c1_157	3536	9198	1527	509	2300	1.1(10)-238	Escherichia coli	b2890	[pn:lysyl tRNA synthetase] [gn:lyss]
CONTIG493	6813751_c1_164	3537	9199	636	212	392	1.7(10)-36	Escherichia coli	b0957	[pn:outer membrane protein a] [gn:ompa]
CONTIG493	10320330_c1_169	3538	9200	1035	345	280	1.3(10)-24	Bacillus subtilis	yxnA	[pn:hypothetical protein]
CONTIG493	16971016_c1_180	3539	9201	297	99	343	2.7(10)-31	Escherichia coli	b4126	[pn:hypothetical 8.6 kd protein in dcub-lysu intergenic region]
CONTIG493	22464665_c2_198	3540	9202	282	94	431	1.3(10)-40	Escherichia coli	b2897	[pn:hypothetical protein] [gn:ygy]
CONTIG493	21519790_c2_203	3541	9203	771	257	914	8.3(10)-92	Escherichia coli	b2865	[pn:hypothetical protein]
CONTIG493	23679591_c2_206	3542	9204	702	234	354	1.8(10)-32	Escherichia coli	b0897	[pn:hypothetical 23.1 kd protein in dmse 3'''region] [gn:ycac]
CONTIG493	4103838_c2_217	3543	9205	456	152	142	5.2(10)-10	Bacillus subtilis	ydrR	[pn:hypothetical protein]
CONTIG493	26597818_c2_226	3544	9206	279	93	408	3.5(10)-38	Escherichia coli	b4127	[pn:hypothetical 10.5 kd protein in dcub-lysu intergenic region]
CONTIG493	30365655_c3_228	3545	9207	2958	986	4326	0	Escherichia coli	b2903	[pn:glycine dehydrogenase] [gn:gcvp]
CONTIG493	35726525_c3_244	3546	9208	432	144	478	1.3(10)-45	Escherichia coli	b2896	[pn:hypothetical protein]
CONTIG493	29507090_c3_245	3547	9209	927	309	1341	4.7(10)-137	Escherichia coli	b2894	[pn:site-specific integrase/recombinase, with xerc] [gn:xerd]
CONTIG493	6500302_c3_247	3548	9210	1743	581	2600	1.8(10)-270	Escherichia coli	b2892	[pn:single-stranded dna-specific exonuclease] [gn:recj]
CONTIG493	23476500_c3_252	3549	9211	1260	420	197	7.5(10)-15	Escherichia coli	b2344	[pn:long-chain fatty acid transport protein precursor] [gn:fadl]
CONTIG493	36141327_c3_253	3550	9212	648	216	209	4.2(10)-17	Escherichia coli	b0375	[pn:hypothetical protein] [gn:yaiv]

CONTIG493	12110131_c3_254	3551	9213	2034	678	1566	6.7(10)-161	Escherichia coli	b4083	[pn:hypothetical 73.7 kd protein in fdhF-phnP intergenic region]
CONTIG493	807137_c3_257	3552	9214	585	195	426	4.2(10)-40	Escherichia coli	b1537	[pn:hypothetical 18.3 kd protein in marB-dcp intergenic region]
CONTIG494	14860206_f1_2	3553	9215	1092	364	1449	1.7(10)-148	Escherichia coli	b0888	[pn:thioredoxin reductase] [gn:trxb]
CONTIG494	14484500_f1_3	3554	9216	1797	599	2423	1.0(10)-251	Escherichia coli	b0887	[pn:transport atp-binding protein cydd] [gn:cydd]
CONTIG494	34376252_f1_26	3555	9217	819	273	679	6.7(10)-67	Escherichia coli	b0875	[pn:aquaporin z] [gn:aqpz]
CONTIG494	32659515_f1_27	3556	9218	942	314	1044	1.3(10)-105	Escherichia coli	b0874	[pn:hypothetical protein]
CONTIG494	115887_f1_28	3557	9219	1677	559	2665	2.2(10)-277	Escherichia coli	b0873	[pn:hypothetical protein]
CONTIG494	15745303_f1_65	3558	9220	744	248	1094	7.0(10)-111	Escherichia coli	b0864	[pn:arginine transport atp-binding protein artp] [gn:artp]
CONTIG494	26828515_f1_66	3559	9221	750	250	1117	2.6(10)-113	Escherichia coli	b0863	[pn:arginine-binding periplasmic protein 1 precursor] [gn:arti]
CONTIG494	33714207_f1_68	3560	9222	684	228	975	2.8(10)-98	Escherichia coli	b0861	[pn:arginine transport system permease protein artm] [gn:artm]
CONTIG494	26363577_f2_79	3561	9223	219	73	362	2.6(10)-33	Escherichia coli	b0884	[pn:initiation factor if-1] [gn:ifia]
CONTIG494	14160432_f2_84	3562	9224	288	96	364	1.6(10)-33	Escherichia coli	b0880	[pn:cold shock-like protein espd] [gn:cspd]
CONTIG494	4114637_f2_89	3563	9225	990	330	743	1.1(10)-73	Escherichia coli	b0877	[pn:hypothetical protein]
CONTIG494	4415901_f2_95	3564	9226	972	324	1441	1.2(10)-147	Escherichia coli	b0872	[pn:hypothetical protein]
CONTIG494	4423318_f2_99	3565	9227	1011	337	1468	1.6(10)-150	Escherichia coli	b0870	[pn:hypothetical protein] [gn:ybjv]
CONTIG494	14642176_f2_102	3566	9228	1110	370	1597	3.5(10)-164	Escherichia coli	b0868	[pn:hypothetical protein]
CONTIG494	12994811_f2_135	3567	9229	696	232	1090	1.8(10)-110	Escherichia coli	b0862	[pn:arginine transport system permease protein artq] [gn:artq]
CONTIG494	2995933_f3_140	3568	9230	1776	592	2045	1.2(10)-211	Escherichia coli	b0886	[pn:transport atp-binding protein cyde] [gn:cyde]
CONTIG494	5865882_f3_141	3569	9231	750	250	1032	2.6(10)-104	Escherichia coli	b0885	[pn:leucyl/phenylalanyl-tRNA--protein transferase] [gn:aat]



CONTIG494	15117842_β_159	3570	9232	1734	578	2699	5.9(10)-281	Escherichia coli	b0871	[pn:cytochrome] [gn:poxb]
CONTIG494	33875912_β_161	3571	9233	1476	492	2080	2.2(10)-215	Escherichia coli	b0869	[pn:hypothetical protein]
CONTIG494	36597081_β_183	3572	9234	1818	606	393	2.8(10)-36	Escherichia coli	b3323	[pn:putative general secretion pathway protein b] [gn:yhed]
CONTIG494	23631550_β_186	3573	9235	540	180	729	3.2(10)-72	Escherichia coli	b0865	[pn:hypothetical protein]
CONTIG494	13150827_c1_199	3574	9236	675	225	177	1.0(10)-13	Escherichia coli	b0375	[pn:hypothetical protein] [gn:yaiv]
CONTIG494	15761437_c1_206	3575	9237	1293	431	847	1.0(10)-84	Escherichia coli	b3327	[pn:putative general secretion pathway protein f] [gn:hoff]
CONTIG494	134652_c1_207	3576	9238	453	151	474	3.5(10)-45	Escherichia coli	b3328	[pn:putative general secretion pathway protein g precursor] [gn:hofg]
CONTIG494	16838437_c1_208	3577	9239	387	129	144	3.2(10)-10	Escherichia coli	b3330	[pn:putative general secretion pathway protein i precursor] [gn:yheh]
CONTIG494	32632827_c1_212	3578	9240	486	162	102	0.00013	Escherichia coli	b3334	[pn:putative general secretion pathway protein m] [gn:pshm]
CONTIG494	26735627_c1_213	3579	9241	2721	907	2118	7.2(10)-248	Aeromonas caviae	U09139	or:aeromonas caviae pn:chitinase protein precursor le:154 re:2751 di:direct
CONTIG494	30664092_c1_230	3580	9242	1689	563	2468	1.8(10)-256	Escherichia coli	b0876	[pn:hypothetical protein]
CONTIG494	16975466_c1_248	3581	9243	210	70	172	3.5(10)-13	Escherichia coli	b0889	[pn:leucine-responsive regulatory protein] [gn:lrp]
CONTIG494	32055135_c2_255	3582	9244	909	303	1016	1.3(10)-102	Escherichia coli	b0867	[pn:hypothetical protein]
CONTIG494	1447187_c2_263	3583	9245	777	259	155	1.5(10)-9	Escherichia coli	b3324	[pn:putative general secretion pathway protein c] [gn:yhec]
CONTIG494	32694192_c2_265	3584	9246	1500	500	1407	4.7(10)-144	Escherichia coli	b3326	[pn:type ii traffic] [gn:yheg]
CONTIG494	36222885_c2_268	3585	9247	1116	372	285	3.7(10)-25	Escherichia coli	b3332	[pn:putative general secretion pathway protein k] [gn:yhej]
CONTIG494	24641037_c2_269	3586	9248	1140	380	307	1.7(10)-27	Escherichia coli	b3333	[pn:putative general secretion pathway protein l] [gn:yhek]

CONTIG494	24726625_c2_276_3587	9249	1845	615	299	6.2(10)-26	Oryza sativa	L37289	or:oryza sativa pn:chitinase ec:3.2.1.14 le:43 re:1044 di:direct sr:oryza sativa (strain ir36) seedling etiolated leaf odna to mrna
CONTIG494	21767817_c2_299_3588	9250	1980	660	2714	1.5(10)-282	Escherichia coli	b0879	[pn:hypothetical protein]
CONTIG494	26449052_c3_323_3589	9251	339	113	493	3.3(10)-47	Escherichia coli	b0866	[pn:hypothetical protein]
CONTIG494	24015932_c3_328_3590	9252	1944	648	1216	8.3(10)-124	Escherichia coli	b3325	[pn:putative general secretion pathway protein d precursor] [gn:yhef]
CONTIG494	4899063_c3_332_3591	9253	498	166	158	1.1(10)-11	Escherichia coli	b3329	[pn:putative general secretion pathway protein h precursor] [gn:hofh]
CONTIG494	35678963_c3_334_3592	9254	708	236	142	5.2(10)-10	Escherichia coli	b3331	[pn:putative general secretion pathway protein j precursor] [gn:yhei]
CONTIG494	32594556_c3_336_3593	9255	849	283	366	9.8(10)-34	Escherichia coli	b2972	[pn:hypothetical protein]
CONTIG494	21672156_c3_337_3594	9256	537	179	208	5.4(10)-17	Escherichia coli	b2854	[pn:hypothetical protein]
CONTIG494	2914181_c3_353_3595	9257	1140	380	1393	1.5(10)-142	Escherichia coli	b0878	[pn:hypothetical protein]
CONTIG494	32507211_c3_355_3596	9258	393	131	528	6.7(10)-51	Escherichia coli	b0881	[pn:hypothetical protein] [gn:yjia]
CONTIG494	975293_c3_356_3597	9259	2304	768	3480	0	Escherichia coli	b0882	[pn:atp-dependent clp protease atp-binding subunit clpa] [gn:clpa]
CONTIG495	11724086_fl_17_3598	9260	618	206	772	9.3(10)-77	Escherichia coli	b3465	[pn:hypothetical 21.7 kd protein in ftsy-nika intergenic region]
CONTIG495	20976555_fl_25_3599	9261	681	227	999	8.1(10)-101	Escherichia coli	b3471	[pn:hypothetical 25.3 kd protein in ftsy-nika intergenic region]
CONTIG495	13095332_fl_26_3600	9262	600	200	874	1.3(10)-87	Escherichia coli	b3472	[pn:hypothetical protein]
CONTIG495	32300666_fl_27_3601	9263	1113	371	1421	1.6(10)-145	Escherichia coli	b3474	[pn:hypothetical 38.5 kd protein in ftsy-nika intergenic region] [gn:yhht]
CONTIG495	5195317_fl_32_3602	9264	1356	452	120	0.0032	Bacillus subtilis	srfAB	[pn:surfactin synthetase] [gn:coml]

CONTIG495	22369816_f1_34	3603	9265	1728	576	173	2.6(10)-12	Methanobacterium thermoautotrophicum	MTH136	[pn:dolichyl-phosphate mannose synthase]
CONTIG495	14869217_f1_35	3604	9266	426	142	119	1.5(10)-7	Bacillus subtilis	ynp	[pn:hypothetical protein]
CONTIG495	16912907_f1_38	3605	9267	1251	417	354	1.8(10)-32	Escherichia coli	b1095	[pn:3-oxoacyl-acyl-carrier-protein synthase ii] [gn:fabf]
CONTIG495	10626391_f1_39	3606	9268	330	110	96	0.00289	Homo sapiens	AF004884	[PN:neuronal calcium channel alpha 1A subunit] [SR:human] [DE:Human neuronal calcium channel alpha 1A subunit isoform A-1mRNA, complete cds.] [LE:237] [RE:7769] [DI:direct]
CONTIG495	15808466_f1_40	3607	9269	1242	414	624	4.5(10)-61	Bacillus subtilis	yjaY	[pn:hypothetical protein]
CONTIG495	14229752_f2_48	3608	9270	432	144	498	1.0(10)-47	Escherichia coli	b3459	[pn:hypothetical 14.5 kd protein in livk-livj intergenic region] [gn:yhhk]
CONTIG495	24619625_f2_58	3609	9271	408	136	341	4.4(10)-31	Escherichia coli	b3466	[pn:hypothetical 10.3 kd protein in ftsy-nika intergenic region] [gn:yhhj]
CONTIG495	10644062_f2_60	3610	9272	753	251	833	3.2(10)-83	Escherichia coli	b3468	[pn:hypothetical protein] [gn:yhhm]
CONTIG495	16269817_f2_71	3611	9273	921	307	122	1.3(10)-5	Helicobacter pylori	HP1348	[pn:1-acyl-glycerol-3-phosphate acyltransferase] [gn:plsc]
CONTIG495	11064191_f2_72	3612	9274	330	110	94	6.5(10)-5	Helicobacter pylori	HP0559	[pn:acyl carrier protein] [gn:acpp]
CONTIG495	31347015_f2_82	3613	9275	813	271	400	2.3(10)-37	Bacillus subtilis	fabG	[pn:3-oxoacyl-acyl-carrier protein reductase] [gn:yipf]
CONTIG495	4884707_f2_86	3614	9276	798	266	552	1.8(10)-53	Bacillus subtilis	yvaG	[pn:hypothetical protein]
CONTIG495	2071916_f3_103	3615	9277	2244	748	2482	5.7(10)-258	Escherichia coli	b3469	[pn:hypothetical protein] [gn:yhho]
CONTIG495	447686_f3_113	3616	9278	291	97	92	0.00011	Leuothrix mucor	P80920	
CONTIG495	13016416_f3_120	3617	9279	384	128	95	0.00034	Streptomyces fradiae	P20186	hypothetical 35.5 kd protein in transposon tn4556.

CONTIG495	11924157_f3_129	3618	9280	543	181	107	2.6(10)-5	Brucella abortus	U51683	or:brucella abortus pn:fabz gn:fabz ec:4.2.1.- le:6377 re:6844 di:direct sr:brucella abortus strain=s2308 nt:similar to swiss-prot accession number p21774
CONTIG495	6540832_f3_133	3619	9281	609	203	705	1.2(10)-69	Escherichia coli	b3475	[pn:hypothetical 21.8 kd protein in ftsy-nika intergenic region]
CONTIG495	30104052_f3_135	3620	9282	927	309	225	8.5(10)-19	Escherichia coli	b0504	[pn:hypothetical protein] [gn:ybbs]
CONTIG495	1307918_c1_142	3621	9283	318	106	229	2.2(10)-18	Rhizobium sp.	P50360	hypothetical 29.3 kd protein in region 2 of sym plasmid (no1265).
CONTIG495	20984450_c1_143	3622	9284	468	156	432	9.9(10)-41	Escherichia coli	S70162	,
CONTIG495	29329043_c1_187	3623	9285	699	233	1086	4.9(10)-110	Escherichia coli	b3463	[pn:cell division atp-binding protein ftsl] [gn:ftsle]
CONTIG495	29713291_c1_190	3624	9286	1344	448	1578	3.6(10)-162	Escherichia coli	b1302	[pn:gaba-aminotransferase] [gn:goag]
CONTIG495	4957587_c1_192	3625	9287	336	112	387	5.7(10)-36	Escherichia coli	b3458	[pn:leucine-specific binding protein precursor] [gn:livk]
CONTIG495	10282891_c2_194	3626	9288	1098	366	1388	4.9(10)-142	Plasmid R478	L38824	or:plasmid r478 gn:terc le:2277 re:3317 di:direct sr:plasmid r478 dna nt:putative
CONTIG495	5189842_c2_202	3627	9289	546	182	478	1.3(10)-45	Bacillus subtilis	padC	[pn:ferulate decarboxylase] [gn:yveh]
CONTIG495	14504207_c2_216	3628	9290	246	82	90	0.00017	Haemophilus influenzae	H11355	[pn:hypothetical protein]
CONTIG495	24271091_c2_227	3629	9291	285	95	391	2.2(10)-36	Escherichia coli	b3470	[pn:hypothetical 9.1 kd protein in ftsy-nika intergenic region]
CONTIG495	21568791_c2_228	3630	9292	1002	334	395	8.3(10)-37	Escherichia coli	b3579	[pn:hypothetical 36.0 kd protein in avta-selb intergenic region] [gn:yiao]
CONTIG495	32086658_c2_229	3631	9293	522	174	114	5.0(10)-7	Haemophilus influenzae	H11030	[pn:sp]
CONTIG495	16838955_c2_240	3632	9294	1467	489	1423	1.3(10)-152	Escherichia coli	b3464	[pn:cell division protein ftsy] [gn:ftsyl]
CONTIG495	4587833_c2_241	3633	9295	1083	361	1361	3.6(10)-139	Escherichia coli	b3462	[pn:cell division protein ftsx] [gn:ftsxl]
CONTIG495	15837807_c3_277	3634	9296	1290	430	548	5.0(10)-53	Haemophilus influenzae	H11029	[pn:sp]

CONTIG495	9767340_c3_283	3635	9297	369	123	211	2.6(10)-17	Escherichia coli	b3467	[pn:hypothetical 13.5 kd protein in ftzy-nika intergenic region] [gn:yhbm]
CONTIG495	26363452_c3_287	3636	9298	858	286	1211	2.7(10)-123	Escherichia coli	b3461	[pn:rna polymerase sigma-32 subunit] [gn:rpoh]
CONTIG495	3257762_c3_291	3637	9299	1164	388	1772	1.0(10)-182	Escherichia coli	b3460	[pn:leu/ile/val-binding protein precursor] [gn:livj]
CONTIG496	4304068_fl_8	3638	9300	2394	798	680	5.2(10)-67	Porphyromonas gingivalis	P49008	beta-hexosaminidase precursor (ec 3.2.1.52) (n-acetyl-beta-glucosaminidase) (beta-glcnaase) (beta-n-acetylhexosaminidase) (beta-nahase).
CONTIG496	29589130_fl_21	3639	9301	468	156	670	6.0(10)-66	Escherichia coli	b4252	[pn:hypothetical 17.3 kd protein in pyl-argi intergenic region] [gn:yjgk]
CONTIG496	24353408_fl_32	3640	9302	429	143	358	6.9(10)-33	Escherichia coli	b4255	[pn:hypothetical 15.6 kd protein in argi-vals intergenic region] [gn:yjgd]
CONTIG496	30369653_fl_49	3641	9303	3330	1110	1018	7.9(10)-103	Escherichia coli	b4308	[pn:hypothetical 38.0 kd protein in feci-fimb intergenic region] [gn:yjhr]
CONTIG496	23609515_fl_50	3642	9304	294	98	419	2.3(10)-39	Enterobacter agglomerans	A38965	hypothetical protein a (insertion sequence is 1222) -enterobacter agglomerans
CONTIG496	3381711_f2_55	3643	9305	2775	925	3895	0	Escherichia coli	b4242	[pn:mg2+ transport atpase, p-type 1] [gn:mgta]
CONTIG496	25522918_f2_76	3644	9306	2025	675	3216	0	Escherichia coli	b0269	[pn:hypothetical 69.4 kd protein in per-argf intergenic region] [gn:yagf]
CONTIG496	30332633_f2_81	3645	9307	1725	575	2884	1.5(10)-300	Escherichia coli	b0271	[pn:hypothetical protein] [gn:yagh]
CONTIG496	34480032_f2_86	3646	9308	936	312	1124	4.5(10)-114	Salmonella typhimurium	Q08015	tma-(ms[2]io[6]a)-hydroxylase (ec 1.1.1.1)
CONTIG496	3984643_f3_102	3647	9309	489	163	183	2.3(10)-14	Vibrio cholerae	S81006	or: vibrio cholerae pn:hcp gn:hcp le:690 re:1208 di:direct sr:vibrio cholerae o17 nt:28 kda secreted hydrophilic protein; this sequence

CONTIG496	954637_f3_104	3648	9310	222	74	100	1.5(10)-5	Escherichia coli	D21157	[PN:unknown] [SR:Escherichia coli (sub_strain W3110, strain K-12) (library: librar] [DE:E.coli gene for unknown product, partial cds.] [NT:the coding frame was determined by the Lac fusion] [LE:57] [RE:192] [DI:direct]
CONTIG496	4470443_f3_123	3649	9311	918	306	1207	7.4(10)-123	Escherichia coli	b0268	[pn:hypothetical 33.3 kd protein in perr-argf intergenic region]
CONTIG496	2849025_f3_126	3650	9312	1413	471	2155	2.6(10)-223	Escherichia coli	b0270	[pn:hypothetical 50.6 kd protein in perr-argf intergenic region]
CONTIG496	9896067_f3_151	3651	9313	186	62	196	1.0(10)-15	Enterobacter agglomerans	B38965	hypothetical protein b (insertion sequence is1222) -enterobacter agglomerans
CONTIG496	26370678_c1_160	3652	9314	1488	496	2397	5.9(10)-249	Escherichia coli	I41293	ecoe type i restriction modification enzyme m subunit -escherichia coli
CONTIG496	29863508_c1_183	3653	9315	951	317	739	2.8(10)-73	Haemophilus influenzae	HI0595	[pn:carbamate kinase] [gn:arcc]
CONTIG496	11214091_c1_185	3654	9316	1551	517	756	4.0(10)-83	Haemophilus influenzae	HI0594	[pn:hypothetical protein]
CONTIG496	33603340_c1_195	3655	9317	300	100	94	6.5(10)-5	Saccharomyces cerevisiae	X85757	or:saccharomyces cerevisiae pn:unknown gn:internal orf g1669 le:6964 re:7365 di:direct sr:baker's yeast
CONTIG496	6251280_c2_204	3656	9318	2466	822	3981	0	Escherichia coli	I41292	ecoe type i restriction-modification enzyme r subunit -escherichia coli
CONTIG496	2599015_c2_216	3657	9319	915	305	1072	1.5(10)-108	Escherichia coli	b0272	[pn:hypothetical transcriptional regulator in perr-argf intergenic region] [gn:yagi]
CONTIG496	31328187_c2_226	3658	9320	1014	338	759	2.2(10)-75	Escherichia coli	U70214	or:escherichia coli le:112561 re:113301 di:complement nt:similar to e. coli yjhh
CONTIG496	21954407_c2_234	3659	9321	993	331	1409	2.8(10)-144	Escherichia coli	b4245	[pn:aspartate carbomoyltransferase catalytic subunit] [gn:pyrb]
CONTIG496	14964218_c2_235	3660	9322	474	158	679	6.7(10)-67	Escherichia coli	b4244	[pn:aspartate carbomoyltransferase regulatory subunit] [gn:pyri]
CONTIG496	32547931_c3_258	3661	9323	1794	598	1274	5.9(10)-130	Citrobacter freundii	X17591	or:citrobacter freundii pn:hsds polypeptide, part of cfra family gn:hsds le:234 re:1970 di:direct

CONTIG496	30657775_c3_266	3662	9324	1038	346	1701	3.2(10)-175	Escherichia coli	b0273	[pn:ornithine carbamoyltransferase chain f] [gn:argf]
CONTIG496	24347205_c3_275	3663	9325	213	71	117	6.2(10)-7	Escherichia coli	U70214	or:escherichia coli ie:112561 re:113301 di:complement nt:similar to e. coli yjhh
CONTIG496	4469557_c3_276	3664	9326	1275	425	912	1.3(10)-91	Pseudomonas aeruginosa	P13981	arginine deiminase (ec 3.5.3.6).
CONTIG496	24406311_c3_279	3665	9327	1026	342	1230	2.7(10)-125	Escherichia coli	b4254	[pn:ornithine carbamoyltransferase] [gn:argi]
CONTIG496	10329756_c3_283	3666	9328	708	236	206	8.8(10)-17	Escherichia coli	b3237	[pn:arginine repressor] [gn:argr]
CONTIG496	9955041_c3_285	3667	9329	408	136	642	5.5(10)-63	Escherichia coli	b4243	[pn:13.5 kd protein in mgta-pyri intergenic region] [gn:yjgf]
CONTIG496	10179702_c3_286	3668	9330	1449	483	377	6.7(10)-35	Streptomyces olivaceoviridis	S55000	alpha-chitin binding protein precursor - streptomycetolivaceoviridis (strain
CONTIG496	31406300_c3_297	3669	9331	978	326	1381	2.7(10)-141	Escherichia coli	b4241	[pn:trehalose operon repressor] [gn:trtr]
CONTIG497	26736576_fl_1	3670	9332	1272	424	1955	4.0(10)-202	Escherichia coli	b2551	[pn:serine hydroxymethyltransferase] [gn:glya]
CONTIG497	25969791_fl_2	3671	9333	1206	402	1637	2.0(10)-168	Escherichia coli	b2536	[pn:hypothetical protein] [gn:yfhs]
CONTIG497	4181268_fl_15	3672	9334	333	111	452	7.5(10)-43	Escherichia coli	b2528	[pn:hypothetical protein in hscsa 5'''region] [gn:yfhi]
CONTIG497	34235307_fl_17	3673	9335	1896	632	2657	1.7(10)-276	Escherichia coli	b2526	[pn:heat shock protein hscsa] [gn:hscsa]
CONTIG497	12397806_fl_27	3674	9336	4983	1661	6480	0	Escherichia coli	b2520	[pn:hypothetical protein]
CONTIG497	11953427_fl_43	3675	9337	535	179	637	1.8(10)-62	Escherichia coli	b2512	[pn:hypothetical protein]
CONTIG497	16266057_fl_47	3676	9338	1008	336	406	5.7(10)-38	Haemophilus influenzae	HI1248	[pn:hypothetical protein]
CONTIG497	12219836_fl_50	3677	9339	537	179	789	1.5(10)-78	Escherichia coli	b2527	[pn:hypothetical 20.1 kd protein in hscsa 5'''region] [gn:yfhe]
CONTIG497	11988587_fl_52	3678	9340	363	121	539	4.5(10)-52	Escherichia coli	b2525	[pn:ferredoxin, 2fe-2s] [gn:fdx]
CONTIG497	31650080_fl_70	3679	9341	2343	781	3091	0	Escherichia coli	b2519	[pn:hypothetical protein] [gn:pbpc]

CONTIG497	21759667_f2_72	3680	9342	633	211	585	6.0(10)-57	Escherichia coli	b0895	[pn:anaerobic dimethyl sulfoxide reductase chain b] [gn:dmsb]
CONTIG497	15751633_f2_74	3681	9343	912	304	126	1.5(10)-7	Methanobacterium thermoautotrophicum	MTH1241	[pn:polyferredoxin]
CONTIG497	4805165_f2_76	3682	9344	1170	390	1691	3.7(10)-174	Escherichia coli	b2515	[pn:gcpe protein] [gn:gcpe]
CONTIG497	879700_f2_77	3683	9345	1323	441	2088	3.2(10)-216	Escherichia coli	b2514	[pn:histidyl-trna synthetase] [gn:hiss]
CONTIG497	22479766_f3_82	3684	9346	651	217	356	1.1(10)-32	Haemophilus influenzae	HI1249	[pn:hypothetical protein precursor]
CONTIG497	31464586_f3_85	3685	9347	807	269	1032	2.6(10)-104	Escherichia coli	b2532	[pn:hypothetical protein]
CONTIG497	3009663_f3_86	3686	9348	495	165	640	9.0(10)-63	Escherichia coli	b2531	[pn:hypothetical protein]
CONTIG497	4956303_f3_87	3687	9349	1320	440	2020	5.2(10)-209	Escherichia coli	b2530	[pn:hypothetical protein]
CONTIG497	23616080_f3_88	3688	9350	411	137	642	5.5(10)-63	Escherichia coli	b2529	[pn:hypothetical protein]
CONTIG497	5275331_f3_92	3689	9351	264	88	346	1.3(10)-31	Escherichia coli	b2524	[pn:hypothetical 7.7 kd protein in fdx 3'''region] [gn:yfhj]
CONTIG497	34022075_f3_93	3690	9352	1308	436	1833	3.3(10)-189	Escherichia coli	b2523	[pn:hypothetical protein in fdx 3'''region] [gn:yfhi]
CONTIG497	9961718_f3_94	3691	9353	804	268	1144	3.5(10)-116	Escherichia coli	b2522	[pn:sseb protein] [gn:sseb]
CONTIG497	34492187_f3_104	3692	9354	2418	806	1485	2.6(10)-152	Escherichia coli	b1588	[pn:hypothetical protein]
CONTIG497	4557805_f3_106	3693	9355	819	273	133	7.0(10)-7	Escherichia coli	b1590	[pn:hypothetical protein]
CONTIG497	3259683_f3_108	3694	9356	516	172	696	1.1(10)-68	Escherichia coli	b2518	[pn:nucleoside diphosphate kinase] [gn:ndk]
CONTIG497	30267625_f3_109	3695	9357	1311	437	1854	2.0(10)-191	Escherichia coli	b2517	[pn:hypothetical 43.1 kd protein in ndk-gcpe intergenic region]
CONTIG497	13869066_f3_110	3696	9358	1011	337	1100	1.6(10)-111	Escherichia coli	b2516	[pn:hypothetical 36.2 kd protein in ndk-gcpe intergenic region]
CONTIG497	16839666_f3_113	3697	9359	678	226	695	1.3(10)-68	Escherichia coli	b2513	[pn:hypothetical protein]



CONTIG497	24424167_c1_147	3698	9360	951	317	1162	4.4(10)-118	Escherichia coli	b2521	[pn:putative thiosulfate sulfurtransferase] [gn:ssea]
CONTIG497	31879212_c1_163	3699	9361	1353	451	1344	2.2(10)-137	Escherichia coli	b2535	[pn:stationary phase inducible protein csie] [gn:csie]
CONTIG497	10052078_c2_204	3700	9362	1572	524	691	3.6(10)-68	Escherichia coli	b1621	[pn:pts system, maltose and glucose-specific ii abc component] [gn:malx]
CONTIG497	24245462_c2_205	3701	9363	1380	460	1013	2.7(10)-102	Bacillus subtilis	glvA	[pn:6-phospho-alpha-glucosidase] [gn:glvg]
CONTIG497	5180338_c2_223	3702	9364	813	271	1258	2.8(10)-128	Escherichia coli	b2533	[pn:extragenic suppressor protein subb] [gn:subb]
CONTIG497	13705037_c2_229	3703	9365	468	156	583	9.9(10)-57	Escherichia coli	b2543	[pn:hypothetical protein]
CONTIG497	4896032_c3_266	3704	9366	783	261	137	3.0(10)-7	Bacillus subtilis	ybbH	[pn:hypothetical protein]
CONTIG498	5947212_fl_1	3705	9367	447	149	651	6.2(10)-64	Escherichia coli	b1778	[pn:hypothetical protein] [gn:yeaa]
CONTIG498	16228431_fl_6	3706	9368	555	185	729	3.2(10)-72	Escherichia coli	b1765	[pn:hypothetical 20.1 kd protein in seld-sppa intergenic region]
CONTIG498	6336575_fl_8	3707	9369	1926	642	2996	0	Escherichia coli	b1763	[pn:dna topoisomerase iii] [gn:topb]
CONTIG498	24740925_fl_14	3708	9370	636	212	479	1.0(10)-45	Escherichia coli	b1753	[pn:hypothetical protein]
CONTIG498	19535307_fl_22	3709	9371	1251	417	1800	1.1(10)-185	Escherichia coli	b1748	[pn:hypothetical protein] [gn:csic]
CONTIG498	34039076_fl_34	3710	9372	516	172	491	5.5(10)-47	Escherichia coli	b1743	[pn:hypothetical protein] [gn:spy]
CONTIG498	34567181_fl_39	3711	9373	339	113	454	4.5(10)-43	Escherichia coli	b1738	[pn:pts system, cellobiose-specific iib component] [gn:cela]
CONTIG498	36516660_fl_43	3712	9374	1374	458	2120	1.3(10)-219	Escherichia coli	b1734	[pn:phospho-beta-glucosidase b] [gn:celf]
CONTIG498	14492058_fl_56	3713	9375	570	190	609	1.7(10)-59	Escherichia coli	b1726	[pn:hypothetical protein]
CONTIG498	32539012_fl_60	3714	9376	936	312	1221	2.3(10)-124	Escherichia coli	b1722	[pn:hypothetical protein]
CONTIG498	4876318_fl_62	3715	9377	1938	646	3261	0	Escherichia coli	b1719	[pn:threonyl-tRNA synthetase] [gn:thrs]
CONTIG498	36125268_fl_63	3716	9378	321	107	373	1.8(10)-34	Escherichia coli	b1718	[pn:initiation factor if-3] [gn:inf3]

CONTIG498	33594575_f2_80	3717	9379	705	235	788	1.8(10)-78	Escherichia coli	b1758	[pn:hypothetical protein]
CONTIG498	33789193_f2_88	3718	9380	1500	500	2058	4.9(10)-213	Escherichia coli	b1746	[pn:hypothetical protein]
CONTIG498	32320932_f2_96	3719	9381	360	120	548	5.0(10)-53	Escherichia coli	b1739	[pn:osmotically inducible protein e precursor] [gn:osme]
CONTIG498	5289837_f3_111	3720	9382	291	97	398	4.0(10)-37	Escherichia coli	b1777	[pn:hypothetical protein]
CONTIG498	11057961_f3_120	3721	9383	1056	352	1608	2.3(10)-165	Escherichia coli	b1764	[pn:selenophosphate synthase] [gn:seld]
CONTIG498	892316_f3_124	3722	9384	342	114	335	1.8(10)-30	Escherichia coli	b1760	[pn:hypothetical protein]
CONTIG498	3252266_f3_141	3723	9385	1140	380	1521	4.0(10)-156	Escherichia coli	b1747	[pn:hypothetical protein]
CONTIG498	29980152_f3_144	3724	9386	1359	453	1724	1.2(10)-177	Escherichia coli	b1745	[pn:hypothetical protein]
CONTIG498	4504818_f3_145	3725	9387	975	325	977	1.8(10)-98	Escherichia coli	b1744	[pn:hypothetical protein] [gn:ydis]
CONTIG498	17052038_f3_146	3726	9388	708	236	644	3.3(10)-63	Escherichia coli	b1742	[pn:hypothetical protein]
CONTIG498	11720093_f3_151	3727	9389	1371	457	1873	2.0(10)-193	Escherichia coli	b1737	[pn:pts system, cellobiose-specific iic component] [gn:celb]
CONTIG498	26181562_f3_152	3728	9390	351	117	467	1.8(10)-44	Escherichia coli	M93573	or:escherichia coli pn:pts enzyme iii cel gn:celc le:1 re:351 di:direct sr:escherichia coli (individual isolate fn23/human/sweden, strain eco nt:putative
CONTIG498	33651711_f3_153	3729	9391	834	278	1136	2.5(10)-115	Escherichia coli	b1735	[pn:cel operon repressor] [gn:celd]
CONTIG498	14657188_f3_155	3730	9392	786	262	949	1.6(10)-95	Escherichia coli	b1733	[pn:ydc]
CONTIG498	125063_f3_159	3731	9393	267	89	359	5.4(10)-33	Escherichia coli	b1731	[pn:hypothetical protein]
CONTIG498	33869002_c1_181	3732	9394	294	98	295	3.2(10)-26	Escherichia coli	b1724	[pn:hypothetical protein]
CONTIG498	22536630_c1_182	3733	9395	888	296	1320	7.9(10)-135	Escherichia coli	b1725	[pn:hypothetical protein]
CONTIG498	270402_c1_186	3734	9396	615	205	968	1.6(10)-97	Escherichia coli	b1728	[pn:hypothetical protein]

CONTIG498	26734393_c1_190	3735	9397	2283	761	3404	0	Escherichia coli	b1732	[pn:catalase hpi1] [gn:kate]
CONTIG498	35647706_c1_197	3736	9398	951	317	1117	2.6(10)-113	Escherichia coli	b1741	[pn:hypothetical protein]
CONTIG498	32109831_c1_203	3737	9399	483	161	95	0.001	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:59 re:580 di:direct nt:orf2
CONTIG498	16688291_c1_210	3738	9400	1545	515	635	2.0(10)-106	Escherichia coli	b1755	[pn:hypothetical protein]
CONTIG498	1961575_c1_212	3739	9401	1404	468	1916	5.5(10)-198	Escherichia coli	b1757	[pn:hypothetical protein]
CONTIG498	24728175_c1_216	3740	9402	429	143	435	4.7(10)-41	Escherichia coli	b1759	[pn:hypothetical protein]
CONTIG498	29890942_c2_244	3741	9403	924	308	1015	1.7(10)-102	Escherichia coli	b2842	[pn:2-deoxy-d-gluconate 3-dehydrogenase] [gn:kdud]
CONTIG498	14730277_c2_245	3742	9404	1407	469	1752	1.3(10)-180	Escherichia coli	b1729	[pn:hypothetical protein]
CONTIG498	16614825_c2_276	3743	9405	882	294	1322	4.7(10)-135	Escherichia coli	b1749	[pn:exodeoxyribonuclease iii] [gn:xtha]
CONTIG498	24663132_c2_277	3744	9406	690	230	173	2.7(10)-13	Escherichia coli	b1750	[pn:hypothetical protein]
CONTIG498	32453418_c2_278	3745	9407	1176	392	1434	6.5(10)-147	Escherichia coli	b1754	[pn:hypothetical protein]
CONTIG498	3922338_c2_288	3746	9408	1359	453	2125	3.8(10)-220	Escherichia coli	b1761	[pn:nadp-specific glutamate dehydrogenase] [gn:gdha]
CONTIG498	13100905_c2_297	3747	9409	1128	376	1636	2.6(10)-168	Escherichia coli	b1767	[pn:l-asparaginase i] [gn:ansa]
CONTIG498	4392318_c2_298	3748	9410	651	217	969	1.2(10)-97	Escherichia coli	b1768	[pn:hypothetical 23.4 kd protein in ansa 3'''region] [gn:ydb]
CONTIG498	22738257_c2_299	3749	9411	1365	455	403	5.4(10)-37	Bacillus circulans	P20533	chitinase a1 precursor (ec 3.2.1.14).
CONTIG498	4354837_c3_304	3750	9412	945	315	1266	4.2(10)-129	Escherichia coli	b1723	[pn:6-phosphofructokinase isozyme] [gn:pfkb]
CONTIG498	5907943_c3_307	3751	9413	684	228	980	8.4(10)-99	Escherichia coli	b1727	[pn:hypothetical protein]
CONTIG498	24266652_c3_320	3752	9414	936	312	1257	3.7(10)-128	Escherichia coli	b1740	[pn:nh3-dependent nad synthetase] [gn:nade]
CONTIG498	12230453_c3_336	3753	9415	1143	381	568	3.7(10)-55	Escherichia coli	b1756	[pn:hypothetical protein]

CONTIG498	2860216_c3_340	3754	9416	261	87	91	0.00093	Pseudomonas aeruginosa	M32077	or:pseudomonas aeruginosa lc:2079 re:3137 di:direct sr.p.aeruginosa (strain pao, isolate pa02003) dna, from patient nt:alginate regulatory protein p; (put.); putative
CONTIG498	4461068_c3_346	3755	9417	1866	622	2525	1.6(10)-262	Escherichia coli	b1766	[pn:protease iv] [gn:sppa]
CONTIG499	22150281_fl_23	3756	9418	1575	525	2423	1.0(10)-251	Escherichia coli	b0074	[pn:2-isopropylmalate synthase] [gn:leua]
CONTIG499	26750286_fl_37	3757	9419	1521	507	2481	7.4(10)-258	Escherichia coli	b0062	[pn:l-arabinose isomerase] [gn:araa]
CONTIG499	14570792_fl_42	3758	9420	2919	973	4561	0	Escherichia coli	b0059	[pn:probable atp-dependent helicase hepa] [gn:hepa]
CONTIG499	24089208_fl_43	3759	9421	499	167	724	1.1(10)-71	Escherichia coli	b0058	[pn:hypothetical 24.9 kd protein in sura-hepa intergenic region] [gn:yabo]
CONTIG499	4736717_f2_82	3760	9422	1413	471	2250	2.2(10)-233	Escherichia coli	b0072	[pn:3-isopropylmalate dehydratase] [gn:leuc]
CONTIG499	32538577_f2_90	3761	9423	1611	537	2192	3.1(10)-227	Escherichia coli	b0067	[pn:hypothetical 59.6 kd protein in arac-tbpa intergenic region] [gn:yabk]
CONTIG499	22445442_f3_131	3762	9424	1116	372	1535	1.3(10)-157	Escherichia coli	b0073	[pn:3-isopropylmalate dehydrogenase] [gn:leub]
CONTIG499	15057762_f3_132	3763	9425	621	207	946	3.3(10)-95	Escherichia coli	b0071	[pn:3-isopropylmalate dehydratase] [gn:leud]
CONTIG499	33640625_f3_134	3764	9426	1680	560	2374	1.6(10)-246	Escherichia coli	b0069	[pn:hypothetical 63.9 kd protein in tbpa-leud intergenic region] [gn:yabn]
CONTIG499	34245791_f3_135	3765	9427	1047	349	1413	1.1(10)-144	Escherichia coli	b0068	[pn:thiamine-binding periplasmic protein precursor] [gn:tbpa]
CONTIG499	29941042_f3_138	3766	9428	921	307	828	1.1(10)-82	Escherichia coli	b0066	[pn:hypothetical abc transporter in arac-tbpa intergenic region]
CONTIG499	10937566_f3_141	3767	9429	1770	590	2439	2.1(10)-253	Escherichia coli	b0063	[pn:l-ribulokinase] [gn:arab]
CONTIG499	15755192_f3_145	3768	9430	828	276	1193	2.2(10)-121	Escherichia coli	b0061	[pn:l-ribulose-5-phosphate 4-epimerase] [gn:arad]
CONTIG499	31765657_f3_146	3769	9431	2514	838	3674	0	Escherichia coli	b0060	[pn:dna polymerase ii] [gn:polb]

CONTIG499	3211058_c1_165	3770	9432	804	268	920	1.8(10)-92	Escherichia coli	b0065	[pn:hypothetical 26.3 kd protein in arac-tbpa intergenic region]
CONTIG499	33722680_c1_171	3771	9433	1263	421	1669	8.1(10)-172	Escherichia coli	b0070	[pn:hypothetical 42.7 kd protein in tbpa-leud intergenic region]
CONTIG499	34632808_c1_183	3772	9434	513	171	758	2.7(10)-75	Escherichia coli	b0078	[pn:acetolactate synthase isozyme iii small subunit] [gn:ilvh]
CONTIG499	15752150_c1_187	3773	9435	399	133	519	6.0(10)-50	Escherichia coli	b0083	[pn:cell division protein ftsl] [gn:ftsll]
CONTIG499	32539093_c1_188	3774	9436	1782	594	2669	8.8(10)-278	Escherichia coli	b0084	[pn:penicillin-binding protein 3 precursor] [gn:ftsll]
CONTIG499	26256317_c1_191	3775	9437	1437	479	1871	3.2(10)-193	Escherichia coli	b0086	[pn:d-alanyl] [gn:murf]
CONTIG499	12265765_c1_192	3776	9438	1278	426	1637	2.0(10)-168	Escherichia coli	b0089	[pn:cell division protein ftsw] [gn:ftsll]
CONTIG499	832575_c1_197	3777	9439	564	188	674	2.2(10)-66	Escherichia coli	b0093	[pn:cell division protein ftsq] [gn:ftsll]
CONTIG499	6527_c2_236	3778	9440	1011	337	1318	1.3(10)-134	Escherichia coli	b0076	[pn:leuo]
CONTIG499	4492650_c2_237	3779	9441	987	329	1500	6.7(10)-154	Escherichia coli	b0082	[pn:hypothetical 34.9 kd protein in frur-ftsll intergenic region] [gn:yabc]
CONTIG499	16987800_c2_240	3780	9442	1509	503	2125	3.8(10)-220	Escherichia coli	b0085	[pn:meso-diaminopimelate-adding enzyme] [gn:mure]
CONTIG499	3398507_c2_244	3781	9443	1416	472	1859	6.0(10)-192	Escherichia coli	b0088	[pn:udp-n-acetylmutamoyl-l-alanine] [gn:murd]
CONTIG499	13869067_c2_247	3782	9444	1479	493	2196	1.2(10)-227	Escherichia coli	b0091	[pn:udp-n-acetylmutamate] [gn:mure]
CONTIG499	1172057_c3_266	3783	9445	1023	341	1299	1.3(10)-132	Escherichia coli	b0064	[pn:arabinose operon regulatory protein] [gn:arac]
CONTIG499	14485081_c3_287	3784	9446	1755	585	2612	9.6(10)-272	Escherichia coli	b0077	[pn:acetolactate synthase isozyme iii large subunit] [gn:ilvi]
CONTIG499	5897968_c3_288	3785	9447	1014	338	1684	2.1(10)-173	Escherichia coli	b0080	[pn:fructose repressor] [gn:frur]
CONTIG499	36016382_c3_290	3786	9448	513	171	705	1.2(10)-69	Escherichia coli	b0081	[pn:hypothetical 17.4 kd protein in frur-ftsll intergenic region] [gn:yabb]
CONTIG499	35333290_c3_295	3787	9449	1128	376	1620	1.3(10)-166	Escherichia coli	b0087	[pn:phospho-n-acetylmutamoyl-pentapeptide- transferase] [gn:mray]
CONTIG499	35244787_c3_298	3788	9450	1107	369	1655	2.5(10)-170	Escherichia coli	b0090	[pn:udp-n-acetylglucosamine] [gn:murg]

CONTIG499	2927042_c3_299	3789	9451	948	316	1322	4.7(10)-135	Escherichia coli	b0092	[pn:d-alanine] [gn:ddlb]
CONTIG5	14265875_c3_6	3790	9452	465	155	225	9.3(10)-25	Escherichia coli	b0368	[pn:hypothetical protein] [gn:taud]
CONTIG50	31847506_f2_1	3791	9453	729	243	1149	1.0(10)-116	Escherichia coli	b0096	[pn:udp-3-o-3-hydroxymyristoyl n-acetylglucosamine deacetylase] [gn:lpxc]
CONTIG500	16464136_f1_8	3792	9454	846	282	1294	4.5(10)-132	Escherichia coli	b3697	[pn:hypothetical 29.7 kd protein in ibpa-gyrb intergenic region]
CONTIG500	31297917_f1_10	3793	9455	978	326	1268	2.6(10)-129	Escherichia coli	b3693	[pn:hypothetical 31.4 kd protein in ibpa-gyrb intergenic region]
CONTIG500	26380382_f1_11	3794	9456	336	112	92	0.0015	Escherichia coli	b3692	[pn:hypothetical 64.0 kd protein in ibpa-gyrb intergenic region]
CONTIG500	14316958_f1_12	3795	9457	1203	401	1929	2.2(10)-199	Escherichia coli	b3692	[pn:hypothetical 64.0 kd protein in ibpa-gyrb intergenic region]
CONTIG500	1265891_f1_13	3796	9458	1398	466	2061	2.3(10)-213	Escherichia coli	b3691	[pn:hypothetical 48.8 kd protein in ibpa-gyrb intergenic region]
CONTIG500	5328280_f1_14	3797	9459	1281	427	1771	1.3(10)-182	Escherichia coli	b3689	[pn:hypothetical 46.4 kd protein in ibpa-gyrb intergenic region]
CONTIG500	24712882_f1_16	3798	9460	465	155	614	5.0(10)-60	Escherichia coli	b3686	[pn:hsls] [gn:ibpb]
CONTIG500	36455080_f1_17	3799	9461	1851	617	2260	1.8(10)-234	Escherichia coli	b3685	[pn:hypothetical 58.9 kd protein in ibpb 3'''region] [gn:yide]
CONTIG500	24254062_f1_20	3800	9462	1674	558	1724	1.2(10)-177	Escherichia coli	b3683	[pn:pts system arbutin-like iic component] [gn:glvc]
CONTIG500	9847285_f1_24	3801	9463	711	237	160	6.5(10)-12	Escherichia coli	D90866	or: escherichia coli pn:d-serine dehydratase transcriptional activator gn:dsdc le:11564 re:11818 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [pir accession number a28674];
CONTIG500	979582_f1_26	3802	9464	954	318	1262	1.1(10)-128	Escherichia coli	b2364	[pn:d-serine deaminase activator] [gn:dsdc]
CONTIG500	24228532_f1_27	3803	9465	1116	372	93	0.098	Clostridium sp.	P23340	endoglucanase c307 precursor (ec 3.2.1.4) (endo-1,4-beta-glucanase) (cellulase).
CONTIG500	4554818_f1_45	3804	9466	462	154	598	2.5(10)-58	Escherichia coli	b3663	[pn:hypothetical 18.2 kd protein in nlpA-uhpt intergenic region]

CONTIG500	2994757_f1_47	3805	9467	1233	411	1251	1.6(10)-127	Escherichia coli	b3662	[pn:hypothetical 43.6 kd protein in nlpa 3 <sup>'''</sup> region] [gn:yicm]
CONTIG500	4551942_f2_55	3806	9468	1125	375	1762	1.1(10)-181	Escherichia coli	b3700	[pn:protein] [gn:recf]
CONTIG500	24417336_f2_60	3807	9469	1038	346	333	3.1(10)-30	Bacillus subtilis	yogA	[pn:hypothetical protein]
CONTIG500	33750965_f2_61	3808	9470	717	239	556	7.2(10)-54	Escherichia coli	b3695	[pn:hypothetical transcriptional regulator in ibpa-gyrb intergenic region] [gn:yidw]
CONTIG500	3210443_f2_63	3809	9471	630	210	669	7.5(10)-66	Escherichia coli	b3692	[pn:hypothetical 64.0 kd protein in ibpa-gyrb intergenic region]
CONTIG500	15806417_f2_73	3810	9472	378	126	371	2.8(10)-34	Escherichia coli	b3676	[pn:hypothetical 12.8 kd protein in ivb1-ibpb intergenic region]
CONTIG500	6362807_f2_79	3811	9473	459	153	97	3.1(10)-5	Escherichia coli	D90866	or:escherichia coli pn:d-serine dehydratase transcriptional activator gn:dsdc le:10864 re:11133 di:complement sr:escherichia coli (strain:k12) dna, clone lib:kohara lambda minise nt:similar to [pir accession number a28674];
CONTIG500	4898593_f2_86	3812	9474	429	143	390	2.7(10)-36	Escherichia coli	b3082	[pn:hypothetical 15.0 kd protein in ebgc-exut intergenic region] [gn:yglm]
CONTIG500	32541507_f2_87	3813	9475	1731	577	2464	4.7(10)-256	Escherichia coli	b3671	[pn:acetoxyhydroxy acid synthase i, small subunit] [gn:ilvb]
CONTIG500	24790908_f2_88	3814	9476	291	97	435	4.7(10)-41	Escherichia coli	b3670	[pn:acetoxyhydroxy acid synthase i, small subunit] [gn:ilvn]
CONTIG500	4822086_f2_90	3815	9477	1515	505	1640	9.6(10)-169	Escherichia coli	b3668	[pn:sensor protein uhpB] [gn:uhpb]
CONTIG500	29430341_f2_96	3816	9478	1017	339	167	2.0(10)-10	Bacillus subtilis	yfhM	[pn:hypothetical protein]
CONTIG500	22462782_f2_99	3817	9479	531	177	94	9.5(10)-5	Helicobacter pylori	HP0641	[pn:h]
CONTIG500	14316406_f3_105	3818	9480	1110	370	1669	8.1(10)-172	Escherichia coli	b3701	[pn:dna polymerase iii beta-subunit] [gn:dnan]
CONTIG500	26432887_f3_106	3819	9481	2430	810	3700	0	Escherichia coli	b3699	[pn:dna gyrase, subunit b] [gn:gyrb]
CONTIG500	22870125_f3_120	3820	9482	429	143	607	2.7(10)-59	Escherichia coli	b3687	[pn:hslT] [gn:ibpa]

CONTIG500	34156516_f3_129	3821	9483	1479	493	1696	1.1(10)-174	Bacillus subtilis	glvA	[pn:6-phospho-alpha-glucosidase] [gn:glvg]
CONTIG500	30707515_f3_130	3822	9484	363	121	287	2.2(10)-25	Escherichia coli	b3675	[pn:hypothetical 13.8 kd protein in ivb1-ibpb intergenic region]
CONTIG500	15085840_f3_141	3823	9485	342	114	349	6.2(10)-32	Escherichia coli	b3083	[pn:hypothetical 12.1 kd protein in ebgc-exut intergenic region] [gn:ygin]
CONTIG500	14930291_f3_143	3824	9486	768	256	932	1.0(10)-93	Escherichia coli	b3669	[pn:transcriptional regulatory protein uhpa] [gn:uhpa]
CONTIG500	4333318_f3_145	3825	9487	1320	440	1875	1.2(10)-193	Escherichia coli	b3667	[pn:regulatory protein uhpc] [gn:uhpc]
CONTIG500	12582291_f3_146	3826	9488	1524	508	2073	1.3(10)-214	Escherichia coli	b3666	[pn:hexosephosphate transport protein] [gn:uhpt]
CONTIG500	5116067_c1_160	3827	9489	315	105	223	1.3(10)-18	Bacillus subtilis	licA	[pn:phosphotransferase system] [gn:celc]
CONTIG500	24226552_c1_161	3828	9490	1407	469	190	3.7(10)-12	Escherichia coli	b4036	[pn:phage lambda receptor protein] [gn:lamb]
CONTIG500	24101517_c1_162	3829	9491	1050	350	1083	1.0(10)-109	Escherichia coli	b3660	[pn:hypothetical 33.1 kd protein in nlpa 5''' region] [gn:yic]
CONTIG500	4332811_c1_168	3830	9492	912	304	426	4.2(10)-40	Pseudomonas putida	P10183	transcriptional activator protein nahr.
CONTIG500	10837763_c1_190	3831	9493	1341	447	1507	1.2(10)-154	Escherichia coli	b2365	[pn:dsdx permease] [gn:dsdx]
CONTIG500	34557265_c1_199	3832	9494	756	252	928	2.7(10)-93	Escherichia coli	b3684	[pn:hypothetical transcriptional regulator in ilvo-ibpb intergenic region] [gn:yidp]
CONTIG500	17004052_c1_215	3833	9495	342	114	99	5.5(10)-5	Escherichia coli	U27192	or:escherichia coli pn:modd gn:modd le:5360 re:6055 di:direct
CONTIG500	25878136_c2_226	3834	9496	1368	456	1543	1.8(10)-158	Bacillus subtilis	ydhP	[pn:hypothetical protein]
CONTIG500	23940636_c2_255	3835	9497	855	285	100	0.00839	Haemophilus influenzae	H11538	[pn:lic-1 operon protein] [gn:licb]
CONTIG500	34415953_c2_277	3836	9498	336	112	448	2.0(10)-42	Escherichia coli	b3688	[pn:ol35] [gn:yidq]
CONTIG500	10626535_c2_291	3837	9499	948	316	407	4.4(10)-38	Escherichia coli	b0208	[pn:hypothetical transcriptional regulator in rnh-dnir intergenic region] [gn:yafc]
CONTIG500	33729167_c2_292	3838	9500	1446	482	291	4.5(10)-28	Haloferax sp.	P21562	hypothetical 80.2 kd protein in the 5' region of gyra and gyrb (orf 4).



CONTIG500	31273568_c2_293	3839	9501	1059	353	349	5.0(10)-31	Haloflex sp.	P21562	hypothetical 80.2 kd protein in the S' region of gyra and gyrb (orf 4).
CONTIG500	13677158_c3_326	3840	9502	291	97	136	2.2(10)-9	Escherichia coli	U70214	or:escherichia coli le:84358 re:84669 di:complement nt:hypothetical
CONTIG500	1984555_c3_330	3841	9503	1230	410	1118	2.0(10)-113	Escherichia coli	b3673	[pn:multidrug resistance protein d] [gn:emrd]
CONTIG500	4859538_c3_336	3842	9504	1389	463	1854	2.0(10)-191	Escherichia coli	b2366	[pn:d-serine dehydratase] [gn:dsda]
CONTIG501	34093886_fl_3	3843	9505	1983	661	3188	0	Escherichia coli	b4069	[pn:acetyl-coa synthetase] [gn:acs]
CONTIG501	10039711_fl_4	3844	9506	1680	560	2193	2.3(10)-227	Escherichia coli	b4067	[pn:hypothetical 59.2 kd protein in soxr-acs intergenic region] [gn:yjcg]
CONTIG501	26306592_fl_52	3845	9507	1203	401	1883	1.7(10)-194	Escherichia coli	b4034	[pn:periplasmic maltose-binding protein] [gn:male]
CONTIG501	17074051_fl_56	3846	9508	909	303	1253	9.9(10)-128	Escherichia coli	b4032	[pn:maltose transport inner membrane protein] [gn:malg]
CONTIG501	21501708_fl_63	3847	9509	282	94	99	0.0006	Streptomyces ambifaciens	Z46913	or:streptomyces ambifaciens pn:polyketide synthase le:<1 re:>3596 di:direct nt:putative
CONTIG501	14554662_f2_71	3848	9510	348	116	366	9.8(10)-34	Escherichia coli	b4068	[pn:hypothetical 11.7 kd protein in soxr-acs intergenic region] [gn:yjch]
CONTIG501	4977000_f2_75	3849	9511	897	299	540	3.6(10)-52	Bacillus subtilis	ywbl	[pn:hypothetical protein] [gn:ipa-24d]
CONTIG501	3409812_f2_83	3850	9512	240	80	204	1.3(10)-16	Escherichia coli	b4062	[pn:regulatory protein soxs] [gn:soxs]
CONTIG501	898566_f2_89	3851	9513	333	111	356	1.1(10)-32	Escherichia coli	b4060	[pn:hypothetical 13.0 kd protein in ssb-soxs intergenic region] [gn:yjcb]
CONTIG501	5267327_f2_91	3852	9514	2823	941	4604	0	Escherichia coli	b4058	[pn:excision nuclease] [gn:uvra]
CONTIG501	26421891_f2_94	3853	9515	303	101	175	1.7(10)-13	Escherichia coli	Z26592	or:escherichia coli pn:dna binding protein sp:p36558 le:1086 re:1376 di:direct
CONTIG501	2552281_f2_100	3854	9516	1110	370	1519	6.5(10)-156	Escherichia coli	b4051	[pn:quinone oxidoreductase] [gn:qor]
CONTIG501	14569216_f2_109	3855	9517	2502	834	3793	0	Escherichia coli	b4041	[pn:glycerol-3-phosphate acyltransferase] [gn:plsB]
CONTIG501	12307842_f2_120	3856	9518	1557	519	2116	3.5(10)-219	Escherichia coli	b4033	[pn:maltose transport inner membrane protein] [gn:malF]

CONTIG501	16147918_f3_151	3857	9519	552	184	150	5.9(10)-10	Pseudomonas aeruginosa	U50396	or:pseudomonas aeruginosa pn:wbpn gn:wbpn le:22302 re:23693 di:direct
CONTIG501	15680438_f3_152	3858	9520	522	174	248	1.2(10)-20	Pseudomonas aeruginosa	U50396	or:pseudomonas aeruginosa pn:wbpn gn:wbpn le:22302 re:23693 di:direct
CONTIG501	276642_f3_161	3859	9521	552	184	849	6.4(10)-85	Escherichia coli	b4046	[pn:hypothetical 21.7 kd protein in difn-qor intergenic region] [gn:yjbl]
CONTIG501	35242955_c1_184	3860	9522	216	72	93	0.00259	Saccharomyces cerevisiae	YJR151C	[pn:similarity to mucin proteins, ykl224c, sta1p] [gn:j2223]
CONTIG501	11883290_c1_189	3861	9523	2139	713	3366	0	Escherichia coli	b4029	[pn:hypothetical 78.5 kd protein in pgi-xyle intergenic region] [gn:yjbl]
CONTIG501	14113277_c1_190	3862	9524	468	156	475	2.7(10)-45	Escherichia coli	b4030	[pn:hypothetical 15.6 kd protein in pgi-xyle intergenic region] [gn:yjbl]
CONTIG501	5868877_c1_202	3863	9525	570	190	756	4.5(10)-75	Escherichia coli	b4039	[pn:chorismate lyase] [gn:ubic]
CONTIG501	24821088_c1_203	3864	9526	882	294	1175	1.8(10)-119	Escherichia coli	b4040	[pn:4-hydroxybenzoate-octaprenyl transferase] [gn:ubia]
CONTIG501	16087_c1_206	3865	9527	381	127	416	4.9(10)-39	Escherichia coli	b4042	[pn:diacylglycerol kinase] [gn:dgka]
CONTIG501	22298201_c1_207	3866	9528	633	211	949	1.6(10)-95	Escherichia coli	b4043	[pn:lexa] [gn:lexa]
CONTIG501	33753275_c1_208	3867	9529	1380	460	1701	3.2(10)-175	Escherichia coli	b4044	[pn:dna-damage-inducible protein f] [gn:dinf]
CONTIG501	4425037_c1_212	3868	9530	1005	335	1606	3.8(10)-165	Escherichia coli	b4049	[pn:hypothetical 38.4 kd protein in difn-qor intergenic region] [gn:yjbl]
CONTIG501	4859693_c1_216	3869	9531	1089	363	1628	1.8(10)-167	Escherichia coli	b4053	[pn:alanine racemase] [gn:alr]
CONTIG501	12579812_c1_221	3870	9532	447	149	631	8.0(10)-62	Escherichia coli	b4056	[pn:hypothetical 15.7 kd protein in tyrb-uvra intergenic region]
CONTIG501	11958316_c1_225	3871	9533	225	75	98	2.5(10)-5	Escherichia coli	J01721	or:escherichia coli le:124 re:300 di:complement sr:escherichia coli, pdr1996 plasmid dna nt:single-stranded dna-binding protein (ssb)
CONTIG501	3908567_c1_231	3872	9534	468	156	749	2.5(10)-74	Escherichia coli	b4063	[pn:soxr protein] [gn:soxr]
CONTIG501	32453126_c1_236	3873	9535	1794	598	2089	2.6(10)-216	Escherichia coli	b4065	[pn:hypothetical 60.5 kd protein in soxr-acs intergenic region] [gn:yjce]
CONTIG501	17036340_c1_240	3874	9536	756	252	257	3.5(10)-22	Bacillus subtilis	ywbG	[pn:hypothetical protein] [gn:ipa-22r]

CONTIG501	24035252_c1_248	3875	9537	1260	420	1530	4.4(10)-157	Escherichia coli	b4077	[pn:glutamate- aspartate carrier] [gn:gltp]
CONTIG501	36343956_c2_251	3876	9538	840	280	805	3.0(10)-80	Escherichia coli	b4028	[pn:hypothetical 26.3 kd protein in pgi-xyle intergenic region] [gn:yibg]
CONTIG501	23884712_c2_273	3877	9539	369	123	350	4.9(10)-32	Escherichia coli	b4050	[pn:hypothetical 17.4 kd protein in dnf-qor intergenic region] [gn:yibo]
CONTIG501	22477280_c2_278	3878	9540	1209	403	1795	3.7(10)-185	Escherichia coli	b4054	[pn:tyrosine aminotransferase] [gn:tyrb]
CONTIG501	9932188_c2_284	3879	9541	549	183	531	3.2(10)-51	Escherichia coli	b4059	[pn:single-strand dna-binding protein] [gn:ssb]
CONTIG501	2236375_c2_285	3880	9542	1476	492	220	5.0(10)-15	Escherichia coli	b1285	[pn:hypothetical protein] [gn:ycir]
CONTIG501	1581_c2_288	3881	9543	738	246	305	2.8(10)-27	Saccharomyces cerevisiae	P40582	hypothetical 26.8 kd protein in hyr1 3'region.
CONTIG501	35287516_c2_289	3882	9544	1419	473	1796	2.8(10)-185	Escherichia coli	b4064	[pn:hypothetical 45.7 kd protein in soxr-acs intergenic region] [gn:yjcd]
CONTIG501	13958261_c3_305	3883	9545	1893	631	2763	9.6(10)-288	Escherichia coli	b4025	[pn:glucose-6-phosphate isomerase] [gn:pgi]
CONTIG501	35551431_c3_307	3884	9546	681	227	961	8.6(10)-97	Escherichia coli	b4027	[pn:hypothetical 25.0 kd lipoprotein in pgi-xyle intergenic region]
CONTIG501	24824066_c3_316	3885	9547	1125	375	1774	6.0(10)-183	Escherichia coli	b4035	[pn:cytoplasmic membrane protein for maltose uptake] [gn:malk]
CONTIG501	956308_c3_317	3886	9548	1353	451	1872	2.5(10)-193	Escherichia coli	b4036	[pn:phage lambda receptor protein] [gn:lamb]
CONTIG501	32547893_c3_318	3887	9549	996	332	1128	1.8(10)-114	Escherichia coli	b4037	[pn:maltose operon periplasmic protein] [gn:malm]
CONTIG501	14276661_c3_329	3888	9550	225	75	355	1.3(10)-32	Escherichia coli	b4045	[pn:hypothetical protein] [gn:yjbi]
CONTIG501	12315630_c3_336	3889	9551	1452	484	2320	8.5(10)-241	Escherichia coli	b4052	[pn:replicative dna helicase] [gn:dnab]
CONTIG501	20516561_c3_341	3890	9552	738	246	973	4.7(10)-98	Escherichia coli	b4055	[pn:hypothetical 26.1 kd protein in tyrb-uvra intergenic region]
CONTIG501	24079387_c3_343	3891	9553	384	128	515	1.6(10)-49	Escherichia coli	b4057	[pn:hypothetical 13.4 kd protein in tyrb-uvra intergenic region] [gn:yjbr]
CONTIG501	31895161_c3_349	3892	9554	588	196	155	6.9(10)-10	Micrococcus luteus	JQ0406	hypothetical protein 1246 (uvra region) - micrococcus luteus(fragment)
CONTIG501	32212775_c3_354	3893	9555	1650	550	1657	1.5(10)-170	Escherichia coli	b4061	[pn:hypothetical 60.8 kd protein in ssb-soxs intergenic region] [gn:yjcc]

CONTIG501	6719025_c3_363	3894	9556	477	159	145	2.6(10)-10	Bacillus subtilis	ywbH	[pn:hypothetical protein] [gn:ipa-23r]
CONTIG502	4093891_fl_14	3895	9557	1533	511	1600	1.7(10)-164	Escherichia coli	b0621	[pn:hypothetical protein] [gn:dcuc]
CONTIG502	19792580_fl_16	3896	9558	354	118	372	2.2(10)-34	Escherichia coli	b2387	[pn:hypothetical protein]
CONTIG502	11897505_fl_17	3897	9559	1275	425	1432	1.1(10)-146	Escherichia coli	b2386	[pn:hypothetical protein]
CONTIG502	15908192_fl_18	3898	9560	1122	374	1339	7.7(10)-137	Escherichia coli	b2384	[pn:hypothetical protein]
CONTIG502	5941376_fl_32	3899	9561	984	328	780	1.3(10)-77	Escherichia coli	b0603	[pn:hypothetical protein]
CONTIG502	14882962_fl_38	3900	9562	621	207	306	2.2(10)-27	Bacillus subtilis	ykrY	[pn:hypothetical protein]
CONTIG502	6040765_fl_44	3901	9563	1311	437	763	8.3(10)-76	Bacillus subtilis	ykrI	[pn:hypothetical protein]
CONTIG502	24243791_fl_53	3902	9564	1113	371	1416	5.2(10)-145	Escherichia coli	b0599	[pn:hypothetical protein in csta 3"]
CONTIG502	32286450_fl_54	3903	9565	801	267	419	2.3(10)-39	Escherichia coli	b4249	[pn:hypothetical 24.6 kd protein in pylr-argi intergenic region] [gn:yjgi]
CONTIG502	4971937_fl_55	3904	9566	372	124	101	1.2(10)-5	Methanococcus jannaschii	MJ1103	[pn:conserved hypothetical protein]
CONTIG502	34178885_f2_75	3905	9567	891	297	1078	3.5(10)-109	Escherichia coli	b0611	[pn:ribonuclease i precursor] [gn:rma]
CONTIG502	12370750_f2_77	3906	9568	447	149	441	1.1(10)-41	Escherichia coli	b2385	[pn:hypothetical protein]
CONTIG502	26250966_f2_80	3907	9569	2457	819	2860	5.0(10)-298	Escherichia coli	b2383	[pn:hypothetical protein]
CONTIG502	6923316_f2_82	3908	9570	426	142	501	4.7(10)-48	Escherichia coli	b0610	[pn:regulator of nucleoside diphosphate kinase] [gn:rnk]
CONTIG502	16490807_f2_92	3909	9571	684	228	758	2.7(10)-75	Escherichia coli	b0601	[pn:hypothetical protein]
CONTIG502	3301088_f2_103	3910	9572	582	194	121	3.7(10)-6	Hordeum vulgare	S35221	globulin beg1 precursor - barley
CONTIG502	994032_f3_134	3911	9573	831	277	798	1.6(10)-79	Escherichia coli	b2385	[pn:hypothetical protein]
CONTIG502	14664811_f3_146	3912	9574	444	148	642	5.5(10)-63	Escherichia coli	b0607	[pn:hypothetical protein] [gn:ybdq]

CONTIG502	21932036_f3_149	3913	9575	261	87	124	2.1(10)-13	Burkholderia cepacia	Q02940	beta-lactamase precursor (ec 3.5.2.6) (penicillinase).
CONTIG502	4095150_f3_151	3914	9576	1281	427	1792	7.5(10)-185	Escherichia coli	b0602	[pn:hypothetical protein]
CONTIG502	15820827_f3_156	3915	9577	723	241	943	7.0(10)-95	Klebsiella oxytoca	A49101	enolase-phosphatase e-1 - klebsiella oxytoca
CONTIG502	11036262_c1_190	3916	9578	528	176	591	1.3(10)-57	Escherichia coli	b0597	[pn:hypothetical 15.0 kd protein in enta-csta intergenic region] [gn:ybdb]
CONTIG502	14479186_c1_199	3917	9579	1644	548	803	4.7(10)-80	Bacillus subtilis	rbsA	[pn:ribose abc transporter]
CONTIG502	24415937_c1_200	3918	9580	1068	356	114	0.00046	Escherichia coli	b2548	[pn:hypothetical protein]
CONTIG502	22297582_c1_202	3919	9581	1083	361	555	9.1(10)-54	Methanococcus jannaschii	MJ0454	[pn:translation initiation factor eif-2b, subunit alpha]
CONTIG502	5320781_c1_211	3920	9582	621	207	970	9.6(10)-98	Escherichia coli	b0605	[pn:alkyl hydroperoxide reductase c22 protein] [gn:ahpc]
CONTIG502	24109377_c1_216	3921	9583	732	244	277	2.6(10)-24	Bacillus subtilis	ywrF	[pn:hypothetical protein]
CONTIG502	29970066_c1_233	3922	9584	1596	532	630	1.0(10)-61	Escherichia coli	b0574	[pn:hypothetical protein] [gn:ylcd]
CONTIG502	16975677_c1_236	3923	9585	996	332	832	4.0(10)-83	Escherichia coli	b0622	[pn:hypothetical protein in cspe 5' region] [gn:ybeg]
CONTIG502	1462634_c2_237	3924	9586	1310	436	1591	1.5(10)-163	Escherichia coli	b0593	[pn:isochorismate synthase entc] [gn:entc]
CONTIG502	4319068_c2_238	3925	9587	1620	540	2268	2.7(10)-235	Escherichia coli	b0594	[pn:2,3-dihydroxybenzoate-amp ligase] [gn:ente]
CONTIG502	15830125_c2_239	3926	9588	822	274	978	1.3(10)-98	Escherichia coli	b0596	[pn:2,3-dihydro-2,3-dihydroxybenzoate dehydrogenase] [gn:enta]
CONTIG502	6135002_c2_263	3927	9589	1098	366	999	8.1(10)-101	Escherichia coli	b2382	[pn:hypothetical protein]
CONTIG502	26343841_c2_274	3928	9590	1305	435	125	5.5(10)-5	Alcaligenes sp.	JC4698	divalent cation resistant determinant protein c - alcaligenessp. this protein is a cation/proton antiporter protein, which determines the resistancy for cadmium, zinc, cobalt.
CONTIG502	21768943_c2_278	3929	9591	2124	708	2082	1.3(10)-215	Escherichia coli	b0575	[pn:hypothetical protein in phep 5' region] [gn:ybde]

CONTIG502	29786537_c3_285	3930	9592	1035	345	1312	5.5(10)-134	Escherichia coli	b0595	[pn:isochorismatase] [gn:entb]
CONTIG502	5972011_c3_288	3931	9593	2139	713	2899	3.7(10)-302	Escherichia coli	b0598	[pn:hypothetical protein] [gn:csta]
CONTIG502	14629386_c3_289	3932	9594	312	104	270	1.5(10)-23	Escherichia coli	b4353	[pn:hypothetical 7.7 kd protein in mrr-lsr intergenic region] [gn:yjix]
CONTIG502	13786067_c3_295	3933	9595	1035	345	317	1.5(10)-28	Bacillus subtilis	rbsC	[pn:ribose abc transporter]
CONTIG502	32610082_c3_305	3934	9596	1242	414	1693	2.3(10)-174	Escherichia coli	b0600	[pn:hypothetical protein]
CONTIG502	22661668_c3_310	3935	9597	1605	535	2447	3.0(10)-254	Escherichia coli	b0606	[pn:alkyl hydroperoxide reductase f52a protein] [gn:ahpf]
CONTIG502	22363758_c3_312	3936	9598	1266	422	1759	2.3(10)-181	Escherichia coli	b0608	[pn:hypothetical protein]
CONTIG502	33722793_c3_313	3937	9599	234	78	128	1.6(10)-8	Escherichia coli	b0609	[pn:hypothetical protein]
CONTIG502	203525_c3_328	3938	9600	336	112	96	4.0(10)-5	Escherichia coli	b0573	[pn:hypothetical protein] [gn:ylcc]
CONTIG502	35673516_c3_332	3939	9601	1188	396	1204	1.5(10)-122	Escherichia coli	b0575	[pn:hypothetical protein in phev 5''' region] [gn:ybde]
CONTIG503	34069387_fl_6	3940	9602	1002	334	732	1.6(10)-72	Escherichia coli	b0147	[pn:hypothetical protein in sfsa-mrdb intergenic region] [gn:yadp]
CONTIG503	33994165_fl_7	3941	9603	477	159	689	5.7(10)-68	Escherichia coli	b0145	[pn:dosage-dependent dnak suppressor protein] [gn:dksa]
CONTIG503	261403_fl_14	3942	9604	903	301	1134	4.0(10)-115	Escherichia coli	b0133	[pn:pantoate] [gn:panc]
CONTIG503	87812_fl_15	3943	9605	615	205	140	8.6(10)-10	Escherichia coli	b0141	[pn:hypothetical fimbrial-like protein in ecpd-folk intergenic region] [gn:yadn]
CONTIG503	12944680_fl_16	3944	9606	627	209	183	2.3(10)-14	Escherichia coli	b0138	[pn:hypothetical 20.3 kd protein in panb-htr intergenic region] [gn:yadm]
CONTIG503	32035766_fl_18	3945	9607	1122	374	197	7.5(10)-16	Escherichia coli	b0135	[pn:hypothetical fimbrial-like protein in panb-htr intergenic region] [gn:yadc]
CONTIG503	22380131_fl_45	3946	9608	1803	601	1054	1.2(10)-106	Escherichia coli	b3657	[pn:hypothetical 51.0 kd protein in glts-selc intergenic region] [gn:yic]
CONTIG503	35282686_fl_50	3947	9609	987	329	1201	3.2(10)-122	Escherichia coli	b0109	[pn:nicotinate-nucleotide pyrophosphorylase] [gn:nadc]

CONTIG503	24692676_f1_53	3948	9610	1224	408	1007	1.2(10)-101	Escherichia coli	b0106	[pn:protein transport protein hofc] [gn:hofc]
CONTIG503	10835150_f1_55	3949	9611	639	213	684	2.0(10)-67	Escherichia coli	b0103	[pn:hypothetical 22.5 kd protein in multi-guac intergenic region precursor] [gn:yacc]
CONTIG503	13023261_f2_62	3950	9612	585	195	114	7.2(10)-5	Homo sapiens	Q15428	[GN:SAP62] [SR:HUMAN] [DE:SPLICEOSOME ASSOCIATED PROTEIN 62 (SAP 62) (SF3A66)] [SP:Q15428]
CONTIG503	5097193_f2_67	3951	9613	714	238	956	3.0(10)-96	Escherichia coli	b0146	[pn:sugar fermentation stimulation protein] [gn:sfsa]
CONTIG503	4582717_f2_71	3952	9614	489	163	620	1.2(10)-60	Escherichia coli	b0142	[pn:2-amino-4-hydroxy-6-hydroxymethyldihydropteridine pyrophosphokinase]
CONTIG503	34166637_f2_72	3953	9615	831	277	1237	4.9(10)-126	Escherichia coli	b0134	[pn:3-methyl-2-oxobutanoate hydroxymethyltransferase] [gn:panb]
CONTIG503	6444212_f2_74	3954	9616	393	131	506	1.3(10)-48	Escherichia coli	b0131	[pn:aspartate 1-decarboxylase] [gn:pand]
CONTIG503	23713380_f2_76	3955	9617	840	280	568	3.7(10)-55	Escherichia coli	b0140	[pn:chaperone protein ecpd precursor] [gn:ecpd]
CONTIG503	806542_f2_77	3956	9618	2598	866	1852	3.2(10)-191	Escherichia coli	b0139	[pn:outer membrane usher protein htr precursor] [gn:htrc]
CONTIG503	21754165_f2_79	3957	9619	651	217	159	8.4(10)-12	Escherichia coli	b0136	[pn:hypothetical 21.1 kd protein in panb-htrc intergenic region] [gn:yadk]
CONTIG503	14462840_f2_91	3958	9620	897	299	1402	1.6(10)-143	Escherichia coli	b0121	[pn:spermidine synthase] [gn:spee]
CONTIG503	3320325_f2_92	3959	9621	795	265	1327	1.3(10)-135	Escherichia coli	b0120	[pn:s-adenosylmethionine decarboxylase proenzyme] [gn:sped]
CONTIG503	14241436_f2_112	3960	9622	1392	464	1968	1.7(10)-203	Escherichia coli	b0112	[pn:aromatic amino acid transport protein arop] [gn:arop]
CONTIG503	33625332_f2_113	3961	9623	999	333	90	0.37	Selenomonas ruminantium	AF040720	[de:selenomonas ruminantium xylosidase/arabinosidase (xsa) gene, complete cds.] [pn:xylosidase/arabinosidase] [gn:xsa]
CONTIG503	24645887_f2_121	3962	9624	213	71	95	0.00259	Turnip yellow mosaic virus	AF035403	[de:turnip yellow mosaic blue lake isolate, complete genome.] [pn:replicase protein]

CONTIG503	2230418_f3_138	3963	9625	924	308	1271	1.2(10)-129	Escherichia coli	b0144	[pn:hypothetical 29.3 kd protein in pcnb-dksa intergenic region] [gn:yadb]
CONTIG503	129175_f3_139	3964	9626	1440	480	1951	1.1(10)-201	Escherichia coli	b0143	[pn:poly] [gn:pcnb]
CONTIG503	32048127_f3_147	3965	9627	606	202	268	2.3(10)-23	Escherichia coli	b0137	[pn:hypothetical 21.0 kd protein in panb-hire intergenic region]
CONTIG503	13695127_f3_151	3966	9628	768	256	1029	5.4(10)-104	Escherichia coli	b0126	[pn:hypothetical protein in hpt-pand intergenic region] [gn:yadf]
CONTIG503	23475311_f3_153	3967	9629	2424	808	3386	0	Escherichia coli	b0124	[pn:glucose dehydrogenase] [gn:gcd]
CONTIG503	32448952_f3_156	3968	9630	636	212	625	3.5(10)-61	Escherichia coli	b0122	[pn:hypothetical protein in spee-gcd intergenic region] [gn:yacc]
CONTIG503	22916285_f3_162	3969	9631	1554	518	807	1.8(10)-80	Escherichia coli	b0117	[pn:hypothetical protein in lpda-spel intergenic region] [gn:yach]
CONTIG503	4507318_f3_179	3970	9632	447	149	499	7.9(10)-48	Escherichia coli	b0108	[pn:prepilin peptidase dependent protein d precursor] [gn:ppdd]
CONTIG503	15089211_f3_180	3971	9633	1392	464	1500	6.7(10)-154	Escherichia coli	b0107	[pn:protein transport protein hofb] [gn:hofb]
CONTIG503	33708341_f3_185	3972	9634	837	279	1007	1.2(10)-101	Escherichia coli	b0102	[pn:hypothetical protein in mutt-guac intergenic region] [gn:yacf]
CONTIG503	5181693_f3_186	3973	9635	204	68	312	5.2(10)-28	Escherichia coli	b0101	[pn:yacg]
CONTIG503	10052077_c1_190	3974	9636	2739	913	4104	0	Escherichia coli	b0098	[pn:preprotein translocase seca subunit] [gn:seca]
CONTIG503	24017063_c1_191	3975	9637	450	150	538	5.7(10)-52	Escherichia coli	b0099	[pn:mutator mult protein] [gn:mutt]
CONTIG503	34613891_c1_195	3976	9638	1098	366	1681	4.4(10)-173	Escherichia coli	b0104	[pn:gmp reductase] [gn:guac]
CONTIG503	10042802_c1_201	3977	9639	579	193	839	7.4(10)-84	Escherichia coli	b0110	[pn:ampd protein] [gn:ampd]
CONTIG503	12367317_c1_221	3978	9640	2643	881	4131	0	Escherichia coli	b0118	[pn:aconitate hydratase 2] [gn:acnb]
CONTIG503	7032637_c1_228	3979	9641	555	185	872	2.3(10)-87	Escherichia coli	b0125	[pn:hypoxanthine phosphoribosyltransferase] [gn:hpt]
CONTIG503	4304753_c1_229	3980	9642	936	312	1453	6.4(10)-149	Escherichia coli	b0127	[pn:hypothetical abc transporter in hpt-pand intergenic region]
CONTIG503	2594632_c1_230	3981	9643	468	156	560	2.7(10)-54	Escherichia coli	b0129	[pn:hypothetical protein in hpt-pand intergenic region] [gn:yadi]



CONTIG503	837557_c1_231	3982	9644	1311	437	1647	1.8(10)-169	Escherichia coli	b0130	[pn:hypothetical 46.3 kd protein in hpt-pand intergenic region precursor] [gn:yade]
CONTIG503	34661686_c1_248	3983	9645	519	173	131	7.7(10)-9	Haemophilus influenzae	L20805	or:haemophilus influenzae le:<1 re:172 di:direct nt:truncated sequence, 32.6% identity and 67%
CONTIG503	35244781_c2_277	3984	9646	1929	643	2436	4.2(10)-253	Escherichia coli	b0115	[pn:c2 of pyruvate dehydrogenase] [gn:acef]
CONTIG503	16970218_c2_286	3985	9647	483	161	594	6.7(10)-58	Escherichia coli	b0119	[pn:hypothetical protein in lpda-sped intergenic region] [gn:yaci]
CONTIG503	81953_c2_315	3986	9648	2430	810	3244	0	Escherichia coli	b0148	[pn:atp-dependent helicase hrpb] [gn:hrpb]
CONTIG503	34634452_c2_316	3987	9649	1956	652	2530	9.8(10)-275	Escherichia coli	b0149	[pn:peptidoglycan synthetase] [gn:mrcb]
CONTIG503	22742711_c3_322	3988	9650	522	174	116	2.0(10)-6	Pseudomonas aeruginosa	P24564	hypothetical 19.5 kd protein in pilt region (orf6).
CONTIG503	4486068_c3_325	3989	9651	864	288	1164	2.7(10)-118	Escherichia coli	b0111	[pn:ampe protein] [gn:ampe]
CONTIG503	24355151_c3_332	3990	9652	795	265	1105	4.7(10)-112	Escherichia coli	b0113	[pn:pyruvate dehydrogenase complex repressor] [gn:pdhr]
CONTIG503	13870927_c3_333	3991	9653	2679	893	4352	0	Escherichia coli	b0114	[pn:pyruvate dehydrogenase c1 component] [gn:acee]
CONTIG503	5112963_c3_334	3992	9654	1434	478	2284	5.5(10)-237	Escherichia coli	b0116	[pn:dihydrolipoamide dehydrogenase] [gn:lpda]
CONTIG503	16145137_c3_343	3993	9655	447	149	199	4.9(10)-16	Xanthomonas campestris	P22264	hypothetical protein in clp 5'region (orf1) (fragment).
CONTIG503	22393826_c3_345	3994	9656	1764	588	1537	3.2(10)-210	Escherichia coli	b0123	[pn:hypothetical protein in spee-gcd intergenic region precursor] [gn:yack]
CONTIG503	12692811_c3_351	3995	9657	834	278	1000	6.4(10)-101	Escherichia coli	b0128	[pn:hypothetical 28.5 kd protein in hpt-pand intergenic region]
CONTIG504	24645388_fl_1	3996	9658	237	79	377	6.7(10)-35	Escherichia coli	b2741	[pn:ma polymerase sigma subunit rpos] [gn:rpos]
CONTIG504	5086577_fl_5	3997	9659	603	201	543	1.7(10)-52	Bacillus subtilis	yclB	[pn:hypothetical protein]
CONTIG504	24652343_fl_12	3998	9660	357	119	524	1.8(10)-50	Escherichia coli	b2732	[pn:hypothetical 13.9 kd protein in fhla-muts intergenic region] [gn:ygba]

CONTIG504	32041625_f1_31	3999	9661	624	208	474	3.5(10)-45	Escherichia coli	b2725	[pn:formate hydrogenlyase regulatory protein] [gn:hyca]
CONTIG504	26074191_f1_32	4000	9662	1827	609	1828	1.2(10)-188	Escherichia coli	b2723	[pn:formate hydrogenlyase subunit 3] [gn:hycc]
CONTIG504	1032706_f1_38	4001	9663	807	269	1176	1.3(10)-119	Escherichia coli	b2719	[pn:formate hydrogenlyase subunit 7] [gn:hyccg]
CONTIG504	6407943_f1_39	4002	9664	465	155	687	9.4(10)-68	Escherichia coli	b2717	[pn:hydrogenase 3 maturation protease] [gn:hyci]
CONTIG504	26618750_f1_65	4003	9665	1111	371	1295	3.5(10)-132	Escherichia coli	b2701	[pn:membrane-bound lytic transglycosylase b precursor] [gn:mltb]
CONTIG504	10166031_f2_71	4004	9666	1521	507	1636	2.6(10)-168	Bacillus subtilis	ycIC	[pn:hypothetical protein]
CONTIG504	2038467_f2_76	4005	9667	495	165	288	1.8(10)-25	Escherichia coli	b2848	[pn:hypothetical protein]
CONTIG504	26272556_f2_96	4006	9668	630	210	749	2.5(10)-74	Escherichia coli	b2724	[pn:formate hydrogenlyase subunit 2] [gn:hycb]
CONTIG504	21491325_f2_97	4007	9669	948	316	1094	7.0(10)-111	Escherichia coli	b2722	[pn:formate hydrogenlyase subunit 4] [gn:hycd]
CONTIG504	6917175_f2_98	4008	9670	1725	575	2877	8.0(10)-300	Escherichia coli	b2721	[pn:formate hydrogenlyase subunit 5] [gn:hyce]
CONTIG504	4425068_f2_99	4009	9671	552	184	905	7.5(10)-91	Escherichia coli	b2720	[pn:formate hydrogenlyase subunit 6] [gn:hyccf]
CONTIG504	12525131_f2_108	4010	9672	2373	791	2325	2.5(10)-241	Escherichia coli	b2712	[pn:transcriptional regulatory protein hypf] [gn:hypf]
CONTIG504	16304657_f2_113	4011	9673	1071	357	205	1.3(10)-16	Escherichia coli	U03846	or:escherichia coli le<1 re:183 di:direct nt:putative orf>60aa
CONTIG504	33709461_f3_124	4012	9674	906	302	320	7.2(10)-29	Haemophilus influenzae	HI1364	[pn:hypothetical protein]
CONTIG504	29930206_f3_126	4013	9675	258	86	189	5.5(10)-15	Bacillus subtilis	ycID	[pn:hypothetical protein]
CONTIG504	14855436_f3_130	4014	9676	900	300	563	1.3(10)-54	Escherichia coli	b2847	[pn:hypothetical protein]
CONTIG504	11885418_f3_160	4015	9677	423	141	592	1.1(10)-57	Escherichia coli	b2718	[pn:formate hydrogenlyase maturation protein] [gn:hych]
CONTIG504	32444642_f3_166	4016	9678	1053	351	1336	1.6(10)-136	Escherichia coli	b2714	[pn:asc operon repressor protein] [gn:ascg]
CONTIG504	32501052_f3_167	4017	9679	582	194	850	5.0(10)-85	Escherichia coli	b2713	[pn:4fe-4s iron-sulfur protein] [gn:hydn]

CONTIG504	32660958_f3_170	4018	9680	1239	413	570	2.3(10)-55	Ralstonia eutropha	P23516	high-affinity nickel transport protein.
CONTIG504	4863165_f3_175	4019	9681	1599	533	2152	5.4(10)-223	Escherichia coli	b2709	[pn:hypothetical sigma-54-dependent transcriptional regulator in gutq-hyph intergenic region] [gn:ygaal]
CONTIG504	10050718_c1_191	4020	9682	849	283	1166	1.6(10)-118	Escherichia coli	b2705	[pn:sorbitol-6-phosphate 2-dehydrogenase] [gn:srlb]
CONTIG504	4349193_c1_221	4021	9683	360	120	407	4.4(10)-38	Escherichia coli	b2726	[pn:hypa protein] [gn:hypa]
CONTIG504	33447142_c1_222	4022	9684	867	289	1153	3.8(10)-117	Escherichia coli	b2727	[pn:hydrogenase isoenzymes formation protein hypb] [gn:hypb]
CONTIG504	16252_c1_226	4023	9685	2109	703	2883	1.8(10)-300	Escherichia coli	b2731	[pn:transcriptional activator of the formate hydrogenlyase system] [gn:fhla]
CONTIG504	13672291_c1_235	4024	9686	330	110	92	0.00011	Escherichia coli	b4347	[pn:hypothetical 14.6 kd protein in merb-hsds intergenic region] [gn:yjiw]
CONTIG504	11992307_c2_247	4025	9687	990	330	1231	2.1(10)-125	Escherichia coli	b2703	[gn:srla_2]
CONTIG504	15728588_c2_248	4026	9688	375	125	475	2.7(10)-45	Escherichia coli	b2704	[pn:pts system, glucitol/sorbitol-specific iia component] [gn:srlb]
CONTIG504	25914811_c2_249	4027	9689	372	124	385	9.5(10)-36	Escherichia coli	b2706	[pn:gutm] [gn:gutm]
CONTIG504	25627000_c2_250	4028	9690	837	279	1202	2.5(10)-122	Escherichia coli	b2707	[pn:srlr] [gn:srlr]
CONTIG504	35739757_c2_258	4029	9691	1152	384	1525	1.5(10)-156	Escherichia coli	b2711	[pn:hypothetical protein in hyda 3''' region] [gn:ygbd]
CONTIG504	31894511_c2_262	4030	9692	342	114	103	6.5(10)-5	Plasmid pSB24.2	M32513	or: plasmid psb24.2 pn:neomycin resistance protein lc:1443 re:2756 di:direct sr:plasmid psb24.2 dna
CONTIG504	14259667_c2_293	4031	9693	1014	338	1359	5.7(10)-139	Escherichia coli	b2730	[pn:hydrogenase isoenzymes formation protein hype] [gn:hype]
CONTIG504	4572178_c2_298	4032	9694	765	255	409	2.7(10)-38	Escherichia coli	b4287	[pn:ironiii dicitrate transport atp-binding protein fece] [gn:fece]
CONTIG504	5195253_c2_299	4033	9695	1230	410	263	1.3(10)-22	Escherichia coli	b1065	[pn:hypothetical protein]
CONTIG504	22548313_c2_302	4034	9696	2583	861	3763	0	Escherichia coli	b2733	[pn:dna mismatch repair protein] [gn:muta]

CONTIG504	7156875_c2_303	4035	9697	264	88	99	0.00013	Salmonella typhi	Q56132	rna polymerase sigma factor rpos (sigma-38).
CONTIG504	24335932_c3_316	4036	9698	594	198	697	8.1(10)-69	Escherichia coli	b2702	[pn:pts system, glucitol/sorbitol-specific iibc component] [gn:srla_1]
CONTIG504	24495456_c3_320	4037	9699	978	326	1287	2.5(10)-131	Escherichia coli	b2708	[pn:gutq] [gn:gutq]
CONTIG504	15682878_c3_324	4038	9700	1461	487	2252	1.3(10)-233	Escherichia coli	b2710	[pn:hypothetical protein]
CONTIG504	11730162_c3_333	4039	9701	1497	499	1768	2.7(10)-182	Escherichia coli	b2715	[pn:phosphotransferase enzyme iiaabc asc]
CONTIG504	26054757_c3_334	4040	9702	1443	481	2295	3.7(10)-238	Escherichia coli	b2716	[pn:6-phospho-beta-glucosidase] [gn:ascb]
CONTIG504	22753805_c3_344	4041	9703	243	81	95	0.00033	Homo sapiens	U31468	[de:homo.sapiens homeobox protein (gbx2) gene, complete cds.] [pn:homeobox protein] [gn:gbx2]
CONTIG504	35282013_c3_346	4042	9704	342	114	409	2.7(10)-38	Escherichia coli	b2728	[pn:hydrogenase isoenzymes formation protein hycp] [gn:hycp]
CONTIG504	19790911_c3_347	4043	9705	1125	375	1709	4.7(10)-176	Escherichia coli	b2729	[pn:hydrogenase isoenzymes formation protein hypd] [gn:hypd]
CONTIG504	24646891_c3_352	4044	9706	1005	335	106	2.3(10)-8	Bacillus subtilis	yvrC	[pn:hypothetical protein]
CONTIG504	29869067_c3_353	4045	9707	747	249	268	2.3(10)-23	Bacillus subtilis	fhuB	[pn:ferrichrome abc transporter]
CONTIG504	167342_c3_369	4046	9708	435	145	128	1.6(10)-8	Bacillus subtilis	yybA	[pn:hypothetical protein]
CONTIG504	14181462_c3_372	4047	9709	1416	472	220	1.7(10)-15	Escherichia coli	b2367	[pn:multidrug resistance protein y] [gn:emry]
CONTIG505	3208137_fl_23	4048	9710	1215	405	1418	3.2(10)-145	Escherichia coli	b3124	[pn:hypothetical 42.1 kd protein in mpb-soha intergenic region] [gn:yhad]
CONTIG505	4082005_fl_33	4049	9711	1437	479	1983	4.4(10)-205	Escherichia coli	b3110	[pn:hypothetical 46.6 kd protein in exur-tdec intergenic region] [gn:yhao]
CONTIG505	34007882_fl_48	4050	9712	1515	505	2290	1.3(10)-237	Escherichia coli	b3091	[pn:altronate hydrolase] [gn:uxaa]
CONTIG505	282138_fl_57	4051	9713	1395	465	1887	6.5(10)-195	Escherichia coli	b3084	[pn:hypothetical 43.4 kd protein in ebgc-exut intergenic region] [gn:ygio]

CONTIG505	13144378_f1_66	4052	9714	938	313	602	9.5(10)-59	Escherichia coli	b1514	[pn:hypothetical protein]
CONTIG505	23712803_f2_76	4053	9715	774	258	1052	2.0(10)-106	Escherichia coli	b3126	[pn:hypothetical 27.4 kd protein in rnpb-soha intergenic region] [gn:yhaaf]
CONTIG505	34381931_f2_77	4054	9716	912	304	1312	5.5(10)-134	Escherichia coli	b3125	[pn:hypothetical 31.0 kd protein in rnpb-soha intergenic region] [gn:yhae]
CONTIG505	4104583_f2_80	4055	9717	933	311	1248	3.3(10)-127	Escherichia coli	b3118	[pn:bc operon transcriptional activator] [gn:tdca]
CONTIG505	24630001_f2_81	4056	9718	1029	343	1249	2.6(10)-127	Escherichia coli	b3117	[pn:catabolic threonine dehydratase] [gn:tdcb]
CONTIG505	24416526_f2_99	4057	9719	1509	503	2319	1.1(10)-240	Escherichia coli	b3092	[pn:uronate isomerase] [gn:uxac]
CONTIG505	35728280_f2_119	4058	9720	1575	525	1521	4.0(10)-156	Escherichia coli	b1513	[pn:hypothetical protein]
CONTIG505	4157318_f3_135	4059	9721	813	271	1216	8.3(10)-124	Escherichia coli	b3131	[pn:putative aga operon transcriptional repressor] [gn:agar]
CONTIG505	24103376_f3_140	4060	9722	1335	445	1803	5.2(10)-186	Escherichia coli	b3116	[pn:threonine-serine permease] [gn:tdcc]
CONTIG505	21617157_f3_141	4061	9723	1239	413	1388	4.9(10)-142	Escherichia coli	b3115	[pn:hypothetical protein in exur-tdcc intergenic region] [gn:yhaa]
CONTIG505	2151667_f3_142	4062	9724	2307	769	3452	0	Escherichia coli	b3114	[pn:probable formate acetyltransferase 3] [gn:yhas]
CONTIG505	16048967_f3_144	4063	9725	1398	466	720	3.0(10)-71	Escherichia coli	b3108	[pn:hypothetical 19.4 kd protein in exur-tdcc intergenic region] [gn:yham]
CONTIG505	32313842_f3_147	4064	9726	972	324	1419	2.6(10)-145	Escherichia coli	b3105	[pn:hypothetical transcriptional regulator in exur-tdcc intergenic region] [gn:yhai]
CONTIG505	24117711_c1_190	4065	9727	1611	537	2058	4.9(10)-213	Escherichia coli	b1511	[pn:hypothetical protein] [gn:ydev]
CONTIG505	5109457_c1_194	4066	9728	1005	335	1405	7.7(10)-144	Escherichia coli	b3087	[pn:ygir]
CONTIG505	14337766_c1_195	4067	9729	1245	415	1456	3.1(10)-149	Escherichia coli	b3089	[pn:hypothetical 43.5 kd protein in ebgc-exut intergenic region] [gn:ygu]
CONTIG505	6375378_c1_200	4068	9730	1347	449	1783	6.7(10)-184	Escherichia coli	b3093	[pn:hexuronate transporter] [gn:exut]

CONTIG505	14261275_c1_204	4069	9731	414	138	430	1.6(10)-40	Escherichia coli	b3097	[pn:hypothetical 14.5 kd protein in exur-tdcc intergenic region]
CONTIG505	26036653_c1_206	4070	9732	300	100	428	2.6(10)-40	Escherichia coli	b3100	[pn:hypothetical protein]
CONTIG505	23886067_c1_211	4071	9733	717	239	1030	4.2(10)-104	Escherichia coli	b3106	[pn:hypothetical 25.9 kd protein in exur-tdcc intergenic region] [gn:yhak]
CONTIG505	25915932_c1_227	4072	9734	1575	525	2421	1.7(10)-251	Escherichia coli	b3128	[pn:hypothetical 56.4 kd protein in mpb-soha intergenic region] [gn:yhag]
CONTIG505	12616452_c1_229	4073	9735	1326	442	1876	9.5(10)-194	Escherichia coli	b3132	[pn:putative tagatase 6-phosphate kinase agaz] [gn:agaz]
CONTIG505	29925955_c1_230	4074	9736	942	314	1003	3.1(10)-101	Vibrio fumiisii	U65015	or:vibrio fumiisii pn:pts permease for mannose subunit iibman gn:manz le:1604 re:2485 di:direct nt:manz
CONTIG505	10937880_c2_243	4075	9737	1050	350	996	1.7(10)-100	Escherichia coli	b1512	[pn:hypothetical protein]
CONTIG505	12733401_c2_245	4076	9738	2064	688	2834	2.8(10)-295	Escherichia coli	b3081	[pn:probable nadh-dependent flavin oxidoreductase] [gn:yglj]
CONTIG505	2238537_c2_259	4077	9739	519	173	403	1.2(10)-37	Escherichia coli	b3096	[pn:hypothetical 14.2 kd protein in exur-tdcc intergenic region]
CONTIG505	35759530_c2_260	4078	9740	471	157	282	7.7(10)-25	Escherichia coli	b3099	[pn:hypothetical 15.1 kd protein in exur-tdcc intergenic region]
CONTIG505	2424140_c2_261	4079	9741	414	138	486	1.8(10)-46	Escherichia coli	b3101	[pn:hypothetical 17.2 kd protein in exur-tdcc intergenic region] [gn:yqjf]
CONTIG505	13833558_c2_262	4080	9742	1059	353	1589	2.5(10)-163	Escherichia coli	b3102	[pn:hypothetical 37.4 kd protein in exur-tdcc intergenic region]
CONTIG505	24432937_c2_289	4081	9743	489	163	764	6.5(10)-76	Escherichia coli	b3133	[pn:pts system, n-acetylglucosamine-specific iib component 2] [gn:agav]
CONTIG505	3239382_c2_293	4082	9744	1176	392	704	1.5(10)-69	Escherichia coli	b3135	[pn:putative n-acetylglucosamine-6-phosphate deacetylase] [gn:agaa]
CONTIG505	22369003_c2_294	4083	9745	1146	382	1610	1.5(10)-165	Escherichia coli	b3136	[pn:protein] [gn:agas]
CONTIG505	3395463_c2_295	4084	9746	885	295	1348	8.5(10)-138	Escherichia coli	b3137	[pn:tagatase-bisphosphate aldolase] [gn:agay]
CONTIG505	2395842_c2_296	4085	9747	546	182	377	6.7(10)-35	Escherichia coli	b1621	[pn:pts system, maltose and glucose-specific ii abc component] [gn:malx]

CONTIG505	1270053_c3_309	4086	9748	537	179	770	1.5(10)-76	Escherichia coli	b3085	[pn:hypothetical 20.9 kd protein in ebgc-exut intergenic region] [gn:ygjp]
CONTIG505	1171956_c3_311	4087	9749	999	333	1142	5.7(10)-116	Escherichia coli	b3088	[pn:hypothetical 35.8 kd protein in ebgc-exut intergenic region]
CONTIG505	31284686_c3_322	4088	9750	810	270	1229	3.5(10)-125	Escherichia coli	b3094	[pn:exu regulon regulator] [gn:exur]
CONTIG505	23577_c3_324	4089	9751	663	221	883	1.6(10)-88	Escherichia coli	b3095	[pn:hypothetical protein] [gn:yqja]
CONTIG505	21570341_c3_326	4090	9752	330	110	345	1.6(10)-31	Escherichia coli	b3098	[pn:hypothetical 11.1 kd protein in exur-tdec intergenic region]
CONTIG505	34562826_c3_329	4091	9753	387	129	444	5.2(10)-42	Escherichia coli	b3103	[pn:hypothetical 14.3 kd protein in exur-tdec intergenic region] [gn:yhah]
CONTIG505	26667717_c3_338	4092	9754	864	288	578	3.3(10)-56	Vibrio fumiissii	U65015	or:vibrio fumiissii pn:pts permease for mannose subunit iipman gn:many le:838 re:1614 di:direct nt:many; pel; iidman
CONTIG505	2822002_c3_360	4093	9755	471	157	363	2.0(10)-33	Vibrio fumiissii	U65015	or:vibrio fumiissii pn:pts permease for mannose subunit iipman n gn:manw le:2543 re:2983 di:direct nt:manw; iipman
CONTIG506	33726666_fl_2	4094	9756	1842	614	3110	0	Escherichia coli	b2286	[pn:chain d] [gn:nadh dehydrogenase i chain c]
CONTIG506	36225631_fl_5	4095	9757	996	332	1393	1.5(10)-142	Escherichia coli	b2282	[pn:nadh dehydrogenase i chain h] [gn:nuoh]
CONTIG506	31697625_fl_6	4096	9758	693	231	589	2.2(10)-57	Escherichia coli	b2280	[pn:nadh dehydrogenase i chain j] [gn:nuoj]
CONTIG506	7242961_fl_22	4097	9759	1311	437	1167	1.3(10)-118	Escherichia coli	b2265	[pn:isochorismate synthase] [gn:menf]
CONTIG506	26302091_fl_23	4098	9760	822	274	904	9.5(10)-91	Escherichia coli	b2263	[pn:yfbb]
CONTIG506	25520283_fl_27	4099	9761	1617	539	1589	2.5(10)-163	Escherichia coli	b2260	[pn:o-succinylbenzoic acid-coa ligase] [gn:mene]
CONTIG506	22477163_fl_31	4100	9762	513	171	130	3.1(10)-8	Agrobacterium vitis	U32375	or:agrobacterium vitis pn:unknown le:10379 re:11221 di:complement sr:plasmid ptrab3
CONTIG506	2532787_fl_34	4101	9763	729	243	843	2.7(10)-84	Salmonella typhimurium	P22104	transcriptional regulatory protein tctd.

CONTIG506	22845841_f1_36	4102	9764	366	122	327	1.3(10)-29	Salmonella typhi	AF029846	[de:salmonella typhi tctd and tcte genes, complete cds.] [pn:tcte] [gn:tcte]
CONTIG506	822702_f1_37	4103	9765	672	224	710	3.5(10)-70	Escherichia coli	b2250	[pn:hypothetical protein]
CONTIG506	26676087_f1_45	4104	9766	1371	457	1962	7.2(10)-203	Escherichia coli	b2240	[pn:glycerol-3-phosphatase transporter] [gn:gpt]
CONTIG506	4181558_f2_57	4105	9767	444	148	700	4.0(10)-69	Escherichia coli	b2288	[pn:nadh dehydrogenase i chain a] [gn:nuoa]
CONTIG506	24117711_f2_58	4106	9768	690	230	1142	5.7(10)-116	Escherichia coli	b2287	[pn:nadh dehydrogenase i chain b] [gn:nuob]
CONTIG506	16688506_f2_61	4107	9769	1356	452	2268	2.7(10)-235	Escherichia coli	b2284	[pn:nadh dehydrogenase i chain f] [gn:nuof]
CONTIG506	9822541_f2_62	4108	9770	2760	920	4322	0	Escherichia coli	b2283	[gn:nuog]
CONTIG506	5322878_f2_66	4109	9771	1851	617	2758	3.2(10)-287	Escherichia coli	b2278	[pn:nadh dehydrogenase i chain l] [gn:nuol]
CONTIG506	1207277_f2_67	4110	9772	1536	512	2262	1.2(10)-234	Escherichia coli	b2277	[pn:nadh dehydrogenase i chain m] [gn:nuom]
CONTIG506	14879150_f2_68	4111	9773	1464	488	1829	9.0(10)-189	Escherichia coli	b2276	[pn:nadh dehydrogenase i chain n] [gn:nuon]
CONTIG506	29900800_f2_72	4112	9774	339	113	279	1.6(10)-24	Escherichia coli	b2266	[pn:hypothetical protein] [gn:elab]
CONTIG506	13859718_f2_74	4113	9775	1767	589	2446	3.7(10)-254	Escherichia coli	b2264	[pn:2-succinyl-6-hydroxy-2,4-cyclohexadiene-1-carboxylate synthase] [gn:mend]
CONTIG506	15801416_f2_76	4114	9776	1068	356	1342	3.7(10)-137	Escherichia coli	b2261	[pn:o-succinylbenzoate-coa synthase] [gn:menc]
CONTIG506	16929206_f2_89	4115	9777	1122	374	1024	1.8(10)-103	Salmonella typhi	AF029846	[de:salmonella typhi tctd and tcte genes, complete cds.] [pn:tcte] [gn:tcte]
CONTIG506	30275056_f2_91	4116	9778	1221	407	1644	3.7(10)-169	Escherichia coli	b2249	[pn:hypothetical protein]
CONTIG506	2552202_f2_109	4117	9779	2637	879	3740	0	Escherichia coli	b2231	[pn:dna gyrase subunit a] [gn:gyra]
CONTIG506	32680156_f2_110	4118	9780	2871	957	3706	0	Escherichia coli	b2218	[pn:sensor protein rcsc] [gn:rcsc]
CONTIG506	29305135_f3_117	4119	9781	537	179	844	2.2(10)-84	Escherichia coli	b2285	[pn:nadh dehydrogenase i chain e] [gn:nuoc]



CONTIG506	16605040_f3_122	4120	9782	423	141	374	1.3(10)-34	Escherichia coli	D90859	or:escherichia coli pn:nadh dehydrogenase i chain g (ec 1.6.5.3) gn:nuog le:9026 re:9565 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [swissprot accession number [pn:nadh dehydrogenase i chain i] [gn:nuoi] [pn:nadh dehydrogenase i chain k] [gn:nuok] [pn:hypothetical protein] [gn:elaa]
CONTIG506	6049181_f3_126	4121	9783	651	217	897	5.2(10)-90	Escherichia coli	b2281	
CONTIG506	26041441_f3_128	4122	9784	336	112	319	9.4(10)-29	Escherichia coli	b2279	
CONTIG506	5129635_f3_134	4123	9785	498	166	630	1.0(10)-61	Escherichia coli	b2267	
CONTIG506	14459708_f3_136	4124	9786	207	69	144	3.2(10)-10	Escherichia coli	D90857	or:escherichia coli pn:mend protein gn:mend le:10698 re:10907 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [pir accession number a33860] [pn:naphthoate synthase] [gn:menb]
CONTIG506	11067041_f3_137	4125	9787	969	323	1438	2.5(10)-147	Escherichia coli	b2262	
CONTIG506	3375126_f3_138	4126	9788	1191	397	1545	1.1(10)-158	Escherichia coli	b2239	[pn:glycerophosphoryl diester phosphodiesterase periplasmic precursor] [gn:g pq]
CONTIG506	24494150_c1_169	4127	9789	741	247	1051	2.5(10)-106	Escherichia coli	b2217	[pn:regulator of capsule synthesis b component] [gn:rcsb]
CONTIG506	20605418_c1_178	4128	9790	2328	776	3678	0	Escherichia coli	b2234	[pn:ribonucleoside-diphosphate reductase 1 alpha chain] [gn:nrda]
CONTIG506	14156906_c1_179	4129	9791	1221	407	1844	2.2(10)-190	Escherichia coli	b2235	[pn:ribonucleoside-diphosphate reductase 1 beta chain] [gn:nrdb]
CONTIG506	22869676_c1_197	4130	9792	1530	510	766	4.0(10)-76	Agrobacterium vitis	U32375	[PN:unknown] [SR:plasmid pTrAB3] [DE:Agrobacterium vitis plasmid pTrAB3 tartrate utilization generegion, including LysR-like regulator (tuuA), membrane protein(tuuB), tartrate dehydrogenase (tuuC and tuuC"), enzyme degradingprimary ta

CONTIG506	33464808_c1_211	4131	9793	1050	350	1143	4.5(10)-116	Escherichia coli	b2268	[pn:hypothetical protein] [gn:elac]
CONTIG506	16600443_c2_242	4132	9794	744	248	1177	1.1(10)-119	Escherichia coli	b2232	[pn:3-demethylubiquinone-9 3-methyltransferase] [gn:ubig]
CONTIG506	14469587_c2_244	4133	9795	201	67	96	0.00076	Escherichia coli	K02672	or:escherichia coli gn:nrdh le:3505 re:5835 di:direct sr:escherichia coli k-12 dna, clone pps2 nt:ribonucleoside diphosphate reductase b1 subunit
CONTIG506	34470790_c2_245	4134	9796	288	96	183	3.7(10)-13	Escherichia coli	K02672	or:escherichia coli gn:nrdh le:3505 re:5835 di:direct sr:escherichia coli k-12 dna, clone pps2 nt:ribonucleoside diphosphate reductase b1 subunit
CONTIG506	24412702_c2_250	4135	9797	1659	553	2303	5.4(10)-239	Escherichia coli	b2241	[pn:anaerobic glycerol-3-phosphate dehydrogenase subunit a] [gn:glpa]
CONTIG506	32031937_c2_252	4136	9798	1320	440	1716	8.5(10)-177	Escherichia coli	b2243	[pn:anaerobic glycerol-3-phosphate dehydrogenase subunit c] [gn:glpc]
CONTIG506	4566712_c2_274	4137	9799	1032	344	334	2.3(10)-30	Helicobacter pylori	HP0393	[pn:chemotaxis protein chev] [gn:chev]
CONTIG506	29956660_c3_303	4138	9800	546	182	755	5.9(10)-75	Escherichia coli	b2216	[pn:probable sensor protein yojn] [gn:yojn]
CONTIG506	7267876_c3_313	4139	9801	1053	351	223	2.6(10)-16	Mycobacterium smegmatis	X84077	or:mycobacterium smegmatis gn:orf617 le:3075 re:4925 di:complement nt:val start codon
CONTIG506	22067707_c3_314	4140	9802	1095	365	108	0.00619	Mycobacterium smegmatis	X84077	or:mycobacterium smegmatis gn:orf617 le:3075 re:4925 di:complement nt:val start codon
CONTIG506	31820142_c3_321	4141	9803	291	97	383	1.5(10)-35	Escherichia coli	b2236	[pn:hypothetical 9.3 kd protein in nrdh 5' region] [gn:yfae]
CONTIG506	35286582_c3_328	4142	9804	1329	443	1061	2.2(10)-107	Escherichia coli	b2242	[pn:anaerobic glycerol-3-phosphate dehydrogenase subunit b] [gn:glpb]
CONTIG506	24415957_c3_336	4143	9805	984	328	232	1.6(10)-19	Bacillus subtilis	yflP	[pn:hypothetical protein]
CONTIG507	26738791_fl_3	4144	9806	768	256	146	6.4(10)-9	Haemophilus influenzae	HI0575	[pn:gb]
CONTIG507	4693763_fl_5	4145	9807	348	116	180	5.0(10)-14	Bacillus subtilis	yvaE	[pn:hypothetical protein]
CONTIG507	913562_fl_6	4146	9808	2463	821	835	2.0(10)-83	Escherichia coli	b3951	[pn:formate acetyltransferase 2] [gn:pflD]

CONTIG507	4584643_fl_7	4147	9809	234	78	208	5.4(10)-17	Escherichia coli	b2833	[pn:hypothetical protein]
CONTIG507	7320391_fl_11	4148	9810	1149	383	428	2.6(10)-40	Escherichia coli	b1486	[pn:hypothetical protein]
CONTIG507	16916450_fl_12	4149	9811	966	322	489	9.0(10)-47	Haemophilus influenzae	H11185	[pn:dipeptide transport atp-binding protein] [gn:dppd]
CONTIG507	25391038_fl_18	4150	9812	894	298	949	1.6(10)-95	Klebsiella terrigena	P52666	bud operon transcriptional regulator.
CONTIG507	33724033_fl_25	4151	9813	495	165	95	0.0061	Saccharomyces cerevisiae	YDL037C	[pn:strong similarity to glucan 1,4-alpha-glucosidase]
CONTIG507	24416090_f2_71	4152	9814	807	269	183	7.0(10)-13	Escherichia coli	b1464	[pn:hypothetical 32.3 kd protein in rhse-narv intergenic region] [gn:ydde]
CONTIG507	30745308_f2_78	4153	9815	975	325	437	2.8(10)-41	Escherichia coli	b0824	[pn:hypothetical protein]
CONTIG507	24001537_f2_81	4154	9816	1602	534	436	3.7(10)-41	Escherichia coli	b1487	[pn:hypothetical protein]
CONTIG507	7300265_f2_87	4155	9817	864	288	477	1.7(10)-45	Bacillus subtilis	ykd	[pn:hypothetical protein]
CONTIG507	33806966_f2_96	4156	9818	273	91	270	1.5(10)-23	Escherichia coli	b0581	[pn:hypothetical protein] [gn:ybdk]
CONTIG507	11209657_f2_97	4157	9819	537	179	233	1.2(10)-19	Vibrio cholerae	S81006	or:vibrio cholerae pn:hcp gn:hcp le:690 re:1208 di:direct sr:vibrio cholerae o17 nt:28 kda secreted hydrophilic protein; this sequence
CONTIG507	16884836_f2_117	4158	9820	753	251	129	1.8(10)-8	Saccharomyces cerevisiae	X85757	or:saccharomyces cerevisiae pn:unknown gn:internal orf g1669 le:6964 re:7365 di:direct sr:baker's yeast
CONTIG507	1300636_f2_126	4159	9821	663	221	106	0.00051	Haemophilus influenzae	H10955	[pn:protein homolog] [gn:tlk]
CONTIG507	6923416_f2_130	4160	9822	1452	484	301	5.7(10)-25	Bacillus subtilis	mmr	[pn:methylenomycin a resistance protein]
CONTIG507	4141926_f3_151	4161	9823	429	143	146	2.0(10)-10	Escherichia coli	b0543	[pn:hypothetical protein] [gn:emre]
CONTIG507	12754381_f3_159	4162	9824	879	293	495	2.1(10)-47	Escherichia coli	b1485	[pn:hypothetical protein]
CONTIG507	26679086_f3_180	4163	9825	1140	380	1556	7.7(10)-160	Escherichia coli	b0581	[pn:hypothetical protein] [gn:ybdk]

CONTIG507	23479076_f3_190	4164	9826	267	89	178	8.1(10)-14	Escherichia coli	b0580	[pn:hypothetical protein] [gn:ybdj]
CONTIG507	26063750_f3_193	4165	9827	2181	727	1814	3.5(10)-187	Erwinia chrysanthemi	Q47162	ferrichrysobactin receptor precursor.
CONTIG507	10007212_f3_195	4166	9828	579	193	397	5.0(10)-37	Escherichia coli	b0579	[pn:hypothetical protein in nfsb 5'''region] [gn:ybdj]
CONTIG507	10206325_f3_196	4167	9829	774	258	1084	8.0(10)-110	Enterobacter cloacae	Q01234	oxygen-insensitive nad(p)h nitroreductase (ec 1.-.-.-).
CONTIG507	4723752_f3_199	4168	9830	1302	434	1644	3.7(10)-169	Escherichia coli	b0577	[pn:hypothetical protein in phep-entd intergenic region] [gn:ybdg]
CONTIG507	1197041_f3_212	4169	9831	786	262	125	3.2(10)-5	Escherichia coli	b0457	[pn:hypothetical protein] [gn:ylab]
CONTIG507	34065925_c1_214	4170	9832	297	99	265	4.9(10)-23	Escherichia coli	b0329	[pn:hypothetical protein] [gn:yaho]
CONTIG507	16148586_c1_217	4171	9833	1986	662	96	0.0038	Pseudomonas putida	S64724	uxpb protein - pseudomonas putida (fragment)
CONTIG507	26369168_c1_218	4172	9834	1392	464	1906	6.2(10)-197	Escherichia coli	b0576	[pn:phenylalanine-specific permease] [gn:phep]
CONTIG507	109705_c1_220	4173	9835	1296	432	669	7.5(10)-66	Escherichia coli	b2098	[pn:hypothetical protein]
CONTIG507	36225641_c1_221	4174	9836	1041	347	155	7.9(10)-9	Bacillus subtilis	iolE	[pn:hypothetical protein] [gn:yxde]
CONTIG507	14582660_c1_234	4175	9837	420	140	499	7.9(10)-48	Klebsiella pneumoniae	Q48413	transcriptional activator rama.
CONTIG507	13867217_c1_241	4176	9838	2721	907	1360	4.5(10)-139	Bacillus subtilis	yloB	[pn:hypothetical protein]
CONTIG507	14570343_c1_242	4177	9839	1092	364	480	8.0(10)-46	Escherichia coli	b2074	[pn:hypothetical protein]
CONTIG507	4332318_c1_243	4178	9840	3165	1055	807	1.8(10)-80	Escherichia coli	b0462	[pn:acriflavin resistance protein b] [gn:acrb]
CONTIG507	7083293_c1_247	4179	9841	306	102	185	1.5(10)-14	Mycobacterium tuberculosis	Q50648	hypothetical 26.2 kd protein cy227.28c.
CONTIG507	25474036_c1_248	4180	9842	723	241	906	5.9(10)-91	Haemophilus influenzae	HI0882	[pn:hypothetical protein]
CONTIG507	24620302_c1_252	4181	9843	804	268	1353	2.5(10)-138	Enterobacter aerogenes	P05361	alpha-acetolactate decarboxylase (ec 4.1.1.5).

CONTIG507	25516656_c1_256	4182	9844	834	278	1194	1.8(10)-121	Klebsiella pneumoniae	D86412	or:klebsiella pneumoniae pn:meso-2,3-butanediol dehydrogenase (d-acetoin gn:budc le:319 re:1089 di:direct sr:klebsiella pneumoniae (strain:iam1063) dna
CONTIG507	12994562_c2_290	4183	9845	258	86	93	0.0015	Bacteriophage I3	S26427	structural protein, 70k - phage I3
CONTIG507	5212842_c2_298	4184	9846	1245	415	237	9.0(10)-18	Bacillus subtilis	yfil	[pn:hypothetical protein]
CONTIG507	33485627_c2_299	4185	9847	1020	340	427	3.3(10)-40	Escherichia coli	b3753	[pn:rbs repressor] [gn:rbsr]
CONTIG507	33786037_c2_305	4186	9848	309	103	110	1.3(10)-6	Mycobacterium tuberculosis	Z84498	or:mycobacterium tuberculosis pn:unknown gn:mtey09f9.08c le:3775 re:4224 di:complement nt:mtey09f9.08c, unknown, len: 149 aa
CONTIG507	973760_c2_314	4187	9849	1632	544	252	2.2(10)-19	Escherichia coli	b0585	[pn:enterochelin esterase] [gn:fes]
CONTIG507	25478406_c2_320	4188	9850	894	298	660	6.9(10)-65	Mycobacterium tuberculosis	Q50648	hypothetical 26.2 kd protein cy227.28c.
CONTIG507	12347825_c2_327	4189	9851	1779	593	1407	4.7(10)-144	Bacillus subtilis	alsS	[pn:alpha-acetolactate synthase]
CONTIG507	12992781_c2_328	4190	9852	240	80	120	2.7(10)-7	Klebsiella terrigena	Q04520	acetoin(diacetyl) reductase (ec 1.1.1.5) (acetoin dehydrogenase (ar).
CONTIG507	2035278_c2_335	4191	9853	765	255	437	2.8(10)-41	Saccharomyces cerevisiae	P40586	hypothetical 27.4 kd protein in hyr1 3'region.
CONTIG507	3940838_c3_348	4192	9854	2193	731	1970	1.0(10)-203	Yersinia enterocolitica	Q05202	ferrichrome receptor fcuA precursor.
CONTIG507	14510461_c3_392	4193	9855	489	163	102	9.3(10)-6	Staphylococcus aureus	AF003592	[PN:CspB] [GN:cspB] [DE:Staphylococcus aureus CspB (cspB) gene, complete cds.] [NT:similar to major cold-shock protein] [LE:439] [RE:639] [DI:direct]
CONTIG508	33992790_f1_9	4194	9856	930	310	1402	1.6(10)-143	Escherichia coli	b2916	[pn:chromosome initiation inhibitor] [gn:icia]
CONTIG508	16285416_f1_24	4195	9857	237	79	105	4.5(10)-6	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:car le:59 re:580 di:direct nt:orf2

CONTIG508	31770765_fl_36	4196	9858	369	123	93	0.003	human herpesvirus 2	Z86099	or: human herpesvirus 2 gn:rs1 le: 128076 re: 132032 di: complement [pn:endonuclease i] [gn:enda]
CONTIG508	36438950_fl_52	4197	9859	765	255	1142	5.7(10)-116	Escherichia coli	b2945	[pn:hypothetical protein in endo- gshb intergenic region] [gn:yggj]
CONTIG508	994036_fl_53	4198	9860	750	250	948	2.1(10)-95	Escherichia coli	b2946	[pn:glutathione synthetase] [gn:gshb]
CONTIG508	24807956_fl_54	4199	9861	960	320	1498	1.1(10)-153	Escherichia coli	b2947	[pn:hypothetical protein in pilt region (orf6)]
CONTIG508	35661425_fl_57	4200	9862	318	106	138	1.3(10)-9	Pseudomonas aeruginosa	P24564	[pn:hypothetical protein] [gn:yggv]
CONTIG508	15751010_fl_60	4201	9863	327	109	416	4.9(10)-39	Escherichia coli	b2953	[pn:hypothetical protein] [gn:yggv]
CONTIG508	24790907_fl_61	4202	9864	597	199	910	2.2(10)-91	Escherichia coli	b2954	[pn:hypothetical protein] [gn:yggv]
CONTIG508	29976593_fl_63	4203	9865	387	129	292	6.7(10)-26	Escherichia coli	b1289	[pn:hypothetical 14.0 kd protein in envm-sapf intergenic region] [gn:ycjd]
CONTIG508	10755300_fl_66	4204	9866	1071	357	1684	2.1(10)-173	Escherichia coli	b2961	[pn:a/g-specific adenine glycosylase] [gn:muty]
CONTIG508	10003757_fl_67	4205	9867	1260	420	1648	1.3(10)-169	Escherichia coli	b2963	[pn:yggz] [gn:mltc]
CONTIG508	20353462_fl_81	4206	9868	1683	561	835	2.0(10)-83	Escherichia coli	b0619	[pn:hypothetical protein] [gn:cita]
CONTIG508	10940963_f2_96	4207	9869	639	213	126	3.5(10)-8	Saccharomyce s cerevisiae	P25614	very hypothetical 22.8 kd protein in pgk1 region.
CONTIG508	3925052_f2_102	4208	9870	846	282	1050	3.2(10)-106	Escherichia coli	b2936	[pn:hypothetical 31.8 kd protein in tkta-speb intergenic region] [gn:yggg]
CONTIG508	13675250_f2_123	4209	9871	1314	438	157	1.3(10)-8	Escherichia coli	b4356	[pn:hypothetical 49.4 kd protein in tsr-mdob intergenic region]
CONTIG508	16151390_f2_124	4210	9872	696	232	726	7.0(10)-72	Escherichia coli	M32363	or: escherichia coli le: 271 re: 1287 di: complement sr: escherichia coli (strain k-12) (clone: plc[2-5,5-8,5- 14]) dna nt: orf1; putative
CONTIG508	14493937_f2_127	4211	9873	1425	475	2144	3.7(10)-222	Escherichia coli	b2943	[pn:galactose-proton symport] [gn:galp]
CONTIG508	22078760_f2_132	4212	9874	453	151	659	8.8(10)-65	Escherichia coli	b2949	[pn:hypothetical protein] [gn:yqgf]
CONTIG508	6775790_f2_133	4213	9875	420	140	118	1.7(10)-6	Volvox carteri	S22697	extensin - volvox carteri (fragment)

CONTIG508	31344632_f2_134	4214	9876	717	239	855	1.5(10)-85	Escherichia coli	b2951	[pn:hypothetical protein] [gn:yggs]
CONTIG508	34395318_f2_135	4215	9877	585	195	797	2.1(10)-79	Escherichia coli	b2952	[pn:hypothetical protein] [gn:yggt]
CONTIG508	29845662_f2_136	4216	9878	1173	391	1799	1.3(10)-185	Escherichia coli	b2955	[pn:hypothetical protein] [gn:ygww]
CONTIG508	2911266_f2_143	4217	9879	1344	448	1883	1.7(10)-194	Escherichia coli	b2964	[pn:nucleoside permease nupg]
CONTIG508	4039075_f2_156	4218	9880	828	276	415	6.2(10)-39	Escherichia coli	b0620	[gn:nupg]
CONTIG508	489515_f3_159	4219	9881	396	132	486	1.8(10)-46	Escherichia coli	b2910	[pn:transcriptional regulator crir]
CONTIG508	25985662_f3_160	4220	9882	693	231	748	3.2(10)-74	Escherichia coli	b2912	[gn:citb]
CONTIG508	13073275_f3_184	4221	9883	1041	347	159	2.5(10)-11	Azospirillum brasilense	X70360	[pn:hypothetical 12.7 kd protein in pepp-ssr intergenic region] [gn:ygfe]
CONTIG508	24353412_f3_204	4222	9884	1158	386	1813	4.5(10)-187	Escherichia coli	b2942	[pn:hypothetical 21.1 kd protein in ssr-sera intergenic region] [gn:ygfa]
CONTIG508	14957708_f3_206	4223	9885	570	190	796	2.7(10)-79	Escherichia coli	b2944	or:azospirillum brasilense gn:carr le:<1 re:588 di:direct
CONTIG508	35820762_f3_209	4224	9886	708	236	894	1.1(10)-89	Escherichia coli	b2948	[pn:s-adenosylmethionine synthetase] [gn:metk]
CONTIG508	14583376_f3_218	4225	9887	381	127	436	3.7(10)-41	Escherichia coli	b2962	[pn:hypothetical 19.3 kd protein in galp-enda intergenic region]
CONTIG508	4879557_c1_240	4226	9888	1389	463	960	1.1(10)-96	Bacillus subtilis	yxkJ	[pn:hypothetical protein]
CONTIG508	4567918_c1_242	4227	9889	1032	344	728	4.2(10)-72	Escherichia coli	b0618	[pn:hypothetical protein]
CONTIG508	21661682_c1_243	4228	9890	903	301	935	5.0(10)-94	Escherichia coli	b0616	[pn:hypothetical protein] [gn:cite]
CONTIG508	22520128_c1_247	4229	9891	621	207	285	3.7(10)-25	Escherichia coli	b0614	[pn:hypothetical protein]
CONTIG508	4730001_c1_259	4230	9892	771	257	1178	8.8(10)-120	Escherichia coli	b2960	[pn:hypothetical 27.3 kd protein in muty 5''' region] [gn:yggh]
CONTIG508	1382785_c1_281	4231	9893	1998	666	3068	0	Escherichia coli	b2938	[pn:biosynthetic arginine decarboxylase] [gn:spea]

CONTIG508	24015955_c1_284	4232	9894	438	146	324	2.7(10)-29	Acinetobacter calcoaceticus	AF009672	[PN:unknown] [DE:Acinetobacter calcoaceticus ADP1 vanillate demethylase region, vanillate demethylase (vanB) and vanillate demethylase (vanA) genes, complete cds.] [NT:putative ferredoxin; ORF9] [LE:12779] [RE:13084] [DI:direct]
CONTIG508	16261457_c1_285	4233	9895	1116	372	1157	1.5(10)-117	Acinetobacter calcoaceticus	AF009672	[PN:unknown] [DE:Acinetobacter calcoaceticus ADP1 vanillate demethylase region, vanillate demethylase (vanB) and vanillate demethylase (vanA) genes, complete cds.] [NT:similar to vanillate demethylase (vanA subunit);] [LE:13411] [RE:143]
CONTIG508	25500413_c1_293	4234	9896	552	184	90	0.01799	Bacillus subtilis	ykrZ	[pn:hypothetical protein]
CONTIG508	5114843_c1_294	4235	9897	780	260	120	2.3(10)-5	Bacillus subtilis	yvaM	[pn:hypothetical protein]
CONTIG508	34491313_c1_307	4236	9898	1185	395	1788	2.0(10)-184	Escherichia coli	b2926	[pn:phosphoglycerate kinase] [gn:pgk]
CONTIG508	2914202_c1_309	4237	9899	855	285	1025	1.3(10)-103	Escherichia coli	b2924	[pn:hypothetical 30.9 kd protein in sbm-fba intergenic region] [gn:yggb]
CONTIG508	14462638_c1_312	4238	9900	783	261	780	1.3(10)-77	Escherichia coli	b2922	[pn:hypothetical 26.6 kd protein in sbm-fba intergenic region] [gn:ygge]
CONTIG508	4165941_c1_318	4239	9901	384	128	548	5.0(10)-53	Escherichia coli	b2909	[pn:hypothetical 21.5 kd protein in pepp-ssr intergenic region] [gn:ygfb]
CONTIG508	15128755_c2_328	4240	9902	600	200	194	1.6(10)-15	Haemophilus influenzae	H10024	[pn:acyl lyase subunit] [gn:citd]
CONTIG508	22917825_c2_330	4241	9903	1668	556	1715	1.1(10)-176	Escherichia coli	b0615	[pn:hypothetical protein] [gn:citf]
CONTIG508	35275330_c2_332	4242	9904	807	269	449	1.6(10)-42	Escherichia coli	b0613	[pn:hypothetical protein] [gn:citg]
CONTIG508	6072130_c2_342	4243	9905	984	328	1234	1.0(10)-125	Escherichia coli	b2950	[pn:hypothetical protein]
CONTIG508	36113775_c2_351	4244	9906	243	81	161	5.2(10)-12	Escherichia coli	b2941	[pn:hypothetical 9.5 kd protein in spea-metk intergenic region] [gn:yqgd]



CONTIG508	3261557_c2_352	4245	9907	219	73	125	3.3(10)-8	Escherichia coli	b2939	[pn:hypothetical 5.4 kd protein in spea-metk intergenic region] [gn:yqgb]
CONTIG508	16195806_c2_367	4246	9908	1227	409	467	1.8(10)-44	Escherichia coli	b2542	[pn:hypothetical protein]
CONTIG508	11750417_c2_372	4247	9909	987	329	248	3.1(10)-21	Escherichia coli	b1619	[pn:7-alpha-hydroxysteroid dehydrogenase] [gn:hdha]
CONTIG508	4332693_c2_373	4248	9910	780	260	297	2.0(10)-26	Methanobacterium thermoautotrophicum	MTH973	[pn:conserved protein]
CONTIG508	5197318_c2_374	4249	9911	1479	493	931	1.3(10)-93	Saccharomyces cerevisiae	YER073W	[pn:aldehyde dehydrogenase] [gn:ald3]
CONTIG508	13756568_c2_377	4250	9912	2184	728	3323	0	Escherichia coli	b2935	[pn:transketolase] [gn:tkta]
CONTIG508	876260_c2_381	4251	9913	708	236	1009	7.0(10)-102	Escherichia coli	b2923	[pn:hypothetical protein in sbm-fba intergenic region] [gn:ygga]
CONTIG508	34394790_c2_384	4252	9914	696	232	1048	5.2(10)-106	Escherichia coli	b2914	[pn:ribose 5-phosphate isomerase] [gn:rpia]
CONTIG508	7226077_c2_385	4253	9915	1233	411	1936	4.2(10)-200	Escherichia coli	b2913	[pn:d-3-phosphoglycerate dehydrogenase] [gn:sera]
CONTIG508	32713250_c3_388	4254	9916	1122	374	1722	2.0(10)-177	Escherichia coli	b2965	[pn:ornithine decarboxylase, constitutive] [gn:spec]
CONTIG508	2128266_c3_394	4255	9917	924	308	396	6.5(10)-37	Bacillus subtilis	yisK	[pn:hypothetical protein]
CONTIG508	13135432_c3_411	4256	9918	387	129	571	1.8(10)-55	Escherichia coli	b2959	[pn:hypothetical protein in muty 5''' region] [gn:ygg]
CONTIG508	35745292_c3_412	4257	9919	771	257	942	9.0(10)-95	Escherichia coli	b2958	[pn:hypothetical protein in ansb 5''' region] [gn:ygg]
CONTIG508	597140_c3_430	4258	9920	948	316	1550	3.2(10)-159	Escherichia coli	b2937	[pn:agmatinase] [gn:speb]
CONTIG508	36069667_c3_435	4259	9921	771	257	300	9.6(10)-27	Bacillus subtilis	fabG	[pn:3-oxoacyl-acyl-carrier protein reductase] [gn:yplf]
CONTIG508	5214643_c3_436	4260	9922	801	267	326	1.7(10)-29	Escherichia coli	b0272	[pn:hypothetical transcriptional regulator in per-argf intergenic region] [gn:yagi]
CONTIG508	17048951_c3_437	4261	9923	978	326	177	1.8(10)-11	Ralstonia eutropha	P17296	metapyrocatechase 2 (ec 1.13.11.2) (cat02ase) (catechol 2,3-dioxygenase ii).

CONTIG508	6351031_c3_440	4262	9924	1278	426	247	1.1(10)-18	Escherichia coli	b4356	[pn:hypothetical 49.4 kd protein in tsr-mdb intergenic region]
CONTIG508	26056558_c3_445	4263	9925	1644	548	374	2.2(10)-34	Escherichia coli	b3671	[pn:acetohydroxy acid synthase i, small subunit] [gn:ilvb]
CONTIG508	34645626_c3_447	4264	9926	1035	345	1663	3.6(10)-171	Escherichia coli	b2927	[pn:d-erythrose 4-phosphate dehydrogenase] [gn:epd]
CONTIG508	22147313_c3_448	4265	9927	1179	393	1761	1.5(10)-181	Escherichia coli	b2925	[pn:fructose 1,6-bisphosphate aldolase] [gn:fba]
CONTIG509	10335127_fl_3	4266	9928	1932	644	2853	2.7(10)-297	Escherichia coli	b3846	[pn:large]
CONTIG509	15702_fl_18	4267	9929	1044	348	1362	2.7(10)-139	Escherichia coli	b3566	[pn:xylf] [gn:xylf]
CONTIG509	16536530_fl_19	4268	9930	1614	538	2319	1.1(10)-240	Escherichia coli	b3567	[pn:d-xylose transport atp-binding protein xylg] [gn:xylg]
CONTIG509	14156906_fl_32	4269	9931	1584	528	1928	2.8(10)-199	Escherichia coli	b3580	[pn:cryptic l-xylose kinase] [gn:lyx]
CONTIG509	15822667_fl_35	4270	9932	723	241	1080	2.1(10)-109	Escherichia coli	b3583	[pn:hypothetical 25.6 kd protein in avta-selb intergenic region] [gn:yias]
CONTIG509	22166282_fl_49	4271	9933	1953	651	2652	5.5(10)-276	Escherichia coli	b3599	[pn:mannitol-specific enzyme ii of phosphotransferase system] [gn:mtla]
CONTIG509	15656306_fl_56	4272	9934	1776	592	2236	6.7(10)-232	Escherichia coli	b3603	[pn:l-lactate permease] [gn:ldp]
CONTIG509	17058340_f2_68	4273	9935	525	175	595	5.2(10)-58	Escherichia coli	b3846	[pn:large]
CONTIG509	4800443_f2_69	4274	9936	1173	391	1796	2.8(10)-185	Escherichia coli	b3845	[pn:small] [gn:fada]
CONTIG509	19582686_f2_76	4275	9937	1140	380	1383	1.7(10)-141	Escherichia coli	b3561	[pn:hypothetical 37.6 kd protein in glyq-xyfb intergenic region]
CONTIG509	16213568_f2_82	4276	9938	1194	398	1570	2.5(10)-161	Escherichia coli	b3568	[pn:xylose transport permease protein xylh] [gn:xylh]
CONTIG509	25400283_f2_83	4277	9939	1182	394	1762	1.1(10)-181	Escherichia coli	b3569	[pn:xylose operon regulatory protein] [gn:xylr]
CONTIG509	14069843_f2_87	4278	9940	1398	466	1914	9.0(10)-198	Escherichia coli	b3572	[pn:valine--pyruvate aminotransferase] [gn:avta]
CONTIG509	34645312_f2_91	4279	9941	1017	339	1599	2.2(10)-164	Escherichia coli	b3575	[pn:hypothetical protein] [gn:yiak]
CONTIG509	178442_f2_93	4280	9942	1359	453	464	4.0(10)-44	Escherichia coli	b2246	[pn:hypothetical protein]

CONTIG509	36066655_f2_97	4281	9943	921	307	1257	3.7(10)-128	Escherichia coli	b3582	[pn:hypothetical 33.7 kd protein in avta-selb intergenic region] [gn:yiar]
CONTIG509	251562_f2_100	4282	9944	1428	476	752	1.2(10)-74	Escherichia coli	b3657	[pn:hypothetical 51.0 kd protein in glts-selc intergenic region] [gn:yicj]
CONTIG509	35783391_f2_112	4283	9945	630	210	795	3.3(10)-79	Escherichia coli	b3601	[pn:mannotol operon repressor] [gn:mtlr]
CONTIG509	36503135_f2_123	4284	9946	951	317	1316	2.1(10)-134	Escherichia coli	b3605	[pn:lctd] [gn:lldd]
CONTIG509	188326_f3_148	4285	9947	2154	718	2820	8.8(10)-294	Escherichia coli	b3571	[pn:alpha-amylase] [gn:mals]
CONTIG509	7317883_f3_150	4286	9948	315	105	103	7.2(10)-6	Escherichia coli	P37670	hypothetical 11.9 kd protein in avta-selb intergenic region (o103).
CONTIG509	21932205_f3_152	4287	9949	663	221	673	2.8(10)-66	Escherichia coli	b3576	[pn:hypothetical 17.5 kd protein in avta-selb intergenic region] [gn:yial]
CONTIG509	12756693_f3_155	4288	9950	666	222	949	1.6(10)-95	Escherichia coli	b3581	[pn:hypothetical 23.4 kd protein in avta-selb intergenic region] [gn:yiaq]
CONTIG509	911390_f3_161	4289	9951	2007	669	172	9.0(10)-10	Bacteroides ovatus	U15179	or:bacteroides ovatus le:<1 re:1311 di:direct nt:orf1
CONTIG509	29767327_f3_162	4290	9952	492	164	207	6.9(10)-17	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:59 re:580 di:direct nt:orf2
CONTIG509	24020078_f3_163	4291	9953	1032	344	228	8.5(10)-19	Haemophilus influenzae	H10182	[pn:hypothetical protein]
CONTIG509	7147186_f3_174	4292	9954	1257	419	1561	2.2(10)-160	Escherichia coli	b3600	[pn:mannotol-1-phosphate dehydrogenase] [gn:mtld]
CONTIG509	260937_f3_176	4293	9955	441	147	496	1.6(10)-47	Escherichia coli	b3602	[pn:hypothetical 13.7 kd protein in mtlr-lctp intergenic region] [gn:yibl]
CONTIG509	25495182_f3_186	4294	9956	849	283	1100	1.6(10)-111	Escherichia coli	b3604	[pn:lctr] [gn:lldr]
CONTIG509	3160052_c1_222	4295	9957	825	275	139	2.2(10)-7	Escherichia coli	b3906	[pn:l-rhamnose operon transcriptional activator] [gn:rhar]
CONTIG509	4692912_c1_237	4296	9958	267	89	96	0.00011	Escherichia coli	A27279	valine--pyruvate transaminase (ec 2.6.1.66) - escherichiacoli
CONTIG509	2081888_c1_242	4297	9959	891	297	1215	1.1(10)-123	Escherichia coli	b3570	[pn:hypothetical protein] [gn:bax]
CONTIG509	16220188_c1_265	4298	9960	645	215	828	1.1(10)-82	Escherichia coli	b3848	[pn:hypothetical 21.9 kd protein in pepq-trkh intergenic region]
CONTIG509	30116455_c1_267	4299	9961	663	221	662	4.2(10)-65	Escherichia coli	b3850	[pn:hypothetical protein] [gn:heng]

CONTIG509	36338375_c2_275	4300	9962	606	202	184	1.8(10)-14	Escherichia coli	b0534	[pn:hypothetical fimbrial-like protein in fimz 5''' region] [gn:ybeg]
CONTIG509	21723250_c2_276	4301	9963	705	235	409	2.7(10)-38	Escherichia coli	b3143	[pn:hypothetical 25.7 kd fimbrial chaperone in agai- mtr intergeni] [gn:yrai]
CONTIG509	2931558_c2_286	4302	9964	612	204	834	2.5(10)-83	Escherichia coli	b3592	[pn:hypothetical protein] [gn:yibf]
CONTIG509	9822892_c2_288	4303	9965	1866	622	2493	4.0(10)-259	Escherichia coli	b3590	[pn:selb]
CONTIG509	2939513_c2_307	4304	9966	879	293	1274	5.9(10)-130	Escherichia coli	b3574	[pn:hypothetical transcriptional regulator in avta- selb intergenic region] [gn:yiaj]
CONTIG509	22072201_c2_308	4305	9967	459	153	517	9.8(10)-50	Escherichia coli	b3573	[pn:hypothetical 17.5 kd protein in avta-selb intergenic region]
CONTIG509	913427_c2_325	4306	9968	1518	506	1955	4.0(10)-202	Escherichia coli	b3564	[pn:xylulose kinase] [gn:xylb]
CONTIG509	20817037_c2_326	4307	9969	939	313	1494	2.8(10)-153	Escherichia coli	b3560	[pn:glycine-trna synthetase, alpha subunit] [gn:glyq]
CONTIG509	4432338_c2_327	4308	9970	2079	693	3204	0	Escherichia coli	b3559	[pn:glycine-trna synthetase, beta subunit] [gn:glys]
CONTIG509	23886001_c2_333	4309	9971	1362	454	2160	7.7(10)-224	Escherichia coli	b3847	[pn:xaa-pro dipeptidase] [gn:pepq]
CONTIG509	3377258_c3_345	4310	9972	2544	848	1465	3.3(10)-150	Escherichia coli	b3144	[pn:hypothetical outer membrane usher protein in agal-mtr intergenic region] [gn:yraj]
CONTIG509	4329717_c3_346	4311	9973	1104	368	144	1.5(10)-9	Escherichia coli	b4318	[pn:fimf protein precursor] [gn:fimf]
CONTIG509	32292527_c3_353	4312	9974	1644	548	1778	2.2(10)-183	Escherichia coli	b3591	[pn:l-seryl-trna] [gn:sela]
CONTIG509	4744642_c3_357	4313	9975	1641	547	2423	1.0(10)-251	Escherichia coli	b3588	[pn:aldehyde dehydrogenase b] [gn:aldb]
CONTIG509	24744803_c3_379	4314	9976	267	89	120	2.2(10)-7	Escherichia coli	A27279	valine-pyruvate transaminase (ec 2.6.1.66) - escherichiacoli
CONTIG509	650966_c3_380	4315	9977	369	123	182	3.1(10)-14	Escherichia coli	A27279	valine-pyruvate transaminase (ec 2.6.1.66) - escherichiacoli
CONTIG509	4739012_c3_392	4316	9978	1410	470	2209	4.9(10)-229	Escherichia coli	b3565	[pn:d-xylose isomerase] [gn:xyla]
CONTIG509	26750802_c3_408	4317	9979	1473	491	1941	1.2(10)-200	Escherichia coli	b3849	[pn:trkh]

CONTIG51	4892893_c1_3	4318	9980	297	99	239	2.7(10)-20	Escherichia coli	b1825	[pn:hypothetical protein]
CONTIG510	22735807_fl_2	4319	9981	516	172	424	7.0(10)-40	Escherichia coli	b3929	[pn:menaquinone biosynthesis protein meng] [gn:meng]
CONTIG510	23634682_fl_25	4320	9982	702	234	977	1.8(10)-98	Escherichia coli	b3912	[pn:transcriptional regulatory protein] [gn:cpxr]
CONTIG510	16015963_fl_28	4321	9983	1020	340	270	1.5(10)-23	Escherichia coli	M85158	or:escherichia coli gn:soda le:<1 re:225 di:direct sr:escherichia coli (strain k-12) (library: lambda from kohara et al.
CONTIG510	23835200_fl_29	4322	9984	1149	383	1420	2.0(10)-145	Escherichia coli	b3907	[pn:rhamnose permease] [gn:rhat]
CONTIG510	7212781_fl_59	4323	9985	591	197	914	8.3(10)-92	Escherichia coli	b3894	[pn:formate dehydrogenase-o, major subunit] [gn:fdog]
CONTIG510	14555411_fl_60	4324	9986	2463	821	3796	0	Escherichia coli	b3894	[pn:formate dehydrogenase-o, major subunit] [gn:fdog]
CONTIG510	2992841_fl_61	4325	9987	915	305	1496	1.8(10)-153	Escherichia coli	b3893	[pn:formate dehydrogenase-o, iron-sulfur subunit] [gn:fdoh]
CONTIG510	5197193_fl_72	4326	9988	879	293	1225	9.1(10)-125	Escherichia coli	b3881	[pn:hypothetical 32 kd protein in glna-fdhe intergenic region]
CONTIG510	55413_fl_75	4327	9989	2037	679	3041	0	Escherichia coli	b3878	[pn:hypothetical 77.2 kd protein in glna-fdhe intergenic region]
CONTIG510	16119418_fl_76	4328	9990	1425	475	2125	3.8(10)-220	Escherichia coli	b3877	[pn:hypothetical 51.7 kd protein in glna-fdhe intergenic region]
CONTIG510	33862917_fl_80	4329	9991	783	261	840	5.7(10)-84	Escherichia coli	b3875	[pn:hypothetical protein]
CONTIG510	19782067_fl_85	4330	9992	1455	485	2353	2.7(10)-244	Escherichia coli	b3870	[pn:glutamine synthetase] [gn:glna]
CONTIG510	14875927_fl_92	4331	9993	927	309	1244	8.9(10)-127	Escherichia coli	b3930	[pn:menaquinone biosynthesis protein mena] [gn:mena]
CONTIG510	412557_fl_95	4332	9994	873	291	1060	2.7(10)-107	Escherichia coli	b3927	[pn:glycerol uptake facilitator protein] [gn:glpf]
CONTIG510	35707880_fl_96	4333	9995	1530	510	2456	3.2(10)-255	Escherichia coli	b3926	[pn:glycerol kinase] [gn:glpk]
CONTIG510	20910700_fl_97	4334	9996	1026	342	1516	1.3(10)-155	Escherichia coli	b3925	[pn:glpx protein] [gn:glpx]
CONTIG510	14664712_fl_98	4335	9997	774	258	1179	6.9(10)-120	Escherichia coli	b3924	[pn:ferredoxin--nadp reductase] [gn:fpr]

CONTIG510	32422943_f2_118	4336	9998	1272	424	1966	2.7(10)-203	Escherichia coli	b3903	[pn:l-rhamnose isomerase] [gn:rhaa]
CONTIG510	12975937_f2_119	4337	9999	927	309	1331	5.4(10)-136	Escherichia coli	b3902	[pn:rhamnulose-1-phosphate aldolase] [gn:rhad]
CONTIG510	35444752_f2_125	4338	10000	1272	424	391	2.2(10)-36	Bacillus subtilis	rbsC	[pn:ribose abc transporter]
CONTIG510	16666393_f2_127	4339	10001	342	114	507	1.1(10)-48	Escherichia coli	b3901	[pn:hypothetical 12.3 kd protein in rhad 3'''region] [gn:yiii]
CONTIG510	31283452_f2_146	4340	10002	930	310	1369	5.0(10)-140	Escherichia coli	b3891	[pn:fdhe protein] [gn:fdhe]
CONTIG510	892542_f2_147	4341	10003	954	318	329	8.1(10)-30	Escherichia coli	b0476	[pn:hypothetical protein in hemh-gsk intergenic region] [gn:ybac]
CONTIG510	36214211_f2_151	4342	10004	903	301	1165	2.1(10)-118	Escherichia coli	b3882	[pn:hypothetical 31.2 kd protein in glna-fdhe intergenic region]
CONTIG510	4416318_f2_158	4343	10005	1434	478	1907	4.9(10)-197	Escherichia coli	b3876	[pn:hypothetical protein]
CONTIG510	7225312_f2_163	4344	10006	1431	477	2103	8.4(10)-218	Escherichia coli	b3868	[pn:gng] [gn:gng]
CONTIG510	2166665_f3_165	4345	10007	303	101	413	1.0(10)-38	Escherichia coli	b3931	[pn:heat shock protein hslu] [gn:hslu]
CONTIG510	17085216_f3_174	4346	10008	615	205	666	1.6(10)-65	Escherichia coli	b3920	[pn:hypothetical 21.8 kd protein in tpia 3'''region precursor] [gn:yitq]
CONTIG510	4964458_f3_175	4347	10009	783	261	1130	1.1(10)-114	Escherichia coli	b3919	[pn:triosephosphate isomerase] [gn:tpia]
CONTIG510	10636442_f3_188	4348	10010	1434	478	1806	2.5(10)-186	Escherichia coli	b3911	[pn:cpxa] [gn:cpxa]
CONTIG510	33796877_f3_194	4349	10011	1707	569	2017	1.1(10)-208	Escherichia coli	b3904	[pn:rhamnulokinase] [gn:rhab]
CONTIG510	23939402_f3_198	4350	10012	1083	361	491	5.5(10)-47	Escherichia coli	b1516	[pn:hypothetical protein]
CONTIG510	34632782_f3_199	4351	10013	1521	507	1076	5.7(10)-109	Escherichia coli	b3749	[pn:high affinity ribose transport protein] [gn:rbsa]
CONTIG510	10422077_f3_200	4352	10014	1005	335	414	8.0(10)-39	Escherichia coli	b3750	[pn:high affinity ribose transport protein] [gn:rbsc]
CONTIG510	4964675_f3_202	4353	10015	1164	388	850	5.0(10)-85	Escherichia coli	b3589	[pn:hypothetical 40.2 kd protein in avta-selb intergenic region] [gn:yiaa]

CONTIG510	24642887_f3_211	4354	10016	666	222	172	3.5(10)-13	Streptomyces coelicolor	AL020958	[de:streptomyces coelicolor cosmid 4h8.] [pn:hypothetical protein sc4h8.02] [gn:sc4h8.02] [nt:sc4h8.02, possible membrane [pn:formate dehydrogenase, cytochrome b556] [gn:fdot] [pn:hypothetical 48 kd protein in glna-fdhe intergenic region] [pn:gln] [gn:gln]
CONTIG510	32304817_f3_221	4355	10017	657	219	999	8.1(10)-101	Escherichia coli	b3892	[pn:formate dehydrogenase, cytochrome b556] [gn:fdot] [pn:hypothetical 48 kd protein in glna-fdhe intergenic region] [pn:gln] [gn:gln]
CONTIG510	32283263_f3_230	4356	10018	1287	429	1880	3.6(10)-194	Escherichia coli	b3880	[pn:hypothetical 48 kd protein in glna-fdhe intergenic region] [pn:gln] [gn:gln]
CONTIG510	35272952_f3_245	4357	10019	1050	350	1609	1.8(10)-165	Escherichia coli	b3869	[pn:hypothetical 48 kd protein in glna-fdhe intergenic region] [pn:gln] [gn:gln]
CONTIG510	7166507_c1_252	4358	10020	1485	495	2205	1.3(10)-228	Escherichia coli	b3867	[pn:oxyen-independent coproporphyrinogen iii oxidase] [gn:hemn]
CONTIG510	36203141_c1_256	4359	10021	969	323	311	1.3(10)-27	Transposon mini-Tn3Cm	M84113	or:transposon mini-tn3cm le:30 re:1508 di:complement sr:transposon mini-tn3cm dna; neisseria gonorrhoeae (strain ms11) dna nt:orf1
CONTIG510	31847192_c1_257	4360	10022	570	190	257	1.5(10)-21	Transposon mini-Tn3Cm	M84113	or:transposon mini-tn3cm le:30 re:1508 di:complement sr:transposon mini-tn3cm dna; neisseria gonorrhoeae (strain ms11) dna nt:orf1
CONTIG510	1964051_c1_271	4361	10023	654	218	813	4.2(10)-81	Escherichia coli	b3885	[pn:hypothetical 23.5 kd protein in glna-fdhe intergenic region] [pn:hypothetical 37.1 kd protein in glna-fdhe intergenic region] or:enterobacter cloacae pn:mely gn:mely le:481 re:1758 di:direct sr:enterobacter cloacae (strain:iid977) dna
CONTIG510	12978333_c1_273	4362	10024	1092	364	1480	8.8(10)-152	Escherichia coli	b3888	[pn:hypothetical 23.5 kd protein in glna-fdhe intergenic region] [pn:hypothetical 37.1 kd protein in glna-fdhe intergenic region] or:enterobacter cloacae pn:mely gn:mely le:481 re:1758 di:direct sr:enterobacter cloacae (strain:iid977) dna
CONTIG510	15829202_c1_289	4363	10025	1401	467	1892	1.8(10)-195	Enterobacter cloacae	AB000622	[pn:hypothetical 23.5 kd protein in glna-fdhe intergenic region] [pn:hypothetical 37.1 kd protein in glna-fdhe intergenic region] or:enterobacter cloacae pn:mely gn:mely le:481 re:1758 di:direct sr:enterobacter cloacae (strain:iid977) dna
CONTIG510	3963265_c1_303	4364	10026	879	293	1050	3.2(10)-106	Escherichia coli	b3906	[pn:l-rhamnose operon transcriptional activator] [gn:rhar] [pn:manganese superoxide dismutase] [gn:soda]
CONTIG510	31884586_c1_305	4365	10027	654	218	1074	9.1(10)-109	Escherichia coli	b3908	[pn:manganese superoxide dismutase] [gn:soda]
CONTIG510	1026_c1_306	4366	10028	705	235	966	2.6(10)-97	Escherichia coli	b3910	[pn:hypothetical 26.6 kd protein in kdgt-cpxa intergenic region] [gn:yim]

CONTIG510	5910626_c1_308	4367	10029	603	201	491	5.5(10)-47	Escherichia coli	b3914	[pn:hypothetical 14.4 kd protein in cpxa-pfka intergenic region]
CONTIG510	4579541_c1_309	4368	10030	909	303	1289	1.5(10)-131	Escherichia coli	b3915	[pn:hypothetical 32.9 kd protein in cpxa-pfka intergenic region]
CONTIG510	14119712_c2_326	4369	10031	282	94	198	6.2(10)-16	Escherichia coli	b3866	[pn:hypothetical 19.1 kd protein in pola-henn intergenic region]
CONTIG510	25671932_c2_346	4370	10032	948	316	1040	3.7(10)-105	Escherichia coli	b3883	[pn:hypothetical 31.9 kd protein in glna-fdhe intergenic region]
CONTIG510	34100937_c2_351	4371	10033	462	154	698	6.4(10)-69	Escherichia coli	b3887	[pn:hypothetical 15.9 kd protein in glna-fdhe intergenic region]
CONTIG510	16688543_c2_366	4372	10034	1062	354	1031	3.2(10)-104	Escherichia coli	A35160	repressor protein rafr - escherichia coli
CONTIG510	24426430_c2_367	4373	10035	2229	743	2766	4.7(10)-288	Escherichia coli	P16551	alpha-galactosidase (ec 3.2.1.22) (melibiase).
CONTIG510	4410252_c2_395	4374	10036	1095	365	1462	7.0(10)-150	Escherichia coli	b3916	[pn:6-phosphofructokinase] [gn:pfka]
CONTIG510	33260055_c2_396	4375	10037	1050	350	1469	1.3(10)-150	Escherichia coli	b3917	[pn:periplasmic sulphate binding protein] [gn:sbp]
CONTIG510	31894063_c2_397	4376	10038	819	273	941	1.1(10)-94	Escherichia coli	b3918	[pn:cdp-diglyceride hydrolase] [gn:cdh]
CONTIG510	5914043_c3_418	4377	10039	1833	611	2757	4.2(10)-287	Escherichia coli	b3871	[pn:65.4 kd gtp-binding protein in glna-fdhe intergenic region]
CONTIG510	34163512_c3_431	4378	10040	912	304	1066	6.5(10)-108	Escherichia coli	b3884	[pn:hypothetical transcriptional regulator in glna-fdhe intergenic region] [gn:yihw]
CONTIG510	20969180_c3_432	4379	10041	951	317	997	1.3(10)-100	Escherichia coli	b3886	[pn:hypothetical 32.8 kd protein in glna-fdhe intergenic region]
CONTIG510	30100280_c3_448	4380	10042	912	304	1110	1.3(10)-112	Escherichia coli	b3895	[pn:fdhd protein] [gn:fdhd]
CONTIG510	818893_c3_449	4381	10043	279	93	97	3.1(10)-5	Escherichia coli	b0802	[pn:hypothetical 8.6 kd protein in ding/rarb 3'''region] [gn:ybij]
CONTIG510	14704431_c3_452	4382	10044	642	214	168	9.4(10)-13	Escherichia coli	b1434	[pn:hypothetical protein]
CONTIG510	10156438_c3_483	4383	10045	864	288	1052	2.0(10)-106	Escherichia coli	b3905	[pn:l-rhamnose operon regulatory protein rhas] [gn:rhas]
CONTIG510	22281562_c3_494	4384	10046	1326	442	204	9.0(10)-14	Methanobacterium thermoautotrophicum	MTH788	[pn:sodium/dicarboxylate or sulfate cotransporter]



CONTIG510	20570300_c3_498	4385	10047	501	167	546	8.3(10)-53	Escherichia coli	b3921	[pn:hypothetical 16.5 kd protein in tpa-fpr intergenic region] [gn:yir]
CONTIG510	519167_c3_509	4386	10048	273	91	362	2.6(10)-33	Escherichia coli	b3928	[pn:hypothetical 9.6 kd protein in gpt-hslu intergenic region] [gn:yiu]
CONTIG511	31835915_f1_2	4387	10049	3219	1073	2383	1.8(10)-247	Burkholderia cepacia	U97042	ceob, ceob, similar to cytoplasmic membrane proteins of the rnd
CONTIG511	17070152_f1_9	4388	10050	855	285	159	4.7(10)-10	Escherichia coli	b1782	[pn:hypothetical protein]
CONTIG511	29320338_f1_13	4389	10051	903	301	316	1.8(10)-28	Escherichia coli	b2409	[pn:hypothetical protein]
CONTIG511	26772336_f1_30	4390	10052	1308	436	1961	9.4(10)-203	Escherichia coli	b2497	[pn:uracil permease] [gn:uraa]
CONTIG511	31647632_f1_31	4391	10053	744	248	1118	2.0(10)-113	Escherichia coli	b2496	[pn:hypothetical protein]
CONTIG511	4335802_f1_48	4392	10054	744	248	1185	1.6(10)-120	Escherichia coli	b2476	[pn:phosphoribosylaminoimidazole-succinocarboxamide synthase] [gn:purc]
CONTIG511	1229775_f1_49	4393	10055	882	294	1223	1.5(10)-124	Escherichia coli	b2475	[pn:hypothetical protein]
CONTIG511	900837_f1_52	4394	10056	804	268	1024	1.8(10)-103	Escherichia coli	b2473	[pn:hypothetical protein] [gn:ypfh]
CONTIG511	31509432_f1_64	4395	10057	2058	686	2749	2.8(10)-286	Escherichia coli	b2468	[pn:hypothetical 71.8 kd protein in tkb-narq intergenic region] [gn:yffg]
CONTIG511	25648456_f1_72	4396	10058	2304	768	3460	0	Escherichia coli	b2463	[pn:hypothetical protein]
CONTIG511	23954386_f1_74	4397	10059	525	175	707	7.2(10)-70	Escherichia coli	b2434	[pn:hypothetical protein]
CONTIG511	4772706_f1_77	4398	10060	948	316	1451	1.0(10)-148	Escherichia coli	b2431	[pn:hypothetical protein]
CONTIG511	15634678_f1_79	4399	10061	837	279	1110	1.3(10)-112	Escherichia coli	b2424	[pn:sulfate transport system permease protein cyst] [gn:cysu]
CONTIG511	24878442_f1_80	4400	10062	1131	377	1677	1.2(10)-172	Escherichia coli	b2422	[pn:sulfate transport atp-binding protein cysa] [gn:cysa]
CONTIG511	792317_f1_81	4401	10063	873	291	986	2.0(10)-99	Escherichia coli	b2418	[pn:hypothetical protein] [gn:pdck]

CONTIG511	10035957_f1_83	4402	10064	282	94	92	0.00169	Homo sapiens	U78554	[PN:mucin] [GN:MUC5B] [SR:human] [DE:Homo sapiens mucin (MUC5B) gene, 3 exons, partial cds.] [LE:U78552<1:U78553:266:U78553:733] [RE:526:447:904] [DI:directJoin]
CONTIG511	4866582_f2_85	4403	10065	1275	425	437	2.8(10)-41	Escherichia coli	b0463	[pn:acriflavin resistance protein a precursor] [gn:acra]
CONTIG511	4551528_f2_96	4404	10066	2379	793	2181	4.5(10)-226	Escherichia coli	b2503	[pn:hypothetical protein]
CONTIG511	36025316_f2_102	4405	10067	1434	478	1667	1.3(10)-171	Escherichia coli	b2901	[pn:6-phospho-beta-glucosidase bgla] [gn:bgla]
CONTIG511	22355137_f2_111	4406	10068	891	297	153	6.7(10)-9	Shewanella sp. SCRC-2738	U73935	[de:shewanella sp. scrc-2738 eicosapentaenoic acid (epa) synthesis genecluster, complete sequence.] [pn:unknown] [nt:orf4]
CONTIG511	10000668_f2_120	4407	10069	906	302	1390	3.0(10)-142	Escherichia coli	b2478	[pn:dihydrodipicolinate synthase] [gn:dapa]
CONTIG511	261305_f2_139	4408	10070	291	97	117	3.5(10)-6	Escherichia coli	L34011	or:escherichia coli pn:fe-s center and glutamate synthase (gltD) gn:aeg53.0 le:239 re:2197 di:direct sr:escherichia coli (individual isolate mc4100, strain k-12) (library)
CONTIG511	864182_f2_140	4409	10071	621	207	795	3.3(10)-79	Escherichia coli	b2467	[pn:hypothetical 21.7 kd protein in tkfb-narq intergenic region] [gn:yffh]
CONTIG511	24651580_f2_152	4410	10072	456	152	531	3.2(10)-51	Escherichia coli	b2433	[pn:hypothetical protein]
CONTIG511	114787_f2_153	4411	10073	639	213	700	4.0(10)-69	Escherichia coli	b2432	[pn:hypothetical protein]
CONTIG511	4817593_f2_155	4412	10074	1023	341	1482	5.4(10)-152	Escherichia coli	b2425	[pn:thiosulfate-binding protein precursor] [gn:cysp]
CONTIG511	6289053_f2_160	4413	10075	921	307	1385	1.0(10)-141	Escherichia coli	b2421	[pn:cysteine synthase b] [gn:cysm]
CONTIG511	6025325_f3_197	4414	10076	723	241	1036	9.8(10)-105	Escherichia coli	b2498	[pn:uracil phosphoribosyltransferase] [gn:upp]
CONTIG511	14070177_f3_210	4415	10077	1113	371	1299	1.3(10)-132	Escherichia coli	b2493	[pn:hypothetical protein] [gn:perm]

CONTIG511	5079818_f3_212	4416	10078	1044	348	1537	8.0(10)-158	Escherichia coli	b2477	[pn:lipoprotein-34 precursor] [gn:nlpb]
CONTIG511	22906410_f3_214	4417	10079	1968	656	2029	5.7(10)-210	Escherichia coli	b2474	[pn:hypothetical protein] [gn:ypfi]
CONTIG511	5173152_f3_229	4418	10080	1134	378	1351	4.0(10)-138	Escherichia coli	b2466	[pn:hypothetical protein]
CONTIG511	24344411_f3_243	4419	10081	918	306	1325	2.2(10)-135	Escherichia coli	P16702	sulfate transport system permease protein cysw.
CONTIG511	30603817_c1_249	4420	10082	456	152	719	3.7(10)-71	Escherichia coli	b2416	[pn:phosphoenolpyruvate-protein phosphotransferase] [gn:ptsi]
CONTIG511	2598451_c1_280	4421	10083	387	129	1115	3.8(10)-7	Escherichia coli	P37670	hypothetical 11.9 kd protein in avta-selb intergenic region (o103).
CONTIG511	36130181_c1_297	4422	10084	591	197	853	2.3(10)-85	Escherichia coli	b2479	[pn:gcvr]
CONTIG511	16540667_c1_302	4423	10085	462	154	488	1.2(10)-46	Escherichia coli	b2495	[pn:hypothetical protein]
CONTIG511	4492202_c1_313	4424	10086	696	232	935	5.0(10)-94	Escherichia coli	b2500	[pn:phosphoribosylglycinamide myltransferase] [gn:purn]
CONTIG511	10268800_c1_324	4425	10087	537	179	104	3.2(10)-5	Bacillus subtilis	yhdE	[pn:hypothetical protein]
CONTIG511	4339088_c2_334	4426	10088	519	173	577	4.2(10)-56	Haemophilus influenzae	HI1711	[pn:glucose phosphotransferase enzyme iii-glc] [gn:crr]
CONTIG511	16289055_c2_349	4427	10089	927	309	1468	1.6(10)-150	Escherichia coli	b2436	[pn:coproporphyrinogen iii oxidase] [gn:hemf]
CONTIG511	156276_c2_355	4428	10090	975	325	1380	3.5(10)-141	Escherichia coli	b2464	[pn:hypothetical protein] [gn:tala]
CONTIG511	34176402_c2_371	4429	10091	384	128	480	8.0(10)-46	Escherichia coli	b2471	[pn:hypothetical 13.6 kd protein in dape 5'''region] [gn:yftb]
CONTIG511	30165880_c2_372	4430	10092	1131	377	1862	2.8(10)-192	Escherichia coli	b2472	[pn:succinyl-diaminopimelate desuccinylase] [gn:dape]
CONTIG511	29900316_c2_373	4431	10093	201	67	192	2.7(10)-15	Escherichia coli	D90875	or:escherichia coli le:15855 re:16055 di:direct sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [pir accession number d42959]
CONTIG511	5109818_c2_392	4432	10094	1428	476	614	5.0(10)-60	Haemophilus influenzae	HI1705	[pn:aminopeptidase a/i] [gn:pepa]
CONTIG511	23947283_c2_399	4433	10095	1041	347	1603	8.0(10)-165	Escherichia coli	b2499	[pn:phosphoribosylformylglycinamide cyclo-ligase] [gn:purm]

CONTIG511	36366008_c2_402_4434	10096	2067	689	3137	0	Escherichia coli	b2501	[pn:polyphosphate kinase] [gn:ppk]
CONTIG511	4790881_c2_403_4435	10097	1542	514	2265	5.7(10)-235	Escherichia coli	b2502	[pn:exopolyphosphatase] [gn:ppx]
CONTIG511	12360706_c2_408_4436	10098	225	75	189	5.5(10)-15	Escherichia coli	b2504	[pn:hypothetical protein]
CONTIG511	26600312_c2_410_4437	10099	1236	412	91	0.28	Escherichia coli	b0045	[pn:hypothetical metabolite transport protein in carb-kefc intergenic region] [gn:yauu]
CONTIG511	31752035_c2_413_4438	10100	1407	469	296	1.5(10)-24	Escherichia coli	b2078	[pn:sensor protein baes] [gn:baes]
CONTIG511	14120812_c2_423_4439	10101	321	107	237	4.5(10)-20	Escherichia coli	b1531	[pn:multiple antibiotic resistance protein] [gn:marA]
CONTIG511	12323286_c3_437_4440	10102	909	303	1262	1.1(10)-128	Escherichia coli	b2435	[pn:probable n-acetylmuramoyl-L-alanine amidase] [gn:amia]
CONTIG511	31728382_c3_444_4441	10103	1992	664	3150	0	Escherichia coli	b2465	[pn:transketolase 2] [gn:tktb]
CONTIG511	23488453_c3_449_4442	10104	1695	565	1836	1.7(10)-189	Escherichia coli	b2469	[pn:nitrate/nitrite sensor protein] [gn:narq]
CONTIG511	24351016_c3_450_4443	10105	3141	1047	4723	0	Escherichia coli	b2470	[pn:acriflavin resistance protein d] [gn:acrd]
CONTIG511	12306553_c3_464_4444	10106	474	158	808	1.3(10)-80	Escherichia coli	b2480	[pn:bacterioferritin comigratory protein] [gn:bep]
CONTIG511	24397576_c3_467_4445	10107	1632	544	2010	6.0(10)-208	Escherichia coli	b2494	[pn:hypothetical protein]
CONTIG511	12578456_c3_471_4446	10108	1407	469	846	1.3(10)-84	Bacillus subtilis	ywbA	[pn:hypothetical protein] [gn:ipa-16d]
CONTIG511	36572167_c3_472_4447	10109	960	320	287	2.2(10)-25	Escherichia coli	b0828	[pn:hypothetical protein in moea-grxa intergenic region] [gn:ybk]
CONTIG511	2035_c3_492_4448	10110	729	243	375	1.1(10)-34	Haemophilus influenzae	H11708	[pn:transcriptional regulatory protein]
CONTIG512	32516252_fl_1_4449	10111	999	333	1465	3.3(10)-150	Escherichia coli	b2615	[pn:hypothetical 32.6 kd protein in grpe-recn intergenic region]
CONTIG512	2431957_fl_15_4450	10112	1290	430	146	1.3(10)-7	Bacillus subtilis	yxIH	[pn:hypothetical protein]
CONTIG512	11722677_fl_16_4451	10113	1383	461	656	1.8(10)-64	Methanobacterium thermoautotrophicum	MTH225	[pn:histidinol dehydrogenase]

CONTIG512	3244033_fl_17	4452	10114	768	256	302	5.9(10)-27	Escherichia coli	b2842	[pn:2-deoxy-d-gluconate 3-dehydrogenase] [gn:kdud]
CONTIG512	4884682_fl_37	4453	10115	921	307	431	1.3(10)-40	Escherichia coli	b0900	[pn:hypothetical protein] [gn:ycan]
CONTIG512	4567957_fl_45	4454	10116	309	103	101	0.00017	Nicotiana tabacum	P13983	extensin precursor (cell wall hydroxyproline-rich glycoprotein).
CONTIG512	13022331_fl_51	4455	10117	237	79	99	1.8(10)-5	Drosophila melanogaster	X13625	or:drosophila melanogaster sp.p13008 le:58 re:398 di:direct sr:fruit fly nt:put. ribosomal protein
CONTIG512	4507693_fl_54	4456	10118	1413	471	166	2.5(10)-9	Escherichia coli	b3035	[pn:tolc] [gn:tolc]
CONTIG512	24323967_fl_56	4457	10119	342	114	102	2.2(10)-5	Orf virus	C34768	orf2 protein - orf virus (strain nz2)
CONTIG512	6382182_fl_58	4458	10120	1251	417	293	1.3(10)-25	Bordetella pertussis	P11091	cyad protein.
CONTIG512	10428883_fl_94	4459	10121	1782	594	2335	2.2(10)-242	Escherichia coli	b2616	[pn:recl] [gn:recl]
CONTIG512	29854158_fl_95	4460	10122	378	126	244	8.3(10)-21	Haemophilus influenzae	H10838	[pn:hypothetical protein]
CONTIG512	16145763_fl_114	4461	10123	834	278	302	5.9(10)-27	Escherichia coli	b2245	[pn:hypothetical protein]
CONTIG512	24847006_fl_122	4462	10124	1950	650	265	2.0(10)-27	Escherichia coli	b3533	[pn:hypothetical 101.6 kd protein in delta-dppf intergenic region]
CONTIG512	32457178_fl_127	4463	10125	2727	909	520	2.6(10)-63	Escherichia coli	b2786	[pn:sensor protein bara] [gn:bara]
CONTIG512	15822183_fl_131	4464	10126	468	156	108	6.0(10)-5	Helicobacter pylori	HP0357	[pn:short chain alcohol dehydrogenase]
CONTIG512	22738760_fl_156	4465	10127	387	129	225	8.5(10)-19	Haemophilus influenzae	H11250	[pn:hypothetical protein]
CONTIG512	12925816_fl_164	4466	10128	276	92	100	0.00033	Caenorhabditis elegans	Z81518	[de:caenorhabditis elegans cosmid f28d9, complete sequence.] [pn:f28d9.a] [nt:protein predicted using genefinder; preliminary]
CONTIG512	16103408_fl_166	4467	10129	924	308	385	9.5(10)-36	Escherichia coli	b0254	[pn:peroxide resistance protein perr] [gn:perr]
CONTIG512	10025462_fl_170	4468	10130	891	297	95	0.016	Plasmodium falciparum	M13987	or:plasmodium falciparum pn:histidine-rich protein le:<23 re:679 di:direct sr:p.falciparum (strain 7g8) dna, clone pfhrp-iii

CONTIG512	33723390_f2_178	4469	10131	984	328	1370	4.0(10)-140	Escherichia coli	b0316	[pn:hypothetical protein] [gn:yahb]
CONTIG512	13713887_f3_187	4470	10132	513	171	768	2.5(10)-76	Escherichia coli	b2620	[pn:small protein b] [gn:snpb]
CONTIG512	14164827_f3_188	4471	10133	588	196	395	8.3(10)-37	Escherichia coli	b2622	[pn:slpa integrase] [gn:inta]
CONTIG512	26589591_f3_201	4472	10134	774	258	92	0.04499	Escherichia coli	b3412	[pn:bioh] [gn:bioh]
CONTIG512	6298291_f3_209	4473	10135	306	102	90	0.00059	Aedes aegypti	C40784	hypothetical protein 2 - aedes
CONTIG512	4453208_f3_218	4474	10136	2277	759	110	0.0028	Escherichia coli	b1617	[pn:beta-d-glucuronidase] [gn:uida]
CONTIG512	9772307_f3_224	4475	10137	11517	3839	681	9.8(10)-77	Acinetobacter calcoaceticus	AF011339	[PN:unknown] [DE:Acinetobacter calcoaceticus unknown protein gene, partial cds.] [LE:<1] [RE:2753] [DI:direct]
CONTIG512	22057062_f3_228	4476	10138	2226	742	451	2.1(10)-42	Bacillus subtilis	yknV	[pn:hypothetical protein]
CONTIG512	13864182_f3_231	4477	10139	447	149	408	3.5(10)-38	Escherichia coli	b0483	[pn:hypothetical protein]
CONTIG512	23516311_f3_238	4478	10140	681	227	431	1.3(10)-40	Synechocystis sp.	S77469	[PN:hypothetical protein] [OR:Synechocystis sp.] [SR:PCC 6803,, PCC 6803] [SR:PCC 6803,,]
CONTIG512	4723443_f3_266	4479	10141	528	176	448	2.0(10)-42	Escherichia coli	b2668	[pn:hypothetical protein] [gn:ygap]
CONTIG512	13726077_c1_269	4480	10142	240	80	170	5.7(10)-13	Escherichia coli	b2666	[pn:hypothetical protein]
CONTIG512	32207625_c1_270	4481	10143	1773	591	2214	1.5(10)-229	Escherichia coli	b0321	[pn:hypothetical protein] [gn:yahg]
CONTIG512	5986592_c1_278	4482	10144	1011	337	1145	2.7(10)-116	Rhizobium sp.	P55573	hypothetical transketolase family protein y4mn,
CONTIG512	32079650_c1_294	4483	10145	252	84	142	5.2(10)-10	Escherichia coli	U73857	or:escherichia coli le:5101 re:5604 di:direct nt:hypothetical protein
CONTIG512	4494082_c1_319	4484	10146	342	114	116	3.0(10)-7	Bacillus subtilis	spolIAA	[pn:anti-anti-sigma factor]
CONTIG512	12948567_c1_332	4485	10147	291	97	391	2.2(10)-36	Escherichia coli	I41306	hypothetical protein (argf-lacZ region) - escherichia coli

CONTIG512	10187_c1_337	4486	10148	411	137	133	4.7(10)-9	Escherichia coli	D90888	or:escherichia coli gn:yfjc le:13589 re:13900 di:complement sr:escherichia coli (strain:k12) dna, clone_lib:kohara lambda minise nt:similar to [swissprot accession number p37907]
CONTIG512	34662787_c2_339	4487	10149	624	208	703	1.8(10)-69	Escherichia coli	b0318	[pn:hypothetical protein] [gn:yahd]
CONTIG512	11187791_c2_340	4488	10150	1650	550	2082	1.3(10)-215	Escherichia coli	b0320	[pn:hypothetical protein] [gn:yahf]
CONTIG512	24785843_c2_347	4489	10151	990	330	1312	5.5(10)-134	Escherichia coli	b0323	[pn:hypothetical protein] [gn:yahi]
CONTIG512	31337753_c2_352	4490	10152	1335	445	444	5.2(10)-42	Escherichia coli	b3127	[pn:hypothetical 49.0 kd protein in mpb-soha intergenic region] [gn:yahau]
CONTIG512	30267840_c2_353	4491	10153	762	254	407	4.4(10)-38	Escherichia coli	b1093	[pn:3-oxoacyl-acyl-carrier protein reductase] [gn:fabg]
CONTIG512	14954818_c2_354	4492	10154	858	286	891	2.2(10)-89	Rhizobium sp.	P55574	hypothetical transketolase family protein y4mo,
CONTIG512	15712817_c2_359	4493	10155	1185	395	432	9.9(10)-41	Bacillus subtilis	ydfD	[pn:hypothetical protein]
CONTIG512	1377152_c2_415	4494	10156	435	145	91	0.00309	Bacillus subtilis	rsbW	[pn:switch protein/serine kinase and anti-sigma factor]
CONTIG512	14082025_c2_428	4495	10157	1200	400	93	0.20999	Methanococcus jannaschii	MJ0156	[pn:acetyl-coa decarboxylase/synthase, subunit beta]
CONTIG512	21678202_c2_429	4496	10158	1296	432	720	3.0(10)-71	Escherichia coli	b2020	[pn:histidinol dehydrogenase] [gn:hisd]
CONTIG512	31725412_c2_430	4497	10159	1365	455	284	3.3(10)-23	Methanobacterium thermoautotrophicum	MTH1496	[pn:amidase]
CONTIG512	787577_c2_435	4498	10160	477	159	701	3.1(10)-69	Escherichia coli	b2619	[pn:hypothetical protein] [gn:yfig]
CONTIG512	35625788_c3_438	4499	10161	381	127	231	2.0(10)-19	Escherichia coli	b2669	[pn:dna-binding protein spa] [gn:stpa]
CONTIG512	5989812_c3_448	4500	10162	1632	544	509	6.9(10)-49	Synechocystis sp.	S76103	[PN:hypothetical protein] [OR:Synechocystis sp.] [SR:PCC 6803., PCC 6803] [SR:PCC 6803.]

CONTIG512	31350680_c3_451	4501	10163	1458	486	2018	8.5(10)-209	Escherichia coli	b0324	[pn:hypothetical protein] [gn:yahj]
CONTIG512	26818956_c3_457	4502	10164	861	287	358	6.9(10)-33	Haemophilus influenzae	HI0912	[pn:gp]
CONTIG512	31307931_c3_460	4503	10165	903	301	397	5.0(10)-37	Escherichia coli	b3243	[pn:hypothetical protein] [gn:yhcs]
CONTIG512	24632632_c3_495	4504	10166	744	248	632	6.4(10)-62	Synechocystis sp.	S76993	[PN:hypothetical protein] [OR:Synechocystis sp.] [SR:PCC 6803, PCC 6803] [SR:PCC 6803, ]
CONTIG512	34275752_c3_515	4505	10167	1269	423	210	1.3(10)-16	Haemophilus influenzae	HI0884	[pn:aerobic respiration control protein arca] [gn:arca]
CONTIG512	29383255_c3_527	4506	10168	1209	403	340	5.5(10)-31	Escherichia coli	b2714	[pn:asc operon repressor protein] [gn:aseg]
CONTIG512	30347307_c3_535	4507	10169	984	328	437	2.8(10)-41	Escherichia coli	b3558	[pn:insertion element is150 hypothetical 33.3 kd protein] [gn:yif5b]
CONTIG512	32660955_c3_540	4508	10170	372	124	409	2.7(10)-38	Escherichia coli	b2618	[pn:hypothetical protein]
CONTIG513	23629567_fl_1	4509	10171	2427	809	1447	2.7(10)-148	Escherichia coli	b0124	[pn:glucose dehydrogenase] [gn:gcd]
CONTIG513	7282800_fl_18	4510	10172	378	126	509	6.9(10)-49	Escherichia coli	b1599	[pn:hypothetical protein]
CONTIG513	24406285_fl_26	4511	10173	918	306	1150	8.1(10)-117	Escherichia coli	b1595	[pn:hypothetical protein] [gn:yaf]
CONTIG513	4891293_fl_40	4512	10174	687	229	758	2.7(10)-75	Proteus mirabilis	P07641	chloramphenicol acetyltransferase (ec 2.3.1.28).
CONTIG513	25584832_fl_50	4513	10175	723	241	868	6.2(10)-87	Escherichia coli	b1585	[pn:hypothetical protein] [gn:yafc]
CONTIG513	33876063_fl_52	4514	10176	1116	372	1366	1.1(10)-139	Escherichia coli	b1580	[pn:starvation sensing protein rspb] [gn:rsbp]
CONTIG513	7119082_fl_59	4515	10177	2037	679	2392	2.0(10)-248	Escherichia coli	b1538	[pn:dipeptidyl carboxypeptidase ii] [gn:dep]
CONTIG513	23611652_fl_61	4516	10178	825	275	322	4.5(10)-29	Escherichia coli	b1790	[pn:hypothetical protein]
CONTIG513	24259677_fl_62	4517	10179	495	165	293	5.2(10)-26	Bacillus subtilis	ybbK	[pn:hypothetical protein]
CONTIG513	1048177_fl_70	4518	10180	309	103	138	1.3(10)-9	Escherichia coli	b3238	[pn:hypothetical protein] [gn:yhcn]



CONTIG513	30286561_fl_76	4519	10181	1590	530	2218	5.5(10)-230	Escherichia coli	b1385	[pn:hypothetical protein] [gn:feab]
CONTIG513	24490626_fl_85	4520	10182	999	333	1487	1.6(10)-152	Escherichia coli	b1388	[pn:hypothetical protein]
CONTIG513	14539213_fl_86	4521	10183	300	100	495	2.1(10)-47	Escherichia coli	b1389	[pn:hypothetical protein]
CONTIG513	4713467_fl_91	4522	10184	798	266	1016	1.3(10)-102	Escherichia coli	b1394	[pn:hypothetical protein]
CONTIG513	12276640_fl_96	4523	10185	1332	444	2185	1.7(10)-226	Escherichia coli	b1398	[pn:hypothetical protein]
CONTIG513	26737775_fl_97	4524	10186	963	321	1260	1.8(10)-128	Escherichia coli	b1399	[pn:hypothetical protein]
CONTIG513	4198568_fl_98	4525	10187	615	205	951	1.0(10)-95	Escherichia coli	b1400	[pn:hypothetical protein]
CONTIG513	23489792_fl_104	4526	10188	1548	516	654	3.0(10)-64	Mycobacterium tuberculosis	AL021932	[de:mycobacterium tuberculosis sequence v037.] [pn:putative dehydrogenase] [gn:mtv037.13c] [nt:mtv037.13c, len]
CONTIG513	14878332_fl_105	4527	10189	1227	409	495	2.1(10)-47	Escherichia coli	b1661	[pn:cyclopropane-fatty-acyl-phospholipid synthase] [gn:cfa]
CONTIG513	42312_fl_115	4528	10190	3942	1314	6025	0	Escherichia coli	b1413	[pn:atp-dependent helicase hrpa] [gn:hrpa]
CONTIG513	126541_fl_141	4529	10191	1062	354	154	1.3(10)-8	Bacillus subtilis	ywcH	[pn:hypothetical protein] [gn:ipa-44c]
CONTIG513	885766_fl_153	4530	10192	1290	430	1818	1.3(10)-187	Escherichia coli	b1594	[pn:protein] [gn:mlc]
CONTIG513	23462778_fl_154	4531	10193	699	233	1033	2.0(10)-104	Escherichia coli	b1593	[pn:hypothetical protein]
CONTIG513	21579652_fl_178	4532	10194	351	117	528	6.7(10)-51	Escherichia coli	b1582	[pn:hypothetical protein]
CONTIG513	4535002_fl_179	4533	10195	1254	418	2087	4.2(10)-216	Escherichia coli	b1581	[pn:starvation sensing protein rspa] [gn:rspa]
CONTIG513	24735213_fl_180	4534	10196	1437	479	1821	6.4(10)-188	Escherichia coli	b1543	[pn:hypothetical protein]
CONTIG513	4769191_fl_181	4535	10197	1476	492	2005	2.0(10)-207	Escherichia coli	b1542	[pn:hypothetical protein] [gn:ydfi]
CONTIG513	11754183_fl_195	4536	10198	2148	716	1055	9.5(10)-107	Escherichia coli	b0124	[pn:glucose dehydrogenase] [gn:gcd]

CONTIG513	4804068_f2_202	4537	10199	201	67	208	5.4(10)-17	Escherichia coli	b1382	[pn:hypothetical protein]
CONTIG513	13808563_f2_213	4538	10200	594	198	765	5.0(10)-76	Escherichia coli	b1391	[pn:hypothetical protein]
CONTIG513	26851665_f2_216	4539	10201	783	261	774	5.7(10)-77	Escherichia coli	b1393	[pn:hypothetical protein] [gn:ydb5]
CONTIG513	32692805_f2_217	4540	10202	1581	527	1940	1.6(10)-200	Escherichia coli	b1395	[pn:hypothetical protein] [gn:ydbu]
CONTIG513	12710012_f2_218	4541	10203	1290	430	1760	1.8(10)-181	Escherichia coli	b1397	[pn:hypothetical protein]
CONTIG513	4767043_f2_225	4542	10204	1986	662	1596	4.5(10)-164	Escherichia coli	b4083	[pn:hypothetical 73.7 kd protein in fdhf-phnp intergenic region]
CONTIG513	15676040_f2_227	4543	10205	948	316	203	1.8(10)-16	Bacillus subtilis	dltE	[pn:hypothetical protein] [gn:ipa-lr]
CONTIG513	32506457_f2_229	4544	10206	753	251	231	2.0(10)-19	Mycobacterium tuberculosis	AL021932	[de:mycobacterium tuberculosis sequence v037.] [pn:hypothetical protein mtv037.12c]
CONTIG513	26584505_f2_233	4545	10207	963	321	176	3.2(10)-13	Haemophilus influenzae	H11399	[pn:hypothetical protein]
CONTIG513	14879707_f2_238	4546	10208	546	182	632	6.4(10)-62	Escherichia coli	b0354	[pn:hypothetical protein]
CONTIG513	34194791_f2_255	4547	10209	846	282	868	6.2(10)-87	Escherichia coli	b1414	[pn:hypothetical protein in alda 5'' region] [gn:ydcf]
CONTIG513	22744037_f3_272	4548	10210	984	328	182	4.7(10)-12	Mycobacterium tuberculosis	Z97193	[PN:hypothetical protein MTCY180.06] [GN:MTCY180.06]
CONTIG513	38275_f3_277	4549	10211	432	144	477	1.7(10)-45	Escherichia coli	b1600	[DE:Mycobacterium tuberculosis cosmid Y180.] [NT:MTCY180.06, possible oxidoreductase, Len: 334.] [LE:6203] [RE:7207] [DI:direct]
CONTIG513	859452_f3_299	4550	10212	456	152	142	1.7(10)-9	Escherichia coli	b1328	[pn:hypothetical protein] [gn:ycjz]
CONTIG513	35581281_f3_320	4551	10213	459	153	383	1.5(10)-35	Escherichia coli	b0607	[pn:hypothetical protein] [gn:ybdq]
CONTIG513	5272338_f3_339	4552	10214	183	61	168	9.4(10)-13	Escherichia coli	b1550	[pn:hypothetical protein]

CONTIG513	24644576_f3_344	4553	10215	2646	882	3438	0	Escherichia coli	b1381	[pn:hypothetical protein] [gn:ydbh]
CONTIG513	16535137_f3_345	4554	10216	348	116	299	1.2(10)-26	Escherichia coli	b1383	[pn:hypothetical protein]
CONTIG513	16657590_f3_365	4555	10217	774	258	997	1.3(10)-100	Escherichia coli	b1390	[pn:hypothetical protein]
CONTIG513	16269758_f3_366	4556	10218	1200	400	1585	6.5(10)-163	Escherichia coli	b1392	[pn:hypothetical protein]
CONTIG513	34614416_f3_371	4557	10219	513	171	569	3.0(10)-55	Escherichia coli	b1396	[pn:hypothetical protein]
CONTIG513	22117188_f3_387	4558	10220	978	326	437	2.8(10)-41	Haemophilus influenzae	HI1364	[pn:hypothetical protein]
CONTIG513	22870175_f3_406	4559	10221	483	161	597	3.2(10)-58	Escherichia coli	b1415	[pn:lactaldehyde dehydrogenase a] [gn:alda]
CONTIG513	22069212_c1_408	4560	10222	1668	556	134	1.3(10)-5	Bacillus subtilis	yhcA	[pn:hypothetical protein]
CONTIG513	6539783_c1_410	4561	10223	1452	484	420	1.8(10)-39	Escherichia coli	b0572	[pn:hypothetical protein] [gn:yldb]
CONTIG513	20214775_c1_439	4562	10224	696	232	116	4.2(10)-5	Escherichia coli	b0375	[pn:hypothetical protein] [gn:yaiv]
CONTIG513	23870400_c1_454	4563	10225	2091	697	2967	0	Escherichia coli	b1387	[pn:hypothetical protein] [gn:maoc]
CONTIG513	13944091_c1_480	4564	10226	231	77	247	4.0(10)-21	Escherichia coli	b1541	[pn:hypothetical protein]
CONTIG513	12911066_c1_491	4565	10227	939	313	138	3.1(10)-9	Homo sapiens	Q09327	beta-1,4-mannosyl-glycoprotein beta-1,4-n-acetylglucosaminyl-transferase (ec 2.4.1.144) (n-glycosyl-oligosaccharide-glycoprotein n-acetylglucosaminyltransferase iii) (n-acetylglucosaminyltransferase iii) (gnt-iii) (glcnac-t iii)
CONTIG513	15786281_c1_492	4566	10228	348	116	323	3.5(10)-29	Escherichia coli	b1586	[pn:hypothetical protein]
CONTIG513	30361281_c1_495	4567	10229	270	90	221	3.5(10)-17	Escherichia coli	b1587	[pn:hypothetical protein]
CONTIG513	22925083_c1_502	4568	10230	783	261	97	0.006	Sinorhizobium meliloti	AF031940	[de:sinorhizobium meliloti alcohol dehydrogenase (adha) gene, completecds.] [pn:hypothetical protein] [nt:orf1]

CONTIG513	15679061_c1_504	4569	10231	672	224	291	8.6(10)-26	Bacillus subtilis	opuCB	[pn:glycine betaine/carnitine/choline abc transporter] [gn:yvbd]
CONTIG513	25961525_c1_512	4570	10232	1503	501	823	3.7(10)-82	Bacillus subtilis	ypwA	[pn:hypothetical protein]
CONTIG513	11722677_c1_521	4571	10233	1461	487	1881	2.7(10)-194	Escherichia coli	b1184	[pn:umuc protein] [gn:umuc]
CONTIG513	892531_c1_523	4572	10234	996	332	1546	8.9(10)-159	Klebsiella pneumoniae	P27504	coenzyme pqq synthesis protein b.
CONTIG513	914591_c2_541	4573	10235	1209	403	338	9.0(10)-31	Escherichia coli	b2685	[pn:multidrug resistance protein a] [gn:emra]
CONTIG513	641967_c2_549	4574	10236	858	286	422	1.1(10)-39	Escherichia coli	b0058	[pn:hypothetical 24.9 kd protein in sura-hepa intergenic region] [gn:yabo]
CONTIG513	1969212_c2_550	4575	10237	1041	347	321	5.5(10)-28	Escherichia coli	b3829	[pn:5-methyltetrahydropteroyltriglutamate-homocysteine methyltransferase] [gn:mete]
CONTIG513	32667541_c2_582	4576	10238	2502	834	3407	0	Escherichia coli	b1386	[pn:copper amine oxidase precursor] [gn:tyna]
CONTIG513	14063411_c2_583	4577	10239	2037	679	1604	6.2(10)-165	Escherichia coli	b4083	[pn:hypothetical 73.7 kd protein in fdhf-phnp intergenic region]
CONTIG513	10338967_c2_584	4578	10240	1158	386	268	1.3(10)-21	Escherichia coli	b1025	[pn:hypothetical protein] [gn:yedf]
CONTIG513	3324025_c2_586	4579	10241	438	146	173	2.7(10)-13	Azospirillum brasilense	X70360	or:azospirillum brasilense gn:carr le:59 re:580 di:direct nt:orf2
CONTIG513	30257702_c2_588	4580	10242	951	317	982	5.2(10)-99	Escherichia coli	b1384	[pn:hypothetical protein] [gn:fear]
CONTIG513	23908516_c2_595	4581	10243	1020	340	1498	1.1(10)-153	Escherichia coli	b1380	[pn:d-lactate dehydrogenase] [gn:ldha]
CONTIG513	915887_c2_596	4582	10244	840	280	1066	6.5(10)-108	Pseudomonas pyrocinia	JN0828	chloroperoxidase (ec 1.11.1.-) precursor - pseudomonaspyrocinia this enzyme catalyzes the formation of carbon-halogen bonds in the presence of hydrogen peroxide, halide ions, and a suitable organic substrate. this enzyme is involved in the production o
CONTIG513	9823576_c2_606	4583	10245	696	232	1021	3.7(10)-103	Escherichia coli	b1931	[pn:yedg] [gn:yedk]

CONTIG513	36070465_c2_609	4384	10246	852	284	1113	0.00017	Methanobacterium thermoautotrophicum	MTH882	[pn:conserved protein]
CONTIG513	3250786_c2_620	4385	10247	609	203	870	3.7(10)-87	Escherichia coli	b1584	[pn:spermidine n1-acetyltransferase] [gn:spg]
CONTIG513	25869028_c2_623	4386	10248	2325	775	3450	0	Escherichia coli	b1587	[pn:hypothetical protein]
CONTIG513	14572206_c2_624	4387	10249	621	207	1110	1.3(10)-112	Escherichia coli	b0895	[pn:anaerobic dimethyl sulfoxide reductase chain b] [gn:dmsb]
CONTIG513	13126632_c2_626	4388	10250	723	241	764	6.5(10)-76	Escherichia coli	b1591	[pn:hypothetical protein]
CONTIG513	24475938_c2_629	4389	10251	291	97	156	8.4(10)-11	Pseudomonas aeruginosa	Q01609	hypothetical 40.7 kd protein in opde 3region (orf2).
CONTIG513	14322166_c2_631	4390	10252	738	246	305	2.8(10)-27	Bacillus subtilis	opuCB	[pn:glycine betaine/carnitine/choline abc transporter] [gn:yvbd]
CONTIG513	9899186_c2_632	4391	10253	912	304	362	2.6(10)-33	Bacillus subtilis	opuCC	[pn:osmoprotectant-binding protein] [gn:yvbc]
CONTIG513	164182_c2_641	4392	10254	1338	446	1829	9.0(10)-189	Escherichia coli	b1596	[pn:hypothetical protein] [gn:yvfm]
CONTIG513	12597675_c2_643	4393	10255	843	281	1086	4.9(10)-110	Escherichia coli	b1598	[pn:hypothetical protein]
CONTIG513	16602291_c2_658	4394	10256	438	146	424	7.0(10)-40	Klebsiella pneumoniae	P27506	coenzyme pqq synthesis protein d.
CONTIG513	14744537_c2_660	4395	10257	2298	766	2069	3.3(10)-214	Klebsiella pneumoniae	P27508	coenzyme pqq synthesis protein f (ec 3.4.99.-).
CONTIG513	21729677_c3_684	4396	10258	666	222	928	2.7(10)-93	Escherichia coli	b1412	[pn:acyl carrier protein phosphodiesterase] [gn:acpd]
CONTIG513	31502177_c3_754	4397	10259	1299	433	434	8.9(10)-49	Escherichia coli	b2796	[pn:putative serine transporter] [gn:sdac]
CONTIG513	1206557_c3_755	4398	10260	246	82	263	8.0(10)-23	Salmonella typhimurium	Q56031	virulence protein msga.
CONTIG513	1257206_c3_763	4399	10261	897	299	1144	3.5(10)-116	Escherichia coli	b1539	[pn:hypothetical oxidoreductase in dcp-noha intergenic region]
CONTIG513	24813151_c3_764	4600	10262	708	236	998	1.0(10)-100	Escherichia coli	b1540	[pn:hypothetical protein]
CONTIG513	2189155_c3_775	4601	10263	372	124	408	3.5(10)-38	Escherichia coli	b1583	[pn:hypothetical protein]

CONTIG513	14473425_c3_783	4602	10264	927	309	932	1.0(10)-93	Escherichia coli	b1590	[pn:hypothetical protein]
CONTIG513	10945780_c3_785	4603	10265	453	151	236	5.7(10)-20	Escherichia coli	b1848	[pn:hypothetical 10.7 kd protein in purt 5' region] [gn:yebg]
CONTIG513	6025635_c3_790	4604	10266	1257	419	806	2.2(10)-80	Bacillus subtilis	opuBA	[pn:choline abc transporter] [gn:prov]
CONTIG513	13159433_c3_791	4605	10267	1416	472	1077	4.4(10)-109	Escherichia coli	b1592	[pn:hypothetical protein]
CONTIG513	29882802_c3_803	4606	10268	1065	355	1065	8.3(10)-108	Escherichia coli	b1601	[pn:hypothetical protein]
CONTIG513	10662961_c3_806	4607	10269	363	121	561	2.1(10)-54	Escherichia coli	b0410	[pn:yajd]
CONTIG513	36382212_c3_807	4608	10270	1038	346	1150	8.1(10)-117	Klebsiella pneumoniae	P27509	hypothetical protein in pqqa 5' region (orf x) (fragment).
CONTIG513	16069816_c3_808	4609	10271	843	281	1153	3.8(10)-117	Klebsiella pneumoniae	P27505	coenzyme pqq synthesis protein c.
CONTIG513	24066262_c3_810	4610	10272	1185	395	1943	7.5(10)-201	Klebsiella pneumoniae	P27507	coenzyme pqq synthesis protein e.
CONTIG52	33807686_fl_1	4611	10273	465	155	723	1.3(10)-71	Escherichia coli	b0683	[pn:ferric uptake regulation protein] [gn:fur]
CONTIG52	4167842_c3_6	4612	10274	240	80	250	1.8(10)-21	Escherichia coli	b0682	[pn:hypothetical protein] [gn:ybfm]
CONTIG53	10359689_c2_3	4613	10275	425	141	660	6.9(10)-65	Escherichia coli	b1827	[pn:hypothetical protein]
CONTIG54	14978382_fl_1	4614	10276	356	119	192	9.4(10)-15	Thiobacillus ferrooxidans	P20085	mobilization protein mobil.
CONTIG56	4070952_f2_2	4615	10277	303	101	92	0.00021	Escherichia coli	b0530	[pn:hypothetical protein] [gn:sfma]
CONTIG58	10947127_fl_1	4616	10278	459	153	543	1.7(10)-52	Escherichia coli	b1938	[pn:flagellar basal-body m-ring protein] [gn:flif]
CONTIG58	23641281_c2_7	4617	10279	477	159	186	1.2(10)-14	Bacillus subtilis	ydfF	[pn:hypothetical protein]
CONTIG59	36350016_c2_5	4618	10280	744	248	1104	6.0(10)-112	Escherichia coli	b3232	[pn:hypothetical 43.1 kd protein in rplm-hhoA intergenic region] [gn:yhem]
CONTIG6	29877090_c2_8	4619	10281	339	113	92	0.00067	Bacillus subtilis	ccpA	[pn:transcriptional regulator] [gn:grar]
CONTIG6	32494527_c3_9	4620	10282	426	142	217	6.0(10)-18	Haemophilus influenzae	HI0522	[pn:sp]

CONTIG60	29322524_c1_3	4621	10283	385	128	325	2.2(10)-29	Bacteriophage phi-80	P14819	repressor protein ci.
CONTIG61	35550041_c2_6	4622	10284	1290	430	1593	9.3(10)-164	Escherichia coli	b0955	[pn:hypothetical protein]
CONTIG62	33261442_f1_1	4623	10285	1028	343	101	0.034	Thiobacillus ferrooxidans	AF032884	[de:thiobacillus ferrooxidans n-acetylglucosamine-1-phosphateuridylyltransferase (glmU) gene, partial cds; glucosamine synthase(glmS) and regg (regg) genes, complete cds; and transposon tn5468,complete sequence"] [pn:tnsd]
CONTIG63	13071907_f2_3	4624	10286	246	82	339	7.0(10)-31	Escherichia coli	b0957	[pn:outer membrane protein a] [gn:ompa]
CONTIG63	22276682_c1_6	4625	10287	483	161	709	4.4(10)-70	Escherichia coli	b0956	[pn:hypothetical protein in ompa 3"] [gn:ycbg]
CONTIG64	22847502_c1_6	4626	10288	198	66	100	8.0(10)-5	Escherichia coli	b1987	[pn:transcriptional regulator] [gn:cbl]
CONTIG64	26453393_c3_7	4627	10289	822	274	1178	8.8(10)-120	Escherichia coli	b1987	[pn:transcriptional regulator] [gn:cbl]
CONTIG65	25838388_c1_1	4628	10290	219	73	269	1.8(10)-23	Escherichia coli	b0958	[pn:cell division inhibitor] [gn:sula]
CONTIG66	4866552_f2_2	4629	10291	426	142	120	1.1(10)-7	Escherichia coli	L43373	or:escherichia coli pn:pilin le:22 re:567 di:direct sr:escherichia coli (strain 31a/o6) dna
CONTIG67	13175643_f1_1	4630	10292	1026	342	135	1.7(10)-8	Haemophilus influenzae	H11488	[pn:e16 protein] [gn:muel6]
CONTIG7	31267540_f2_2	4631	10293	282	94	326	1.7(10)-29	Escherichia coli	b1734	[pn:phospho-beta-glucosidase b] [gn:celf]
CONTIG7	11729152_f3_3	4632	10294	345	115	138	1.1(10)-8	Escherichia coli	b1734	[pn:phospho-beta-glucosidase b] [gn:celf]
CONTIG70	22066625_f3_2	4633	10295	687	229	961	8.6(10)-97	Escherichia coli	b0055	[pn:hypothetical 30.6 kd protein in sura-hepa intergenic region] [gn:yabh]
CONTIG71	23880458_f2_2	4634	10296	579	193	962	6.7(10)-97	Escherichia coli	b0675	[pn:nagd protein] [gn:nagd]
CONTIG71	16251037_f3_4	4635	10297	345	115	370	5.7(10)-34	Escherichia coli	b0674	[pn:asparagine synthetase b] [gn:asnb]

CONTIG72	3943817_c1_3	4636	10298	684	228	104	0.00289	Yersinia enterocolitica (type 0:8)	U46859	or:yersinia enterocolitica (type 0:8) pn:rfbx gn:rfbx le:6526 re:7815 di:direct
CONTIG73	33719031_c3_4	4637	10299	594	198	1003	3.1(10)-101	Escherichia coli	b0223	[pn:hypothetical protein in gmha-thia intergenic region] [gn:yafj]
CONTIG74	431332_f3_1	4638	10300	987	329	94	0.07	Bacillus subtilis	yvrC	[pn:hypothetical protein]
CONTIG77	32145043_c2_7	4639	10301	222	74	235	7.5(10)-20	synthetic construct	M15619	or:artificial sequence le:29 re:>232 di:direct sr:e.coli (strain se5000) synthetic dna, clone pkl nt:orf16-lacZ fusion protein
CONTIG78	31901556_c2_6	4640	10302	1035	345	1101	1.3(10)-111	Escherichia coli	X05874	or:escherichia coli pn:mature receptor protein le:244 re:2367 di:direct nt:author-given protein sequence is in conflict with
CONTIG80	23629716_f3_2	4641	10303	614	205	909	2.7(10)-91	Escherichia coli	b0054	[pn:organic solvent tolerance protein precursor] [gn:imp]
CONTIG82	14570453_c3_4	4642	10304	366	122	159	8.4(10)-12	Escherichia coli	b1160	[pn:hypothetical protein]
CONTIG83	22166288_f3_3	4643	10305	801	267	1289	1.5(10)-131	Escherichia coli	b0432	[pn:cytochrome o ubiquinol oxidase subunit ii] [gn:cyoa]
CONTIG84	35663506_c1_7	4644	10306	441	147	419	2.3(10)-39	Escherichia coli	b3344	[pn:hypothetical protein] [gn:yhem]
CONTIG84	12203427_c2_8	4645	10307	441	147	396	6.5(10)-37	Escherichia coli	b3345	[pn:hypothetical 13.6 kd protein in rpsl-fkpa intergenic region]
CONTIG85	26573386_f3_3	4646	10308	615	205	328	1.0(10)-29	Escherichia coli	b1025	[pn:hypothetical protein] [gn:ycdt]
CONTIG85	2080192_c3_7	4647	10309	390	130	545	1.1(10)-52	Escherichia coli	b1642	[pn:hypothetical protein] [gn:slya]
CONTIG86	36150166_c1_2	4648	10310	468	156	100	0.00064	Haemophilus influenzae	H10676	[pn:integrase-recombinase protein] [gn:xere]
CONTIG87	12308130_c1_4	4649	10311	402	134	334	2.3(10)-30	Escherichia coli	b1090	[pn:plsx protein] [gn:plsx]
CONTIG87	20738833_c1_5	4650	10312	519	173	762	1.1(10)-75	Escherichia coli	b1091	[pn:3-oxoacyl-acyl-carrier-protein synthase iii] [gn:fabh]
CONTIG88	6362961_c1_4	4651	10313	789	263	1231	2.1(10)-125	Escherichia coli	b0931	[pn:nicotinate phosphoribosyltransferase] [gn:pncb]
CONTIG89	22078757_c2_4	4652	10314	522	174	259	2.1(10)-22	Escherichia coli	b1374	[pn:hypothetical protein]



CONTIG90	2150290_f1_1	4653	10315	651	217	781	1.0(10)-77	Escherichia coli	b0212	[pn:hypothetical protein] [gn:glob]
CONTIG90	4538138_c3_6	4654	10316	420	140	511	4.2(10)-49	Escherichia coli	b0213	[pn:hypothetical protein] [gn:yafs]
CONTIG91	135805_c2_5	4655	10317	849	283	853	1.6(10)-104	Escherichia coli	b2581	[pn:hypothetical 37.8 kd protein in ung 3''' region] [gn:yfif]
CONTIG92	117202_f1_1	4656	10318	429	143	734	9.9(10)-73	Escherichia coli	b3231	[pn:50s ribosomal subunit protein 113] [gn:rplm]
CONTIG92	25525308_f1_2	4657	10319	396	132	645	2.7(10)-63	Escherichia coli	b3230	[pn:30s ribosomal subunit protein s9] [gn:rpsi]
CONTIG93	32478807_f1_1	4658	10320	609	203	849	6.4(10)-85	Escherichia coli	b1818	[pn:pts system, mannose-specific iic component] [gn:many]
CONTIG93	26056711_f1_2	4659	10321	474	158	748	3.2(10)-74	Escherichia coli	b1819	[pn:pts system, mannose-specific iid component] [gn:manz]
CONTIG94	3412917_c1_4	4660	10322	414	138	313	4.0(10)-28	Escherichia coli	b1678	[pn:hypothetical protein]
CONTIG96	128251_c2_4	4661	10323	648	216	958	1.8(10)-96	Escherichia coli	b0222	[pn:phosphoheptose isomerase] [gn:gmha]
CONTIG98	10161416_c1_4	4662	10324	870	290	152	9.5(10)-9	Escherichia coli	b1959	[pn:hypothetical 32.2 kd protein in vsr 5''' region] [gn:yeda]
CONTIG99	13867212_f1_1	4663	10325	270	90	326	1.7(10)-29	Haemophilus influenzae	H11713	[pn:phosphohistidinoprotein-hexose phosphotransferase] [gn:ptsh]
CONTIG99	3964643_f1_2	4664	10326	845	282	1035	1.3(10)-104	Escherichia coli	b2416	[pn:phosphoenolpyruvate-protein phosphotransferase] [gn:ptsj]
CONTIG99	12298262_c1_4	4665	10327	357	119	269	1.8(10)-23	Salmonella typhimurium	S04160	hypothetical protein k - salmonella typhimurium (fragment)
CONTIG103	26692751_f3_2	4666	10328	192	64					
CONTIG104	4086013_f1_1	4667	10329	231	77					
CONTIG107	2461718_f3_2	4668	10330	426	142					
CONTIG107	160902_c2_4	4669	10331	198	66					
CONTIG108	16922530_f2_1	4670	10332	369	123					
CONTIG109	10550041_f3_3	4671	10333	258	86					
CONTIG112	24298378_c2_2	4672	10334	411	137					
CONTIG112	267327_c3_3	4673	10335	483	161					
CONTIG114	24328157_c2_2	4674	10336	924	308					
CONTIG116	32086557_c3_10	4675	10337	348	116					
CONTIG118	24508568_c3_17	4676	10338	183	61					
CONTIG12	24883437_c2_3	4677	10339	240	80					
CONTIG123	15647957_c2_8	4678	10340	189	63					

[illegible]

CONTIG172	5204702_f1_1	4718	10380	921	307						
CONTIG172	26359452_f1_2	4719	10381	531	177						
CONTIG172	5893755_f2_3	4720	10382	669	223						
CONTIG172	195825_f2_4	4721	10383	498	166						
CONTIG172	19535931_f2_5	4722	10384	270	90						
CONTIG172	2117802_f3_6	4723	10385	195	65						
CONTIG177	4870127_f1_1	4724	10386	234	78						
CONTIG178	25392128_f1_2	4725	10387	186	62						
CONTIG178	1070192_f3_5	4726	10388	231	77						
CONTIG178	1318937_c1_6	4727	10389	507	169						
CONTIG178	2925042_c1_8	4728	10390	204	68						
CONTIG178	24422680_c3_11	4729	10391	213	71						
CONTIG179	35391652_f1_2	4730	10392	717	239						
CONTIG179	33752182_f2_3	4731	10393	585	195						
CONTIG179	6250026_f3_5	4732	10394	690	230						
CONTIG181	24650693_f3_2	4733	10395	510	170						
CONTIG181	4739702_f3_3	4734	10396	258	86						
CONTIG181	14539091_c2_5	4735	10397	183	61						
CONTIG183	12207031_f1_1	4736	10398	183	61						
CONTIG183	36134701_f1_2	4737	10399	711	237						
CONTIG183	36134701_f1_3	4738	10400	711	237						
CONTIG183	16595443_f1_4	4739	10401	204	68						
CONTIG185	969827_c3_5	4740	10402	198	66						
CONTIG189	22083507_c1_8	4741	10403	186	62						
CONTIG191	9980208_f3_5	4742	10404	303	101						
CONTIG194	4494002_f1_1	4743	10405	222	74						
CONTIG194	16616077_f3_3	4744	10406	516	172						
CONTIG194	4485092_f3_4	4745	10407	501	167						
CONTIG194	9767885_c3_12	4746	10408	186	62						
CONTIG196	23541552_f2_3	4747	10409	201	67						
CONTIG198	12986016_c3_9	4748	10410	189	63						
CONTIG201	22902166_f1_2	4749	10411	234	78						
CONTIG204	24408531_f2_3	4750	10412	231	77						
CONTIG205	564203_c2_6	4751	10413	1110	370						
CONTIG207	5261287_f3_7	4752	10414	276	92						
CONTIG208	33886591_f2_3	4753	10415	459	153						
CONTIG208	22273462_f3_4	4754	10416	369	123						
CONTIG208	25600262_f3_7	4755	10417	555	185						
CONTIG208	22833262_c3_16	4756	10418	237	79						



[illegible]

[illegible]

[illegible]

CONTIG318	6814128_f3_13	4913	10575	213	71						
CONTIG318	26363308_f3_18	4914	10576	309	103						
CONTIG318	30579507_c2_25	4915	10577	195	65						
CONTIG319	14957657_f1_2	4916	10578	186	62						
CONTIG319	10440713_f1_6	4917	10579	381	127						
CONTIG319	30339027_f1_7	4918	10580	555	185						
CONTIG319	19776557_f3_26	4919	10581	225	75						
CONTIG319	45245692_c2_30	4920	10582	333	111						
CONTIG319	22459705_c2_32	4921	10583	186	62						
CONTIG32	1979167_c1_2	4922	10584	186	62						
CONTIG320	4348250_f3_14	4923	10585	198	66						
CONTIG320	36378428_c1_22	4924	10586	228	76						
CONTIG322	20941652_c3_44	4925	10587	207	69						
CONTIG324	36022916_f1_1	4926	10588	249	83						
CONTIG326	16225827_f2_12	4927	10589	192	64						
CONTIG326	4094687_f2_13	4928	10590	273	91						
CONTIG326	6839662_f3_23	4929	10591	900	300						
CONTIG326	2346907_f3_29	4930	10592	198	66						
CONTIG326	21579680_c1_41	4931	10593	219	73						
CONTIG326	22345463_c1_43	4932	10594	417	139						
CONTIG326	26847916_c2_54	4933	10595	288	96						
CONTIG326	33790916_c3_56	4934	10596	342	114						
CONTIG327	34666516_f1_8	4935	10597	732	244						
CONTIG327	5117968_f1_9	4936	10598	597	199						
CONTIG327	32313591_f3_18	4937	10599	378	126						
CONTIG327	6119561_c2_37	4938	10600	219	73						
CONTIG329	31900713_c3_35	4939	10601	255	85						
CONTIG330	16015787_f1_7	4940	10602	432	144						
CONTIG330	786402_f3_17	4941	10603	267	89						
CONTIG330	22477305_c1_29	4942	10604	228	76						
CONTIG330	9975431_c1_31	4943	10605	525	175						
CONTIG330	11891652_c1_32	4944	10606	1206	402						
CONTIG330	5976381_c1_33	4945	10607	282	94						
CONTIG330	20006305_c1_36	4946	10608	441	147						
CONTIG330	26753427_c1_37	4947	10609	714	238						
CONTIG330	14844816_c2_38	4948	10610	1635	545						
CONTIG330	6719787_c2_42	4949	10611	291	97						
CONTIG330	26292592_c2_44	4950	10612	333	111						
CONTIG330	24663201_c3_47	4951	10613	801	267						



CONTIG330	25554761_c3_48	4952	10614	453	151
CONTIG330	23698253_c3_49	4953	10615	648	216
CONTIG330	36500637_c3_53	4954	10616	390	130
CONTIG331	1056337_f1_4	4955	10617	240	80
CONTIG331	14097265_f3_27	4956	10618	222	74
CONTIG331	3321012_c1_29	4957	10619	288	96
CONTIG331	21677281_c2_37	4958	10620	297	99
CONTIG331	23476027_c2_39	4959	10621	240	80
CONTIG331	16835967_c3_41	4960	10622	1434	478
CONTIG334	25976457_f3_22	4961	10623	372	124
CONTIG335	26836387_f2_13	4962	10624	297	99
CONTIG335	25975292_f2_20	4963	10625	213	71
CONTIG335	4738406_f3_25	4964	10626	234	78
CONTIG335	25604511_c1_35	4965	10627	192	64
CONTIG335	15730042_c1_36	4966	10628	495	165
CONTIG335	34381300_c2_40	4967	10629	312	104
CONTIG335	4816918_c2_43	4968	10630	1068	356
CONTIG335	21754515_c3_45	4969	10631	198	66
CONTIG336	16071968_c3_43	4970	10632	282	94
CONTIG338	24098442_f3_19	4971	10633	207	69
CONTIG338	9899066_c1_25	4972	10634	243	81
CONTIG338	24026930_c1_27	4973	10635	213	71
CONTIG339	4718808_c2_42	4974	10636	225	75
CONTIG340	26595288_f3_25	4975	10637	204	68
CONTIG340	1064000_c1_30	4976	10638	339	113
CONTIG340	30339591_c2_35	4977	10639	654	218
CONTIG340	21742077_c2_37	4978	10640	324	108
CONTIG340	24410292_c3_41	4979	10641	534	178
CONTIG341	11882767_f2_16	4980	10642	309	103
CONTIG342	23563886_f1_5	4981	10643	231	77
CONTIG342	34072137_f3_12	4982	10644	489	163
CONTIG342	4710801_f3_17	4983	10645	219	73
CONTIG343	32521877_c1_42	4984	10646	234	78
CONTIG344	35722591_c1_40	4985	10647	273	91
CONTIG344	21579650_c3_65	4986	10648	249	83
CONTIG346	32422687_f3_27	4987	10649	342	114
CONTIG346	23439037_c2_38	4988	10650	477	159
CONTIG347	34180260_f3_25	4989	10651	461	154
CONTIG348	9765713_c2_32	4990	10652	243	81

[illegible]

[illegible]

[illegible]

CONTIG392	36523503_c3_87	5108	10770	231	77						
CONTIG392	35346907_c3_93	5109	10771	183	61						
CONTIG392	6818751_c3_97	5110	10772	189	63						
CONTIG393	4078393_c2_59	5111	10773	282	94						
CONTIG394	20337881_f3_48	5112	10774	198	66						
CONTIG394	26740637_c2_76	5113	10775	525	175						
CONTIG394	23443808_c3_90	5114	10776	441	147						
CONTIG395	32714562_f1_20	5115	10777	192	64						
CONTIG395	978193_f3_36	5116	10778	210	70						
CONTIG397	21564443_f3_33	5117	10779	522	174						
CONTIG397	22839802_c3_73	5118	10780	183	61						
CONTIG398	674191_f1_1	5119	10781	843	281						
CONTIG398	34238555_f2_10	5120	10782	282	94						
CONTIG398	24646910_f2_11	5121	10783	462	154						
CONTIG398	35234806_c1_43	5122	10784	339	113						
CONTIG398	12581253_c1_55	5123	10785	219	73						
CONTIG398	24625127_c2_60	5124	10786	198	66						
CONTIG398	29845262_c2_72	5125	10787	282	94						
CONTIG398	6300465_c3_87	5126	10788	186	62						
CONTIG399	9850631_f3_47	5127	10789	210	70						
CONTIG399	4330033_f3_60	5128	10790	399	133						
CONTIG4	23626662_c3_2	5129	10791	315	105						
CONTIG40	24787507_c2_2	5130	10792	687	229						
CONTIG400	25428882_f2_18	5131	10793	186	62						
CONTIG400	4882068_f2_19	5132	10794	306	102						
CONTIG401	24349015_c2_78	5133	10795	186	62						
CONTIG402	30098465_c3_78	5134	10796	255	85						
CONTIG403	976062_f1_6	5135	10797	339	113						
CONTIG403	26578201_f1_8	5136	10798	324	108						
CONTIG403	31256953_f2_22	5137	10799	282	94						
CONTIG403	22384683_f2_27	5138	10800	570	190						
CONTIG403	24063540_f2_31	5139	10801	306	102						
CONTIG403	15098426_f3_45	5140	10802	297	99						
CONTIG403	24644503_c1_57	5141	10803	417	139						
CONTIG403	13162887_c1_58	5142	10804	582	194						
CONTIG403	16145253_c1_61	5143	10805	213	71						
CONTIG403	13083385_c1_66	5144	10806	1068	356						
CONTIG403	2189012_c1_70	5145	10807	186	62						
CONTIG403	24644661_c2_75	5146	10808	375	125						

CONTIG403	36505133_c2_76	5147	10809	183	61
CONTIG403	969555_c2_83	5148	10810	237	79
CONTIG403	26619755_c3_88	5149	10811	837	279
CONTIG403	22070937_c3_90	5150	10812	219	73
CONTIG404	15132081_f1_25	5151	10813	324	108
CONTIG404	4088917_f3_55	5152	10814	486	162
CONTIG404	788562_c2_94	5153	10815	246	82
CONTIG404	24228382_c3_105	5154	10816	480	160
CONTIG405	16052187_c1_91	5155	10817	231	77
CONTIG407	4487555_f1_7	5156	10818	438	146
CONTIG407	15098163_c2_77	5157	10819	303	101
CONTIG408	36501525_c2_78	5158	10820	231	77
CONTIG409	562758_f1_5	5159	10821	201	67
CONTIG409	33244092_f1_10	5160	10822	615	205
CONTIG409	16257132_f2_33	5161	10823	195	65
CONTIG409	23860287_c1_73	5162	10824	570	190
CONTIG409	978382_c1_74	5163	10825	357	119
CONTIG409	4150463_c2_81	5164	10826	402	134
CONTIG409	26267713_c2_82	5165	10827	444	148
CONTIG409	30180187_c3_90	5166	10828	1122	374
CONTIG410	16535932_f1_1	5167	10829	192	64
CONTIG410	26756876_c2_87	5168	10830	315	105
CONTIG412	9781875_c1_56	5169	10831	468	156
CONTIG412	14067533_c1_59	5170	10832	1089	363
CONTIG415	36519025_f1_6	5171	10833	249	83
CONTIG415	1456966_f1_25	5172	10834	269	90
CONTIG415	7164127_f2_40	5173	10835	186	62
CONTIG415	14333277_f2_41	5174	10836	1011	337
CONTIG415	12375451_f3_45	5175	10837	360	120
CONTIG415	4016588_c1_62	5176	10838	1329	443
CONTIG415	24430387_c1_63	5177	10839	216	72
CONTIG415	16206933_c3_91	5178	10840	249	83
CONTIG415	11213890_c3_93	5179	10841	183	61
CONTIG417	14585875_f2_26	5180	10842	369	123
CONTIG417	4939068_f2_27	5181	10843	330	110
CONTIG417	16449033_f2_28	5182	10844	228	76
CONTIG417	25472775_f2_33	5183	10845	225	75
CONTIG417	2473777_f2_40	5184	10846	201	67
CONTIG417	4689377_f3_49	5185	10847	624	208

CONTIG417	13089755_f3_50	5186	10848	297	99
CONTIG417	13064701_c1_66	5187	10849	924	308
CONTIG417	5113818_c1_67	5188	10850	237	79
CONTIG417	36152043_c1_70	5189	10851	306	102
CONTIG417	12710313_c1_85	5190	10852	279	93
CONTIG417	29797152_c1_86	5191	10853	195	65
CONTIG417	22766715_c2_91	5192	10854	369	123
CONTIG417	1172783_c2_93	5193	10855	354	118
CONTIG417	596062_c2_94	5194	10856	189	63
CONTIG417	20176562_c2_95	5195	10857	192	64
CONTIG417	2830015_c3_106	5196	10858	522	174
CONTIG417	2425063_c3_107	5197	10859	231	77
CONTIG417	26751500_c3_108	5198	10860	243	81
CONTIG417	23944181_c3_109	5199	10861	237	79
CONTIG417	11149130_c3_110	5200	10862	327	109
CONTIG417	4095068_c3_111	5201	10863	291	97
CONTIG417	36573586_c3_112	5202	10864	618	206
CONTIG417	2926025_c3_120	5203	10865	351	117
CONTIG417	31492187_c3_126	5204	10866	537	179
CONTIG419	25527333_f1_2	5205	10867	204	68
CONTIG419	31830442_f3_51	5206	10868	276	92
CONTIG419	17087599_c1_75	5207	10869	298	99
CONTIG419	34115881_c1_102	5208	10870	921	307
CONTIG419	25958441_c3_156	5209	10871	258	86
CONTIG420	17033191_f2_23	5210	10872	186	62
CONTIG422	35671890_f2_36	5211	10873	192	64
CONTIG422	1379750_c2_106	5212	10874	183	61
CONTIG423	24036533_f1_10	5213	10875	252	84
CONTIG423	24253275_f3_61	5214	10876	210	70
CONTIG423	4331568_c1_68	5215	10877	414	138
CONTIG423	35706317_c1_72	5216	10878	405	135
CONTIG423	4022201_c2_81	5217	10879	375	125
CONTIG423	32039193_c2_99	5218	10880	381	127
CONTIG423	32147708_c2_101	5219	10881	1419	473
CONTIG423	7132062_c3_109	5220	10882	498	166
CONTIG423	24506900_c3_110	5221	10883	378	126
CONTIG424	31735031_f3_78	5222	10884	474	158
CONTIG425	22145311_f1_6	5223	10885	243	81
CONTIG425	35292176_f1_16	5224	10886	234	78

[illegible]



[illegible]

CONTIG446	35323312_c3_101	5303	10965	339	113				
CONTIG448	32300026_f1_8	5304	10966	669	223				
CONTIG448	33883541_f2_35	5305	10967	345	115				
CONTIG448	7119500_f2_54	5306	10968	924	308				
CONTIG448	23956880_f3_65	5307	10969	885	295				
CONTIG448	21519626_c1_123	5308	10970	189	63				
CONTIG448	3165933_c2_130	5309	10971	753	251				
CONTIG448	2744655_c2_134	5310	10972	363	121				
CONTIG448	15641942_c2_154	5311	10973	207	69				
CONTIG449	25583402_f2_67	5312	10974	192	64				
CONTIG449	11848555_c2_161	5313	10975	243	81				
CONTIG450	984788_f2_32	5314	10976	222	74				
CONTIG450	24081268_f2_60	5315	10977	354	118				
CONTIG450	14160691_f3_66	5316	10978	198	66				
CONTIG450	522142_c3_176	5317	10979	204	68				
CONTIG451	33208290_f1_13	5318	10980	1626	542				
CONTIG451	26223761_f3_96	5319	10981	429	143				
CONTIG451	32057312_c1_110	5320	10982	333	111				
CONTIG451	20156577_c3_166	5321	10983	243	81				
CONTIG452	12535418_f1_4	5322	10984	204	68				
CONTIG452	29329665_f1_33	5323	10985	312	104				
CONTIG452	251877_f2_48	5324	10986	336	112				
CONTIG452	13682961_c3_162	5325	10987	534	178				
CONTIG453	1353518_f1_26	5326	10988	207	69				
CONTIG453	12548216_f2_46	5327	10989	252	84				
CONTIG453	26456263_f2_48	5328	10990	849	283				
CONTIG453	564203_c1_83	5329	10991	1740	580				
CONTIG453	26745133_c1_92	5330	10992	225	75				
CONTIG453	5100032_c1_100	5331	10993	246	82				
CONTIG453	29535313_c1_103	5332	10994	519	173				
CONTIG453	30484791_c2_121	5333	10995	204	68				
CONTIG453	23837802_c2_123	5334	10996	210	70				
CONTIG453	23444717_c3_133	5335	10997	492	164				
CONTIG453	10726592_c3_148	5336	10998	411	137				
CONTIG454	13790832_f1_22	5337	10999	195	65				
CONTIG454	34188765_f2_44	5338	11000	216	72				
CONTIG455	12206653_f3_77	5339	11001	234	78				
CONTIG455	26375632_c1_128	5340	11002	234	78				
CONTIG455	6429627_c2_154	5341	11003	285	95				

CONTIG456	24719182_f1_10	5342	11004	216	72
CONTIG456	21878392_f1_15	5343	11005	261	87
CONTIG456	14565641_f3_80	5344	11006	243	81
CONTIG456	24782768_c1_130	5345	11007	183	61
CONTIG456	3125905_c3_204	5346	11008	234	78
CONTIG457	16619340_f2_67	5347	11009	210	70
CONTIG457	30578461_f3_95	5348	11010	243	81
CONTIG457	16839066_c2_150	5349	11011	207	69
CONTIG458	21484377_f1_8	5350	11012	264	88
CONTIG458	11117202_c2_132	5351	11013	222	74
CONTIG459	11179693_f2_21	5352	11014	465	155
CONTIG462	284787_f1_14	5353	11015	795	265
CONTIG462	24431415_f1_22	5354	11016	186	62
CONTIG462	22439131_c1_90	5355	11017	339	113
CONTIG462	14745665_c1_91	5356	11018	351	117
CONTIG462	10746018_c2_108	5357	11019	222	74
CONTIG462	2167311_c2_121	5358	11020	447	149
CONTIG462	10815718_c3_142	5359	11021	393	131
CONTIG462	20916702_c3_143	5360	11022	672	224
CONTIG462	32147708_c3_144	5361	11023	1347	449
CONTIG463	22541312_f3_61	5362	11024	207	69
CONTIG463	7213408_f3_62	5363	11025	234	78
CONTIG463	16207251_c1_91	5364	11026	324	108
CONTIG464	16533162_f1_2	5365	11027	1161	387
CONTIG464	23695751_f2_23	5366	11028	1665	555
CONTIG464	32226650_c1_70	5367	11029	228	76
CONTIG464	2131291_c2_85	5368	11030	741	247
CONTIG465	270807_f1_50	5369	11031	261	87
CONTIG465	6677291_c2_165	5370	11032	207	69
CONTIG467	9801927_f2_58	5371	11033	972	324
CONTIG468	17011062_f3_103	5372	11034	510	170
CONTIG468	1442175_f3_105	5373	11035	276	92
CONTIG468	10197555_c1_137	5374	11036	336	112
CONTIG468	23960941_c2_175	5375	11037	768	256
CONTIG468	16285050_c3_177	5376	11038	351	117
CONTIG469	900841_f1_2	5377	11039	195	65
CONTIG469	32691941_c1_105	5378	11040	213	71
CONTIG469	2132827_c3_173	5379	11041	189	63
CONTIG47	21564003_f2_3	5380	11042	207	69

CONTIG470	4429212_f1_11	5381	11043	534	178
CONTIG470	21522283_c1_106	5382	11044	255	85
CONTIG470	26252193_c2_153	5383	11045	207	69
CONTIG470	32117201_c3_204	5384	11046	225	75
CONTIG471	32213278_f1_1	5385	11047	195	65
CONTIG471	10423968_f1_2	5386	11048	198	66
CONTIG471	2129526_f2_27	5387	11049	192	64
CONTIG471	12313528_c2_113	5388	11050	363	121
CONTIG471	11728926_c2_116	5389	11051	225	75
CONTIG472	36117968_f1_1	5390	11052	219	73
CONTIG472	25525333_f1_2	5391	11053	426	142
CONTIG472	32301461_c1_109	5392	11054	195	65
CONTIG472	22283442_c2_149	5393	11055	1359	453
CONTIG472	10320382_c2_151	5394	11056	2247	749
CONTIG472	24491266_c3_177	5395	11057	3009	1003
CONTIG473	116287_f1_5	5396	11058	1971	657
CONTIG473	21742211_f3_70	5397	11059	1236	412
CONTIG473	3400141_c2_149	5398	11060	567	189
CONTIG475	12527018_f1_33	5399	11061	195	65
CONTIG475	10955042_c2_180	5400	11062	249	83
CONTIG475	14661680_c3_220	5401	11063	201	67
CONTIG476	35646916_f2_49	5402	11064	183	61
CONTIG476	25634437_c2_169	5403	11065	216	72
CONTIG477	32116557_f1_24	5404	11066	282	94
CONTIG477	2931552_c1_143	5405	11067	384	128
CONTIG477	32617003_c1_144	5406	11068	183	61
CONTIG477	4144552_c1_162	5407	11069	228	76
CONTIG477	22772687_c1_169	5408	11070	186	62
CONTIG477	4787518_c2_187	5409	11071	336	112
CONTIG478	12697175_f2_69	5410	11072	429	143
CONTIG478	4352253_f3_109	5411	11073	345	115
CONTIG478	1058452_f3_140	5412	11074	234	78
CONTIG478	4551262_c2_204	5413	11075	675	225
CONTIG479	5136015_f1_1	5414	11076	249	83
CONTIG479	23706377_c1_147	5415	11077	186	62
CONTIG479	2538508_c2_193	5416	11078	1002	334
CONTIG479	16284462_c3_238	5417	11079	186	62
CONTIG479	31410277_c3_252	5418	11080	234	78
CONTIG480	34261542_c1_168	5419	11081	1395	465

CONTIG480	19581591_c2_205	5420	11082	2637	879						
CONTIG480	3923507_c3_234	5421	11083	219	73						
CONTIG482	32032765_fl_44	5422	11084	204	68						
CONTIG482	34178455_f3_122	5423	11085	783	261						
CONTIG482	14073293_c2_195	5424	11086	351	117						
CONTIG483	11738212_fl_15	5425	11087	240	80						
CONTIG483	7157901_f3_87	5426	11088	219	73						
CONTIG483	22910842_c1_140	5427	11089	237	79						
CONTIG485	22911318_fl_17	5428	11090	195	65						
CONTIG485	11973425_f2_92	5429	11091	285	95						
CONTIG485	14535938_f3_125	5430	11092	381	127						
CONTIG485	1307942_c1_151	5431	11093	741	247						
CONTIG485	36447656_c1_153	5432	11094	549	183						
CONTIG485	19540843_c1_155	5433	11095	699	233						
CONTIG485	31439827_c1_158	5434	11096	636	212						
CONTIG485	21756332_c1_165	5435	11097	411	137						
CONTIG485	14969588_c1_166	5436	11098	369	123						
CONTIG485	22678752_c1_171	5437	11099	474	158						
CONTIG485	12120966_c1_172	5438	11100	195	65						
CONTIG485	11772250_c1_173	5439	11101	489	163						
CONTIG485	9970466_c1_185	5440	11102	327	109						
CONTIG485	24886586_c2_189	5441	11103	570	190						
CONTIG485	22830405_c2_193	5442	11104	270	90						
CONTIG485	25494591_c2_196	5443	11105	765	255						
CONTIG485	11192582_c2_202	5444	11106	783	261						
CONTIG485	36463300_c2_204	5445	11107	264	88						
CONTIG485	6852282_c2_207	5446	11108	978	326						
CONTIG485	15863536_c2_210	5447	11109	2946	982						
CONTIG485	10960457_c2_211	5448	11110	633	211						
CONTIG485	6067678_c2_218	5449	11111	1557	519						
CONTIG485	32595657_c3_233	5450	11112	786	262						
CONTIG485	5292657_c3_235	5451	11113	540	180						
CONTIG485	12971931_c3_236	5452	11114	537	179						
CONTIG485	9895406_c3_243	5453	11115	405	135						
CONTIG485	667336_c3_245	5454	11116	672	224						
CONTIG485	29782891_c3_249	5455	11117	702	234						
CONTIG485	30746081_c3_256	5456	11118	1008	336						
CONTIG485	9771937_c3_261	5457	11119	231	77						
CONTIG485	33364217_c3_262	5458	11120	204	68						

[illegible]

[illegible]

[illegible]



[illegible]

CONTIG512	24303790_c1_293	5615	11277	201	67						
CONTIG512	4102090_c1_317	5616	11278	951	317						
CONTIG512	4479193_c2_427	5617	11279	981	327						
CONTIG512	26219653_c3_443	5618	11280	183	61						
CONTIG512	24491512_c3_514	5619	11281	1257	419						
CONTIG513	11761717_f1_65	5620	11282	381	127						
CONTIG513	36516877_f1_72	5621	11283	189	63						
CONTIG513	16494757_f1_107	5622	11284	564	188						
CONTIG513	10644651_f2_173	5623	11285	183	61						
CONTIG513	13884401_f2_189	5624	11286	186	62						
CONTIG513	13066307_f2_226	5625	11287	294	98						
CONTIG513	22663931_f3_260	5626	11288	543	181						
CONTIG513	165802_f3_269	5627	11289	186	62						
CONTIG513	14337781_f3_378	5628	11290	438	146						
CONTIG513	22057040_f3_384	5629	11291	528	176						
CONTIG513	12991255_c1_424	5630	11292	468	156						
CONTIG513	6744762_c1_429	5631	11293	279	93						
CONTIG513	25496067_c1_435	5632	11294	765	255						
CONTIG513	29718808_c1_465	5633	11295	660	220						
CONTIG513	31517316_c1_531	5634	11296	258	86						
CONTIG513	14181925_c2_552	5635	11297	573	191						
CONTIG513	24413961_c2_627	5636	11298	198	66						
CONTIG513	1354152_c3_686	5637	11299	1131	377						
CONTIG513	16189443_c3_700	5638	11300	210	70						
CONTIG513	22839138_c3_747	5639	11301	210	70						
CONTIG513	972812_c3_800	5640	11302	330	110						
CONTIG52	29502261_c3_8	5641	11303	219	73						
CONTIG55	11820252_f2_1	5642	11304	231	77						
CONTIG56	21756280_f2_1	5643	11305	462	154						
CONTIG57	197135_c1_2	5644	11306	258	86						
CONTIG58	2469800_f2_4	5645	11307	297	99						
CONTIG60	4178160_c2_4	5646	11308	339	113						
CONTIG68	6922902_f2_1	5647	11309	342	114						
CONTIG68	34551450_f3_2	5648	11310	363	121						
CONTIG69	14460950_c2_5	5649	11311	663	221						
CONTIG69	19553_c3_6	5650	11312	633	211						
CONTIG75	23553182_c1_2	5651	11313	237	79						
CONTIG76	17007692_f2_1	5652	11314	363	121						
CONTIG77	32213278_c1_5	5653	11315	195	65						

CONTIG79	10976406_c3_4	5654	11316	609	203				
CONTIG8	2634382_fl_2	5655	11317	220	74				
CONTIG81	14104500_c3_4	5656	11318	855	285				
CONTIG82	3252313_cl_2	5657	11319	192	64				
CONTIG89	34272811_fl_1	5658	11320	216	72				
CONTIG90	31800306_f2_2	5659	11321	297	99				
CONTIG95	406661_c3_5	5660	11322	237	79				
CONTIG97	5995312_fl_1	5661	11323	210	70				
CONTIG97	836_c3_2	5662	11324	579	193				